

# Hopf Galois (Co)Extensions In Noncommutative Geometry

Mohammad HASSANZADEH



Institut des Hautes Études Scientifiques  
35, route de Chartres  
91440 – Bures-sur-Yvette (France)

Avril 2012

IHES/M/12/06

# HOPF GALOIS (CO)EXTENSIONS IN NONCOMMUTATIVE GEOMETRY

MOHAMMAD HASSANZADEH

ABSTRACT. We introduce an alternative proof, with the use of tools and notions for Hopf algebras, to show that Hopf Galois coextensions of coalgebras are the sources of stable anti Yetter-Drinfeld modules. Furthermore we show that two natural cohomology theories related to a Hopf Galois coextension are isomorphic.

## INTRODUCTION

The general definition of Hopf Galois extensions was introduced by Kreimer and Takeuchi [KT] around the same time when noncommutative geometry [C-Book] started to develop with works of Alain Connes in 1980. In fact Hopf Galois extensions are noncommutative analogue of affine torsors and principal bundles. Descent data related to a Hopf Galois extension has its roots in works of Grothendieck in algebraic geometry [Groth-SGA1]. Basically there are two well-known homology theories which are related to a Hopf Galois extension  $A(B)_H$ . One is the relative cyclic homology of the algebra extension  $B \subseteq A$ , and another one is the Hopf cyclic homology of the Hopf algebra  $H$  involved in the Hopf Galois extension. It is shown in [J-S] that these two homology theories are isomorphic where the Hopf cyclic homology of Hopf algebra  $H$  has coefficients in a stable anti Yetter-Drinfeld (SAYD) module. Not only this isomorphism is given by the canonical isomorphism of the Hopf Galois extension, but also the SAYD coefficients, which is a module-comodule over  $H$ , is constructed by this canonical map. A great idea here is that the cyclic (co)homology of (co)algebra (co)extensions which is not easy to compute is isomorphic to the Hopf cyclic (co)homology of the Hopf algebra of the (co)extension which can be computed more easily. The dual notion of Hopf Galois coextensions is introduced by Schneider in [Schn1] and can be viewed as a noncommutative generalization of the theory of quotients of formal schemes under free actions of formal group schemes [Schhn2].

In this paper we recall the basics of Hopf Galois (co)extensions. Next, we study the module-comodule and specially stable anti Yetter-Drinfeld module structures over the Hopf algebra involved in a Hopf Galois (co)extension. Furthermore we study the related (co)homology theories to a Hopf Galois (co)extension and finally we observe that these (co)homology theories are isomorphic. More precisely, in Section 1 we recall the results of Jara and Stefan in [J-S] for Hopf Galois extensions and in Section 2 we introduce an alternative proof for the dual case of Hopf Galois coextensions using the tools and notions for Hopf algebras. Although the main results of the second section of this paper have been already proved for a general case of Equivariant Hopf Galois coextensions for  $\times$ -Hopf coalgebras in [HR2], a direct

proof for the case of Hopf algebras using the related concepts and notions, which has not been appeared in the literature, has its own advantages.

**Acknowledgments:** The author would like to thank the Institut des Hautes Études Scientifiques, IHES, for its hospitality and financial supports during his visit when the whole work was accomplished.

**Notations:** In this paper we denote a Hopf algebra by  $H$  and its counit by  $\varepsilon$ . We use the Sweedler summation notation  $\Delta(h) = h^{(1)} \otimes h^{(2)}$  for the coproduct of a Hopf algebra. Furthermore the summation notations  $\blacktriangleright(h) = h_{<-1>} \otimes h_{<0>}$  and  $\blacktriangleleft(h) = h_{<0>} \otimes h_{<1>}$  are used for the left and right coactions of a coalgebra, respectively.

## CONTENTS

Introduction	1
1. Hopf cyclic cohomology	2
1.1. Cyclic modules and cyclic duality	2
1.2. Cyclic cohomology of Hopf algebras	3
2. Hopf Galois extensions of algebras	4
2.1. Preliminaries of Hopf Galois extensions	4
2.2. Homology theories related to Hopf Galois extensions	6
3. Hopf Galois coextensions of coalgebras	8
3.1. Preliminaries of Hopf Galois coextensions	9
3.2. Hopf Galois coextensions as sources of anti-Yetter-Drinfeld modules	10
3.3. Cohomology theories related to Hopf Galois coextensions	16
References	20

## 1. HOPF CYCLIC COHOMOLOGY

In this section we review the basics of (co)cyclic modules. We explain cyclic duality to obtain a cyclic module from a cocyclic module and vice versa. As an example we study the Hopf cyclic (co)homology with coefficients in a stable anti Yetter-Drinfeld module.

**1.1. Cyclic modules and cyclic duality.** A cosimplicial module [C-Book], [Lo] contains  $\mathbb{C}$ -modules  $C^n$ ,  $n \geq 0$ , with  $\mathbb{C}$ -module maps  $\delta_i : C^n \rightarrow C^{n+1}$  called cofaces, and  $\sigma_i : C^n \rightarrow C^{n-1}$  called codegeneracies satisfying the following cosimplicial relations;

$$\begin{aligned}
 \delta_j \delta_i &= \delta_i \delta_{j-1}, & \text{if } i < j, \\
 \sigma_j \sigma_i &= \sigma_i \sigma_{j+1}, & \text{if } i \leq j, \\
 \sigma_j \delta_i &= \begin{cases} \delta_i \sigma_{j-1}, & \text{if } i < j, \\ \text{Id}, & \text{if } i = j \text{ or } i = j + 1, \\ \delta_{i-1} \sigma_j, & \text{if } i > j + 1. \end{cases} & (1.1)
 \end{aligned}$$

A cocyclic module is a cosimplicial module with extra morphisms  $\tau : C^n \rightarrow C^n$  which are called cocyclic maps such that the following extra commutativity relations hold.

$$\begin{aligned} \tau\delta_i &= \delta_{i-1}\tau, & 1 \leq i \leq n+1, \\ \tau\delta_0 &= \delta_{n+1}, & \tau\sigma_i = \sigma_{i-1}\tau, & 1 \leq i \leq n, \\ \tau\sigma_0 &= \sigma_n\tau^2, & \tau^{n+1} &= \text{Id}. \end{aligned} \quad (1.2)$$

Dually a cyclic module is given by quadruple  $C = (C_n, \delta_i, \sigma_i, \tau_n)$  where  $C_n$ 's,  $n \geq 0$ , are  $\mathbb{C}$ -modules and there are  $\mathbb{C}$ -module maps  $\delta_i : C_n \rightarrow C_{n-1}$ , called faces,  $\sigma_i : C_n \rightarrow C_{n+1}$ ,  $0 \leq i \leq n$  called degeneracies and  $\tau : C_n \rightarrow C_n$  called cyclic maps satisfying the following commutativity relations;

$$\begin{aligned} \delta_i\delta_j &= \delta_{j-1}\delta_i, & \text{if } i < j, \\ \sigma_i\sigma_j &= \sigma_{j+1}\sigma_i, & \text{if } i \leq j, \\ \delta_i\sigma_j &= \begin{cases} \sigma_{j-1}\delta_i, & \text{if } i < j, \\ \text{Id}, & \text{if } i = j \text{ or } i = j+1, \\ \sigma_j\delta_{i-1}, & \text{if } i > j+1, \end{cases} \end{aligned} \quad (1.3)$$

and;

$$\begin{aligned} \delta_i\tau &= \tau\delta_{i-1}, & 1 \leq i \leq n, \\ \delta_0\tau &= \delta_n, & \sigma_i\tau = \tau\sigma_{i-1}, & 1 \leq i \leq n, \\ \sigma_0\tau &= \tau^2\sigma_n, & \tau^{n+1} &= \text{Id}. \end{aligned} \quad (1.4)$$

Now we recall the duality procedure for (co)cyclic modules [C-Book]. For any cyclic module  $C = (C_n, \delta_i, \sigma_i, \tau)$  we can construct its cyclic dual which is denoted by  $\check{C}$  and defined by  $\check{C}^n = C_n$  with the following cofaces, codegeneracies and cyclic morphisms.

$$\begin{aligned} d_0 &:= \tau_n\sigma_{n-1}, & d_i &:= \sigma_{i-1} : \check{C}^n \rightarrow \check{C}^{n+1}, & 1 \leq i \leq n, \\ s_i &:= \delta_i : \check{C}^n \rightarrow \check{C}^{n-1}, & 0 \leq i \leq n-1, \\ t &:= \tau^{-1}. \end{aligned} \quad (1.5)$$

Conversely for any cocyclic module  $C = (C^n, d_i, s_i, t)$  one has its cyclic dual denoted by  $\tilde{C}$  where;

$$\begin{aligned} \delta_i &:= s_i : \tilde{C}_n \rightarrow \tilde{C}_{n-1}, & 0 \leq i \leq n-1, & \delta_n := \delta_0\tau_n, \\ \sigma_i &:= d_{i+1} : \tilde{C}_n \rightarrow \tilde{C}_{n+1}, & 0 \leq i \leq n-1, \\ \tau &:= t^{-1}. \end{aligned} \quad (1.6)$$

**1.2. Cyclic cohomology of Hopf algebras.** In this subsection, we study some examples of (co)cyclic modules and their dual theories for Hopf algebras which will be used later in Sections 2 and 3. Hopf cyclic cohomology was introduced by Connes and Moscovici in [CM98] and was generalized to Hopf cyclic cohomology with coefficients in a stable anti Yetter-Drinfeld module in [HKRS2].

A left-right stable anti Yetter-Drinfeld (SAYD) module  $M$  over a Hopf algebra  $H$  is a left module and a right comodule over  $H$  satisfying the following

compatibility condition [HKRS1];

$$(h \triangleright m)_{\langle 0 \rangle} \otimes (h \triangleright m)_{\langle 1 \rangle} = h^{(2)} \triangleright m_{\langle 0 \rangle} \otimes h^{(3)} m_{\langle 1 \rangle} S(h^{(1)}). \quad (1.7)$$

We use this type of SAYD modules in Section 2 for Hopf Galois extensions. Similarly a right-left SAYD module  $M$  over  $H$  is a right module and a left comodule satisfying the following compatibility condition,

$$(m \triangleleft h)_{\langle -1 \rangle} \otimes (m \triangleleft h)_{\langle 0 \rangle} = S(h^{(3)}) m_{\langle -1 \rangle} h^{(1)} \otimes m_{\langle 0 \rangle} \triangleleft h^{(2)}. \quad (1.8)$$

This type of SAYD modules will be used in Section 3 for Hopf Galois coextensions. A generalization of Connes-Moscovici cocyclic module for Hopf algebras with coefficients in a SAYD module was introduced in [HKRS2]. The following cyclic module is the dual cyclic module of the mentioned cocyclic module for a left-right  $H$ -SAYD module  $M$  and  $C_n(H, M) = H^{\otimes n+1} \otimes_H M$ .

$$\begin{aligned} \delta_i(h_0 \otimes \cdots \otimes h_n \otimes_H m) &= h_0 \otimes \cdots \otimes \varepsilon(h_i) \otimes \cdots \otimes h_n \otimes_H m, \\ \sigma_i(h_0 \otimes \cdots \otimes h_n \otimes_H m) &= h_0 \otimes \cdots \otimes \Delta(h_i) \otimes \cdots \otimes h_n \otimes_H m, \\ \tau_n(h_0 \otimes \cdots \otimes h_n \otimes_H m) &= h_n m_{\langle 1 \rangle} \otimes h_0 \otimes \cdots \otimes h_{n-1} \otimes_H m_{\langle 0 \rangle}. \end{aligned} \quad (1.9)$$

We use this cyclic module in Section 2 for Hopf Galois extensions. The authors in [KR02] introduced a new cyclic module for Hopf algebras which is a dual for Connes-Moscovici cocyclic module for Hopf algebras in some sense. Later they have shown in [KR03] that this cyclic module is isomorphic to the cyclic dual of Connes-Moscovici cocyclic module for Hopf algebras. A generalization of this cyclic module with coefficients in a SAYD module was introduced in [HKRS2]. The following cocyclic module is the dual cocyclic module of the mentioned cyclic module for a right-left SAYD module  $M$  over Hopf algebra  $H$  and  $C^n(H, M) = H^{\otimes n} \otimes M$ .

$$\begin{aligned} \delta_i(\tilde{h} \otimes m) &= h_1 \otimes \cdots \otimes h_i \otimes 1_H \otimes \cdots \otimes h_n \otimes m, \\ \delta_n(\tilde{b} \otimes m) &= h_1^{(1)} \otimes \cdots \otimes h_n^{(1)} \otimes S(h_1^{(2)} \cdots h_n^{(2)}) m_{\langle -1 \rangle} \otimes m_{\langle 0 \rangle}, \\ \sigma_i(\tilde{h} \otimes m) &= h_1 \otimes \cdots \otimes h_i h_{i+1} \otimes \cdots \otimes h_n \otimes m, \\ \sigma_n(\tilde{h} \otimes m) &= h_1 \otimes \cdots \otimes h_{n-1} \varepsilon(h_n) \otimes m, \\ \tau_n(\tilde{h} \otimes m) &= h_2^{(1)} \otimes \cdots \otimes h_n^{(1)} \otimes S(h_1^{(2)} \cdots h_n^{(2)}) m_{\langle -1 \rangle} \otimes m_{\langle 0 \rangle} \triangleleft h_1^{(1)}. \end{aligned} \quad (1.10)$$

Here  $\tilde{h} = h_1 \otimes \cdots \otimes h_n$ . We use this cocyclic module in Section 3 for Hopf Galois coextensions. For more about Hopf cyclic (co)homology we refer the reader to [CM98], [HKRS2], [Kh] and [Kay].

## 2. HOPF GALOIS EXTENSIONS OF ALGEBRAS

**2.1. Preliminaries of Hopf Galois extensions.** In this section we recall the notion of Hopf Galois extensions. We review the results in [J-S] which imply that Hopf Galois extensions are sources of producing stable anti Yetter-Drinfeld modules. Furthermore we study the relation between the relative cyclic homology of the algebra extension and the Hopf cyclic homology of the Hopf algebra involved in a Hopf Galois extension.

Let  $H$  be a Hopf algebra and  $A$  a right  $H$ -comodule with the coaction  $\rho : A \longrightarrow A \otimes H$ . The set of coinvariants of this coaction  $B = \{a \in A, \rho(a) = a \otimes 1_H\}$  is a subalgebra of  $A$ . The algebra extension  $B \subseteq A$  is Hopf Galois if the canonical map

$$\beta : A \otimes_B A \longrightarrow A \otimes H, \quad a \otimes_B a' \longmapsto aa'_{\langle 0 \rangle} \otimes a'_{\langle 1 \rangle}, \quad (2.1)$$

is bijective. We denote such a Hopf Galois extension by  $A(B)^H$ .

The map  $\beta$  is an isomorphism of left  $A$ -modules and right  $H$ -comodules where the left  $A$ -module structures of  $A \otimes_B A$  and  $A \otimes H$  are given by  $a(a_1 \otimes_B a_2) = aa_1 \otimes_B a_2$  and  $a(a' \otimes h) = aa' \otimes h$ , respectively, and also the right  $H$  comodule structures of  $A \otimes_B A$  and  $A \otimes H$  are given by  $a \otimes a' \longmapsto a \otimes a'_{\langle 0 \rangle} \otimes a'_{\langle 1 \rangle}$  and  $a \otimes h \longmapsto a \otimes h^{(1)} \otimes h^{(2)}$ , respectively, for all  $a, a', a_1, a_2 \in A$  and  $h \in H$ . In fact an extension  $B \subseteq A$  is Hopf Galois if  $A \otimes_B A$  and  $A \otimes H$  are isomorphic in  ${}_A\mathcal{M}^H$ , the category of left  $A$ -modules and right  $H$ -comodules, by the canonical map  $\beta$ . The first natural question is why we call this extension Galois and what is its relation with classical Galois extension of fields.

**Example 2.1.** *Classical Galois extensions of fields:*

Let  $F \subseteq E$  be a field extension with the Galois group  $G$ . This extension is Galois if and only if  $G$  acts faithfully on  $E$ . This is equivalent to  $|[E : F]| = |G|$ . Let  $|G| = n$  and  $G = \{x_1, \dots, x_n\}$ . Furthermore suppose  $\{b_1, \dots, b_n\}$  be the basis of  $E/F$  and  $\{p^1, \dots, p^n\}$  be the dual basis of  $\{x_i\}_i$  in  $(kG)^*$  where  $kG$  is the group algebra of  $G$ . Since  $G$  is finite then  $(kG)^*$  is a Hopf algebra and  $p^n(x_i) = \delta_{ij}$ . Furthermore the action of  $G$  and therefore  $kG$  on  $E$  amounts to a coaction of  $(kG)^*$  on  $E$  as follows;

$$E \longrightarrow E \otimes (kG)^*, \quad a \longmapsto x_i \triangleright a \otimes p^i. \quad (2.2)$$

One checks that  $E$  is a right  $(kG)^*$ -comodule algebra by this coaction. We define the canonical Galois map as follows;

$$\beta : E \otimes_F E \longrightarrow E \otimes_k (kG)^*, \quad a \otimes b \longmapsto a(x_i \triangleright b) \otimes p^i. \quad (2.3)$$

One use the independence of  $\{p^i\}$  to show that  $\beta$  is injective. Furthermore since both tensor products are finite-dimensional  $F$ -algebras then  $\beta$  is a surjection.

**Example 2.2.** A Hopf algebra  $H$  over the field  $F$  is a right  $H$ -comodule algebra where the right coaction is given by the comultiplication of  $H$ . For  $A = H$  we have  $B = F1_H$ . Therefore  $H(F)_H$  is a  $H$ -Galois extension with the canonical map which is given by;

$$\beta : H \otimes_F H \longrightarrow H \otimes_F H, \quad h \otimes k \longmapsto hk^{(1)} \otimes k^{(2)}, \quad h, k \in H, \quad (2.4)$$

with the inverse map  $\beta^{-1}(h \otimes k) = hS(k^{(1)}) \otimes k^{(2)}$ .

The following example shows that Hopf Galois extensions are algebraic analogue of principal bundles.

**Example 2.3.** *Principal bundles:*

Let  $P(M, G)$  denote the principal bundle of the smooth manifold  $P$  over the

base  $M$  with the structure Lie group  $G$ . There is a smooth right action of  $G$  on  $P$  denoted by  $\triangleleft : P \times G \rightarrow P$  which is free, *i.e.*

$$u \triangleleft g = u \triangleleft g' \implies g = g', \quad g, g' \in G, u \in P. \quad (2.5)$$

If  $G$  is finite then freeness of the  $G$ -action is equivalent to the injectivity of the following map;

$$\beta : P \times G \rightarrow P \times P, \quad (u, g) \mapsto (u, u \triangleleft g). \quad (2.6)$$

One notes that the orbit space is isomorphic to the base space  $P/G \cong M$  and the canonical projection  $\pi : P \rightarrow M$  is smooth. Now we apply a noncommutative approach to the map  $\beta$  and we consider the space of functions on the principal bundle. Let  $A = C^\infty(P)$ ,  $B = C^\infty(M)$  and  $H = (kG)^*$ . We dualize the action  $\triangleleft$  to obtain a right coaction  $\blacktriangledown : A \rightarrow A \otimes (kG)^*$ . Therefore the following induced map

$$(m_A \otimes \text{Id}) \circ (\text{Id} \otimes \blacktriangledown) : A \otimes A \rightarrow A \otimes H, \quad (2.7)$$

is surjective where  $m_A$  denotes the multiplication of  $A$ . Now if we restrict the tensor product in the domain of the previous map to the coinvariant space of the coaction  $\blacktriangledown$ , we obtain an isomorphism. This example is a motivation to define the notion of quantum principal bundles. A right  $H$ -comodule algebra  $A$  is called a quantum principal bundle if the Galois map  $\beta$  related to the coaction of  $H$  is an isomorphism. For more about this example we refer the reader to [Haj], [Dur], [BM] and [KZ].

**Remark 2.4.** There are two major difference between Hopf Galois theory and the classical Galois theory of field extensions. Classical Galois extensions can be characterized by the normal, separable field extensions without explicitly mentioning the Galois group. Furthermore, a Galois field extension determines uniquely the Galois group. There is no similar result for Hopf Galois theory, although a characterization exists for Hopf Galois extensions of Hopf algebroids with some finiteness conditions. Another difference is that the fundamental theorem of Galois theory of fields extensions does not hold for the Hopf Galois extensions of algebras. We refer the reader for more in this regard to [Bál-Thesis] and [Ver].

Hopf Galois extensions were generalized later in different ways. For example the authors in [BH] have introduced the notion of coalgebra extensions. Also see [SS]. Furthermore, Galois extensions have been studied for extended versions of Hopf algebras such as Hopf algebroids in [BO] and  $\times$ -Hopf algebras in [BS] and [HR].

**2.2. Homology theories related to Hopf Galois extensions.** Basically there are two well-known homology theories which are related to a Hopf Galois extension  $A(B)_H$ . One is the relative cyclic homology of the algebra extension  $B \subseteq A$  and another one is the Hopf cyclic homology of the Hopf algebra  $H$  involved in the Hopf Galois extension. In this subsection we recall that these two homology theories are isomorphic. Here the Hopf cyclic homology of  $H$  has coefficients in a SAYD module which is constructed by the canonical isomorphism of the Hopf Galois extension. This shows that Hopf Galois extensions are the sources of providing SAYD modules. In this subsection we recall some related results from [J-S] to be able to compare

them with the similar results in the dual case of Hopf Galois coextensions in Section 3.

Suppose  $B$  be an associative algebra over the field of complex numbers. Let  $M$  and  $N$  be  $B$ -bimodules. The cyclic tensor product of  $M$  and  $N$  is defined to be  $M\widehat{\otimes}_B N := (M \otimes_B N) \otimes_{B^e} B, [Q]$ . It can be shown that

$$M\widehat{\otimes}_B N \cong \frac{M \otimes_B N}{[M \otimes_B N, B]}, \quad (2.8)$$

where the bracket stands for the subspace generated by all commutators. In fact,

$$M\widehat{\otimes}_B N = \frac{M \otimes_B N}{\sim}, \quad (2.9)$$

where  $\sim$  is the equivalence relation defined by the following relation;

$$\{bm \otimes_B n = m \otimes_B nb\}. \quad (2.10)$$

Similarly if  $M_1, \dots, M_n$  are  $B$ -bimodules then their cyclic tensor product can be defined as

$$M_1\widehat{\otimes} \dots \widehat{\otimes} M_n := (M_1 \otimes_B \dots \otimes_B M_n) \otimes_{B^e} B. \quad (2.11)$$

Now let  $B \subseteq A$  be an algebra extension and  $M$  a  $B$ -bimodule. Let  $C_n(A(B), M) = M\widehat{\otimes}_B A^{\widehat{\otimes} B}$ . One defines a simplicial module where the faces are given by

$$d_i(m\widehat{\otimes}_B a_1\widehat{\otimes}_B \dots \widehat{\otimes}_B a_n) = \begin{cases} ma_1\widehat{\otimes}_B a_2\widehat{\otimes}_B \dots \widehat{\otimes}_B a_n, & i = 0, \\ m\widehat{\otimes}_B a_1\widehat{\otimes} \dots \widehat{\otimes}_B a_i a_{i+1}\widehat{\otimes}_B \dots \widehat{\otimes}_B a_n, & 0 < i < n, \\ a_n m\widehat{\otimes}_B a_1\widehat{\otimes}_B \dots \widehat{\otimes}_B a_{n-1}, & i = n, \end{cases} \quad (2.12)$$

and the degeneracies are given by

$$s_i(m\widehat{\otimes}_B a_1\widehat{\otimes}_B \dots \widehat{\otimes}_B a_n) = m\widehat{\otimes}_B a_1\widehat{\otimes}_B \dots \widehat{\otimes}_B 1_A \widehat{\otimes}_B \dots \widehat{\otimes}_B a_n. \quad (2.13)$$

It is shown in [J-S] that if  $M = A$ , then the following cyclic operator turns  $C_n(A(B), A)$  into a cyclic module;

$$t_n(a_0\widehat{\otimes}_B a_1\widehat{\otimes}_B \dots \widehat{\otimes}_B a_n) = a_n\widehat{\otimes}_B a_0\widehat{\otimes}_B \dots \widehat{\otimes}_B a_{n-1}. \quad (2.14)$$

The cyclic homology of this cyclic module is called relative cyclic homology of the algebra extension  $B \subseteq A$  and it is denoted by  $HC_*(A(B), A)$ . Although the space  $A \otimes_B A$  has not a well-defined algebra structure, the subspace  $(A \otimes_B A)^B$  is an associative algebra by the following multiplication;

$$(a_1 \otimes_B a'_1)(a_2 \otimes_B a'_2) = a_1 a_2 \otimes_B a'_2 a'_1. \quad (2.15)$$

The canonical map  $\beta$  induces an isomorphism;

$$\bar{\beta}: (A \otimes_B A)^B \longrightarrow A^B \otimes H.$$

Therefore in spite of the fact that the canonical isomorphism of the Hopf Galois extension is not an algebra map, the algebra structure (2.15) enables us to obtain the following anti-algebra map;

$$\kappa: H \longrightarrow (A \otimes_B A)^B, \quad \kappa := \bar{\beta}^{-1} \circ i, \quad (2.16)$$



where  $i : H \longrightarrow A^B \otimes H$  is given by  $h \longmapsto 1_A \otimes h$ . We denote the summation notation  $\kappa(h) = \kappa^1(h) \otimes \kappa^2(h)$  for the image of the map  $\kappa$ . Now let  $M$  be a  $A$ -bimodule. We set;

$$M^B = \{m \in M, \quad bm = mb, \quad \forall b \in B\}, \quad M_B = \frac{M}{[M, B]}. \quad (2.17)$$

The map  $\kappa$  enables us to define a right  $H$ -module structure on  $M^B$  given by

$$mh = \kappa^1(h)m\kappa^2(h). \quad (2.18)$$

Furthermore a left  $H$ -module structure on  $M_B$  can be defined by

$$hm = \kappa^1(h)m\kappa^1(h). \quad (2.19)$$

In a special case when  $M = A$ , the quotient space  $A_B$  is also a right  $H$ -comodule by the original coaction of  $H$  over  $A$ . This coaction with the action defined in (2.19) amounts  $A_B$  to a left-right SAYD module over  $H$ . For any Hopf Galois extension  $A(B)_H$ , one iteratively use the map  $\beta$  to transfer the cyclic structure of  $C_n(A(B), A)$  to  $C_n(H, A_B) = H^{\otimes n} \otimes_H H \otimes A_B$ . The result cyclic module on  $C_n(H, A_B)$  is proved in [J-S] to be the cyclic module associated to Hopf cyclic homology of Hopf algebra  $H$  with coefficients in SAYD module  $A_B$  which is introduced in (1.9). The results of this subsection have been later generalized for Hopf Galois extensions of  $\times$ -Hopf algebras in [BS] and the Equivariant Hopf Galois extensions of  $\times$ -Hopf algebras in [HR]. We summarize headlines of this section to be able to compare them with the dual case of Hopf Galois coextensions in the sequel section.

**Remark 2.5.** For any Hopf Galois extension  $A(B)^H$ , we have;

- i) The relative cyclic module of the algebra extension is a quotient space given by,

$$B \otimes_{B^e} A \otimes_B \otimes_B \cdots \otimes_B A = A \widehat{\otimes}_B \cdots \widehat{\otimes}_B A = \frac{A \otimes_B \cdots \otimes_B A}{\sim}.$$

- ii) The objects  $A \otimes_B A$  and  $A \otimes H$  are isomorphic in the category of  ${}_A\mathcal{M}^H$ .
- iii) The subspace  $A^B$  is an subalgebra of  $A$ .
- iv) The subspace  $(A \otimes_B A)^B$  is an algebra.
- v) The subspace  $A^B$  is a left  $(A \otimes_B A)^B$ -module and a right  $H$ -module.
- vi) The quotient space  $A_B$  is a right  $(A \otimes_B A)^B$ -module and a left  $H$ -module.
- vii) The quotient space  $A_B \cong B \otimes_{B^e} A$  is a right-left SAYD module over  $H$ .
- viii)  $HC_*(H, A_B) \cong HC_*(A(B), A)$ .

### 3. HOPF GALOIS COEXTENSIONS OF COALGEBRAS

In this section we use the notions and tools for Hopf algebras and Hopf cyclic cohomology to prove that Hopf Galois coextensions of coalgebras are the sources of stable anti Yetter-Drinfeld modules. Furthermore we show that the Hopf cyclic cohomology of the Hopf algebra involved in a Hopf Galois coextension with coefficients in a SAYD module, which is dual to  $A_B$ , is isomorphic to the relative cyclic cohomology of coalgebra coextension.

These are the dual results in [J-S] for Hopf Galois extensions which are explained in Subsection 2.2. The authors in [HR2] have proved these results for  $\times$ -Hopf coalgebras and have mentioned that the similar results for Hopf Galois coextensions hold as a special case of their work.

**3.1. Preliminaries of Hopf Galois coextensions.** Let  $H$  be a Hopf algebra and  $C$  a right  $H$ -module coalgebra with the action  $\triangleleft : C \otimes H \rightarrow C$ . The set

$$I = \{c \triangleleft h - \varepsilon(h)c\}, \quad (3.1)$$

is a two-sided coideal of  $C$  and therefore  $D = \frac{C}{I}$  is a coalgebra. The natural surjection  $\pi : C \rightarrow D$  is a right  $H$ -Galois coextension if the canonical map

$$\beta : C \otimes H \rightarrow C \square_D C, \quad c \otimes h \mapsto c^{(1)} \otimes c^{(2)} \triangleleft h, \quad (3.2)$$

is a bijection. Such a Hopf Galois coextensions is denoted by  $C(D)^H$ . Here the  $D$ -bicomodule structure of  $C$  is given by

$$c \mapsto \pi(c^{(1)}) \otimes c^{(2)}, \quad c \mapsto c^{(1)} \otimes \pi(c^{(2)}). \quad (3.3)$$

Since by the definition of  $D$  for all  $h \in H$  and  $c \in C$  we have

$$\pi(c \triangleleft h) = \varepsilon(h)\pi(c), \quad (3.4)$$

the map  $\beta$  defined in (3.2) is well-defined. Let us denote the inverse of Galois map  $\beta$  by the following summation notation;

$$\beta^{-1}(c \square_D c') := \beta_-(c \square_D c') \otimes \beta_+(c \square_D c'), \quad (3.5)$$

where  $Im\beta_- \in C$  and  $Im\beta_+ \in H$ . If there is no confusion we can simply write  $\beta = \beta_- \otimes \beta_+$ . Since  $C \square_D C$  is not a coalgebra, the map  $\beta$  is not an isomorphism of coalgebras. Instead it is an isomorphism of left  $C$ -comodules and right  $H$ -modules. The map  $\beta$  is a left  $C$ -comodule map where the left  $C$ -comodule structures of  $C \square_D C$  and  $C \otimes H$  are given by

$$c \square_D c' \mapsto c^{(1)} \otimes c^{(2)} \square_D c', \quad c \otimes h \mapsto c^{(1)} \otimes c^{(2)} \otimes h.$$

Furthermore it is a right  $H$ -module map where the right  $H$  module structures of  $C \square_D C$  and  $C \otimes H$  are given by

$$(c \square_D c') \triangleleft h = c \square_D c' \triangleleft h, \quad (c \otimes h) \triangleleft h' = c \otimes hh'.$$

The first  $H$ -action introduced above is well-defined by (3.4). In fact the coextension  $\pi : C \rightarrow D$  is Hopf Galois if  $C \otimes H$  and  $C \square_D C$  are isomorphic in  ${}^C\mathcal{M}_H$ , the category of left  $C$ -comodules and right  $H$ -modules, by the canonical map  $\beta$ . One notes that the categories  ${}^C\mathcal{M}_H$  and  ${}_A\mathcal{M}^H$  related to Hopf Galois extensions and coextensions are dual to each other.

**Lemma 3.1.** *Let  $C$  and  $D$  be coalgebras,  $H$  a Hopf algebra,  $C(D)^H$  a Hopf Galois coextension with canonical bijection  $\beta$ . Then the following properties hold.*

- i)  $\beta_-^{(1)} \otimes \beta_-^{(2)} \triangleleft \beta_+ = Id_{C \square_D C}$ .
- ii)  $\beta_-(c^{(1)} \square_D c^{(2)} \triangleleft h) \otimes \beta_+(c^{(1)} \square_D c^{(2)} \triangleleft h) = c \otimes h, \quad c \in C, h \in H$ .
- iii)  $\beta_-(c_1 \square_D c_2) \triangleleft \beta_+(c_1 \square_D c_2) = \varepsilon(c_1)c_2$ .
- iv)  $\varepsilon(\beta_-(c_1 \square_D c_2))\varepsilon(\beta_+(c_1 \square_D c_2)) = \varepsilon(c_1)\varepsilon(c_2)$ .
- v)  $\beta_-(c_1 \square_D c_2 \triangleleft h) \otimes \beta_+(c_1 \square_D c_2 \triangleleft h) = \beta_-(c_1 \square_D c_2) \otimes \beta_+(c_1 \square_D c_2)h$ .

$$\begin{aligned}
\text{vi)} \quad & [\beta_-(c_1 \square_D c_2)]^{(1)} \otimes [\beta_-(c_1 \square_D c_2)]^{(2)} \otimes \beta_+(c_1 \square_D c_2) = \\
& c_1^{(1)} \otimes \beta_-(c_1^{(2)} \square_D c_2) \otimes \beta_+(c_1^{(2)} \square_D c_2). \\
\text{vii)} \quad & \beta_-(c_1 \square_D c_2)^{(1)} \otimes \beta_-(c_1 \square_D c_2)^{(2)} \otimes \beta_-(c_1 \square_D c_2)^{(3)} \triangleleft \beta_+(c_1 \square_D c_2)^{(1)} \otimes \\
& \beta_-(c_1 \square_D c_2)^{(4)} \triangleleft \beta_+(c_1 \square_D c_2)^{(2)} = c_1^{(1)} \otimes c_1^{(2)} \square_D c_2^{(1)} \otimes c_2^{(2)}.
\end{aligned}$$

*Proof.* The relations i) and ii) are equivalent to  $\beta\beta^{-1} = \text{Id}$  and  $\beta^{-1}\beta = \text{Id}$ , respectively. The relation iii) is obtained by applying  $\varepsilon \otimes \text{Id}$  on the both hand sides of i). The relation iv) is proved by applying  $\varepsilon \otimes \varepsilon$  on the both hand sides of i) and  $H$ -module coalgebra property of  $C$ . The relation v) is equivalent to the right  $H$ -module property of the map  $\beta^{-1}$ . The relation vi) is equivalent to the left  $C$ -comodule map property of the map  $\beta^{-1}$  where the left  $C$  comodule structures of  $C \otimes H$  and  $C \square_D C$  are given by  $c \otimes h \mapsto c^{(1)} \otimes c^{(2)} \otimes h$  and  $c_1 \otimes c_2 \mapsto c_1^{(1)} \otimes c_1^{(2)} \otimes c_2$ , respectively. The relation vii) holds by applying  $\Delta_C \otimes \Delta_C$  on the both hand sides of relation (i) and using  $H$ -module coalgebra property of  $C$ .  $\square$

**3.2. Hopf Galois coextensions as sources of anti-Yetter-Drinfeld modules.** In this subsection we show that Hopf Galois coextensions are sources of producing stable anti Yetter-Drinfeld modules.

For any  $D$ -bicomodule  $C$ , one can assign the following  $C$ -submodule;

$$C^D = \{c \in C, \quad c_{\langle 0 \rangle} \varphi(c_{\langle 1 \rangle}) = c_{\langle 0 \rangle} \varphi(c_{\langle -1 \rangle})\}_{\varphi \in D^*}, \quad (3.6)$$

where  $D^*$  is the linear dual of  $D$ . We set

$$C_D = \frac{C}{W}, \quad (3.7)$$

where

$$W = \left\{ c_{\langle \bar{0} \rangle} \varphi(c_{\langle 1 \rangle}) - c_{\langle 0 \rangle} \varphi(c_{\langle -1 \rangle}), \quad c \in C \right\}_{\varphi \in D^*}.$$

For any coalgebra coextension  $\pi : C \rightarrow D$ , we precisely obtain

$$C^D := \left\{ c \in C, \quad c^{(1)} \varphi(\pi(c^{(2)})) = c^{(2)} \varphi(\pi(c^{(1)})), \quad c \in C \right\}_{\varphi \in D^*}, \quad (3.8)$$

and

$$W = \left\{ c^{(1)} \varphi(\pi(c^{(2)})) - c^{(2)} \varphi(\pi(c^{(1)})), \quad c \in C \right\}_{\varphi \in D^*}. \quad (3.9)$$

**Lemma 3.2.** *Let  $C(D)^H$  be a Hopf Galois coextension with the corresponding  $H$ -action  $\triangleleft : C \otimes H \rightarrow C$ . Then  $\triangleleft$  induces the following  $H$ -action,  $\blacktriangleleft : C^D \otimes H \rightarrow C^D$ .*

*Proof.* It is enough to show that  $c \triangleleft h \in C^D$  for all  $c \otimes h \in C^D \otimes H$ . Indeed,

$$\begin{aligned}
(c \triangleleft h)^{(1)} \varphi(\pi(c \triangleleft h)^{(2)}) &= c^{(1)} \triangleleft h^{(1)} \varphi(\pi(c^{(2)} \triangleleft h^{(2)})) \\
&= c^{(1)} \triangleleft h^{(1)} \varphi(\pi(c^{(2)})) \varepsilon(h^{(2)}) = c^{(1)} \triangleleft h \varphi(\pi(c^{(2)})) \\
&= c^{(2)} \triangleleft h \varphi(\pi(c^{(1)})) = c^{(2)} \triangleleft h^{(2)} \varphi(\pi(c^{(1)})) \varepsilon(h^{(1)}) \\
&= c^{(2)} \triangleleft h^{(2)} \varphi(\pi(c^{(1)} \triangleleft h^{(1)})) = (c \triangleleft h)^{(2)} \varphi(\pi(c \triangleleft h)^{(1)}).
\end{aligned}$$

We use the  $H$ -module coalgebra property of  $C$  in the first equality and  $c \in C^D$  in the fourth equality.  $\square$

One can define a  $D$ -bicomodule structure on  $C \square_D C$  as follows;

$$c \square_D c' \mapsto c \square_D c'^{(1)} \otimes \pi(c'^{(2)}), \quad c \square_D c' \mapsto \pi(c^{(1)}) \otimes c^{(2)} \square_D c'. \quad (3.10)$$

It is easy to check that the coactions defined in (3.10) are well-defined. We set

$$(C \square_D C)_D = \frac{C \square_D C}{W}, \quad (3.11)$$

where

$$W = \langle c \otimes c'^{(1)} \varphi(\pi(c'^{(2)})) - c^{(2)} \otimes c' \varphi(\pi(c^{(1)})) \rangle, \quad (3.12)$$

and  $\varphi \in D^*$ ,  $c \otimes c' \in C \square_D C$ . We show the elements of the quotient by an over line. Although  $T \square_S T$  is not a coalgebra, it is proved in [Bál-Thesis, section 6.4, page 93-95] that the quotient space  $(C \square_D C)_D$  is a coassociative coalgebra where the coproduct and counit maps are given by

$$\Delta(\overline{c \otimes c'}) = \overline{c^{(1)} \square_D c'^{(2)} \otimes c'^{(2)} \square_D c'^{(1)}}. \quad \varepsilon(\overline{c \otimes c'}) = \varepsilon(c)\varepsilon(c'). \quad (3.13)$$

The following lemma can be similarly proved as [HR2][Lemma 4.4] when  $C$  and  $D$  are replaced by  $T$  and  $S$ , respectively.

**Lemma 3.3.** *If  $C(D)^H$  be a Hopf-Galois coextension, then  $\beta$  induces a bijection  $\bar{\beta} : C_D \otimes H \rightarrow (C \square_D C)_D$  where*

$$\bar{\beta}(\bar{c} \otimes h) = \overline{\beta(c \otimes h)}.$$

The following lemma can be similarly proved as [HR2][Lemma 4.5] when  $C$ ,  $D$  and  $H$  are replaced by  $T$ ,  $S$  and  $\mathcal{B}$  respectively.

**Lemma 3.4.** *Let  $C(D)^H$  be a Hopf-Galois coextension. Define*

$$\kappa := (\varepsilon \otimes \text{Id}_H) \circ \bar{\beta}^{-1} : (C \square_D C)_D \rightarrow H.$$

*The map  $\kappa$  is an anti coalgebra map.*

The anti coalgebra map property of the map  $\kappa$  is equivalent to

$$\kappa(\overline{c \otimes c'})^{(1)} \otimes \kappa(\overline{c \otimes c'})^{(2)} = \kappa(\overline{c^{(2)} \otimes c'^{(1)}}) \otimes \kappa(\overline{c^{(1)} \otimes c'^{(2)}}). \quad (3.14)$$

The following lemma introduces some properties of the map  $\kappa$ .

**Lemma 3.5.** *If  $C(D)^H$  be a Hopf Galois coextension then the map  $\kappa$  has the following properties.*

- i)  $\kappa(\overline{c^{(1)} \triangleleft h \square_D c^{(2)} \triangleleft g}) = \varepsilon(c)S(h)g, \quad c \in C, g, h \in H,$
- ii)  $\kappa(\overline{c \square_D c'} \triangleleft h) = \kappa(\overline{c \square_D c' \triangleleft h}), \quad h \in H,$
- iii)  $\kappa(\overline{c \triangleleft h \square_D c'}) = S(h)\kappa(\overline{c \square_D c'}), \quad h \in H,$
- iv)  $c^{(1)} \triangleleft \kappa(\overline{c^{(2)} \square_D c'}) = \varepsilon(c)c', \quad c \otimes c' \in C \square_D C.$

*Proof.* The following computation proves the relation i).

$$\begin{aligned} \kappa(\overline{c^{(1)} \triangleleft h \square_D c^{(2)} \triangleleft g}) &= \kappa(\overline{c^{(1)} \triangleleft h^{(1)} \square_D c^{(2)} \triangleleft \varepsilon(h^{(2)})g}) \\ &= \kappa(\overline{c^{(1)} \triangleleft h^{(1)} \square_D c^{(2)} \triangleleft h^{(2)} S(h^{(3)})g}) = \kappa(\overline{c^{(1)} \triangleleft h^{(1)} \square_D (c^{(2)} \triangleleft h^{(2)}) \triangleleft S(h^{(3)})g}) \\ &= \kappa(\overline{(c \triangleleft h^{(1)})^{(1)} \square_D (c \triangleleft h^{(1)})^{(2)} \triangleleft S(h^{(2)})g}) = \varepsilon(c \triangleleft h^{(1)})S(h^{(2)})g \\ &= \varepsilon(c)\varepsilon(h^{(1)})S(h^{(2)})g = \varepsilon(c)S(h)g. \end{aligned}$$

We use Lemma 3.1(ii) in the fifth equality. The relation ii) is obvious by the right  $H$ -module map property of  $\beta^{-1}$  in Lemma 3.1(v). To prove iii), it is

enough to show that the maps  $m_H \circ (S_H \otimes \kappa)$  and  $\kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C)$ , appearing in both hand sides of iii), have the same inverse in the convolution algebra  $\text{Hom}_k(\mathbb{H} \otimes C \square_D C, \mathbb{H})$ . Here  $m_H$  denotes the multiplication map of  $H$ . To do this, first we show that the map  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a left inverse for  $m_H \circ (S \otimes \kappa)$ , with respect to the convolution product denoted by  $\star$ . Let  $h \otimes \overline{c \otimes c'} \in H \otimes (C \square_D C)_D$ . We have;

$$\begin{aligned}
& [m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa]) \star m_H \circ (S \otimes \kappa)](h \otimes \overline{c \otimes c'}) \\
&= S^{-1}(\overline{\kappa(c^{(1)} \otimes c'^{(2)})})h^{(1)}S(h^{(2)})\overline{\kappa(c^{(2)} \otimes c'^{(1)})} \\
&= \varepsilon(h)S^{-1}(\overline{\kappa(c^{(1)} \otimes c'^{(2)})})\overline{\kappa(c^{(2)} \otimes c'^{(1)})} \\
&= \varepsilon(h)S^{-1}[\overline{\kappa((\overline{c \otimes c'})^{(1)})}]\overline{\kappa((\overline{c \otimes c'})^{(2)})} \\
&= \varepsilon(h)S^{-1}[\overline{\kappa(\overline{c \otimes c'})}^{(2)}]\overline{\kappa(\overline{c \otimes c'})}^{(1)} \\
&= \varepsilon(h)\varepsilon(\overline{\kappa(\overline{c \otimes c'})}) \\
&= \varepsilon(h)\varepsilon(\overline{\kappa(c^{(1)} \otimes c'^{(2)})})\varepsilon(\overline{\kappa(c^{(2)} \otimes c'^{(1)})}) \\
&= \varepsilon(h)\varepsilon(c)\varepsilon(c')1_H = \eta \circ \varepsilon(h \otimes \overline{c \otimes c'}).
\end{aligned}$$

Here  $\eta$  is the unit map of  $H$ . In the previous calculation, we use the anti coalgebra map property of  $\kappa$  in the fourth equality and 3.1(iv) in the seventh equality. Now we show that the map  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a right inverse for the map  $m_H \circ (S \otimes \kappa)$ . For the convenience in the following computation, we denote  $\beta^{-1}(c \otimes d) = \beta_- \otimes \beta_+$ .

$$\begin{aligned}
& [m_H \circ (S \otimes \kappa) \star m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])](h \otimes \overline{c \otimes c'}) \\
&= S(h^{(1)})\overline{\kappa(c^{(1)} \otimes c'^{(2)})}S^{-1}(\overline{\kappa(c^{(2)} \otimes c'^{(1)})})h^{(2)} \\
&= S(h^{(1)})\overline{\kappa(\beta_-^{(1)} \otimes \beta_-^{(4)} \triangleleft \beta_+^{(2)})}S^{-1}(\overline{\kappa(\beta_-^{(2)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(1)})})h^{(2)} \\
&= S(h^{(1)})\overline{\kappa(\beta_-^{(1)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(2)})}S^{-1}(\overline{\varepsilon(\beta_-^{(2)})\beta_+^{(1)}})h^{(2)} \\
&= S(h^{(1)})\overline{\kappa(\beta_-^{(1)} \otimes \beta_-^{(2)} \triangleleft \beta_+^{(2)})}S^{-1}(\overline{\beta_+^{(1)}})h^{(2)} \\
&= S(h^{(1)})\varepsilon(\beta_-)\beta_+^{(2)}S^{-1}(\overline{\beta_+^{(1)}})h^{(2)} = \varepsilon(\beta_-)\varepsilon(\beta_+)S(h^{(1)})h^{(2)} \\
&= \varepsilon(c)\varepsilon(c')\varepsilon(h)1_H = \eta \circ \varepsilon(h \otimes \overline{c \otimes c'}).
\end{aligned}$$

We use Lemma 3.1(vii) in the second equality, Lemma 3.1(ii) and definition of  $\kappa$  in the third and fifth equalities and Lemma 3.1(iv) in the penultimate equality. Therefore we have shown that  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a two-sided inverse for  $m_H \circ (S \otimes \kappa)$ . Now let's check  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a left inverse for the map  $\kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C)$  with respect to the

convolution product.

$$\begin{aligned}
 & [m_H \circ tw \circ (H \otimes [S^{-1} \circ \kappa]) \star \kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C)](h \otimes \overline{c \otimes c'}) \\
 &= m_H \circ tw \circ (h^{(1)} \otimes S^{-1}(\kappa(c^{(1)} \otimes c'^{(2)})))\kappa \circ (\triangleleft \otimes C)(c^{(2)} \otimes h^{(2)} \otimes c'^{(1)}) \\
 &= S^{-1}(\kappa(c^{(1)} \otimes c'^{(2)}))h^{(1)}\kappa(c^{(2)} \triangleleft h^{(2)} \otimes c'^{(1)}) \\
 &= S^{-1}(\kappa(\overline{\beta_-^{(1)} \otimes \beta_-^{(4)} \triangleleft \beta_+^{(2)}}))h^{(1)}\kappa(\overline{\beta_-^{(2)} \triangleleft h^{(2)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(1)}}) \\
 &= S^{-1}(\kappa(\overline{\beta_-^{(1)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(2)}}))h^{(1)}\varepsilon(\beta_-^{(2)})S(h^{(2)})\beta_+^{(1)} \\
 &= S^{-1}(\kappa(\overline{\beta_-^{(1)} \otimes \beta_-^{(2)} \triangleleft \beta_+^{(2)}}))\varepsilon(h)\beta_+^{(1)} \\
 &= S^{-1}(\varepsilon(\beta_-)\beta_+^{(2)})\varepsilon(h)\beta_+^{(1)} \\
 &= \varepsilon(\beta_-)\varepsilon(h)\varepsilon(\beta_+)\mathbf{1}_H = \varepsilon(c)\varepsilon(h)\varepsilon(c')\mathbf{1}_H = \iota \circ \varepsilon(h \otimes \overline{c \otimes c'}).
 \end{aligned}$$

We use Lemma 3.1(vii) in the third equality, Lemma 3.5(i) in the fourth equality, definition of  $\kappa$  and Lemma 3.1(ii) in the sixth equality and Lemma 3.1(iv) in the eighth equality. Now we show that  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a right inverse for  $\kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C)$ .

$$\begin{aligned}
 & [\kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C) \star m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])](h \otimes \overline{c \otimes c'}) \\
 &= \kappa(c^{(1)} \triangleleft h^{(1)} \otimes c'^{(2)})S^{-1}(\kappa(c^{(2)} \otimes c'^{(1)}))h^{(2)} \\
 &= \kappa(\overline{\beta_-^{(1)} \triangleleft h^{(1)} \otimes \beta_-^{(4)} \triangleleft \beta_+^{(2)}})S^{-1}(\kappa(\overline{\beta_-^{(2)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(1)}}))h^{(2)} \\
 &= \kappa(\overline{\beta_-^{(1)} \triangleleft h^{(1)} \otimes \beta_-^{(3)} \triangleleft \beta_+^{(2)}})S^{-1}(\varepsilon(\beta_-^{(2)})\beta_+^{(1)})h^{(2)} \\
 &= \kappa(\overline{\beta_-^{(1)} \triangleleft h^{(1)} \otimes \beta_-^{(2)} \triangleleft \beta_+^{(2)}})S^{-1}(\beta_+^{(1)})h^{(2)} \\
 &= \varepsilon(\beta_-)S(h^{(1)})\beta_+^{(2)}S^{-1}(\beta_+^{(1)})h^{(2)} \\
 &= \varepsilon(\beta_-)\varepsilon(\beta_+)S(h^{(1)})h^{(2)} \\
 &= \varepsilon(c)\varepsilon(c')\varepsilon(h)\mathbf{1}_H = \iota \circ \varepsilon(h \otimes \overline{c \otimes c'}).
 \end{aligned}$$

We use Lemma 3.1(vii) in the second equality, Lemma 3.5(i) and the definition of  $\kappa$  in the third equality, Lemma 3.5(i) in the fifth equality and Lemma 3.1(iv) in the penultimate equality. Therefore  $m_H \circ tw \circ (\text{Id}_H \otimes [S^{-1} \circ \kappa])$  is a two-sided inverse for  $\kappa \circ (\triangleleft \otimes \text{Id}_C) \circ (tw \otimes \text{Id}_C)$  with respect to the convolution product. Therefore we have shown

$$\kappa \circ (\mu_C \otimes C) \circ (tw \otimes C) = \mu_H \circ (S \otimes \kappa).$$

To prove the relation iv), we apply  $\text{Id} \otimes \varepsilon \otimes \text{Id}$  on the both hand sides of Lemma 3.1(vi) and therefore we obtain;

$$\begin{aligned}
 & \beta_- \overline{(c \otimes c')} \otimes \beta_+ \overline{(c \otimes c')} = \varepsilon(\beta_- \overline{(c^{(2)} \otimes c')})c^{(1)} \otimes \beta_+ \overline{(c^{(2)} \otimes c')} \\
 &= c^{(1)} \otimes \kappa(\overline{c^{(2)} \otimes c'}).
 \end{aligned} \tag{3.15}$$

By applying the right action of  $H$  on the previous equation we obtain;

$$\begin{aligned}
 & c^{(1)} \triangleleft \kappa(\overline{c^{(2)} \otimes c'}) = \varepsilon(\beta_- \overline{(c^{(2)} \otimes c')})c^{(1)} \triangleleft \beta_+ \overline{(c^{(2)} \otimes c')} \\
 &= \beta_- \overline{(c \otimes c')} \triangleleft \beta_+ \overline{(c \otimes c')} = \varepsilon(c)c'.
 \end{aligned}$$

We use the Lemma 3.1(iii) on the last equality.  $\square$

**Lemma 3.6.** *Let  $C(D)^H$  be a Hopf Galois coextension. Then  $C^D$  is a right  $(C \square_D C)_D$ -comodule and a left  $H$ -comodule by the following coactions,*

$$\blacktriangledown^{C^D} : C^D \longrightarrow C^D \otimes (C \square_D C)_D, \quad c \longmapsto c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}},$$

and

$$\nabla^{C^D} : C^D \longrightarrow H \otimes C^D, \quad c \longmapsto \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)}. \quad (3.16)$$

*Proof.* First we show that the coaction  $\blacktriangledown^{C^D}$  is well-defined. The following computation proves  $c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}} \in C^D \otimes C \otimes C$ .

$$\begin{aligned} c^{(3)} \varphi(\pi(c^{(2)})) \otimes \overline{c^{(4)} \square_D c^{(1)}} &= c^{(3)} \otimes \overline{c^{(4)} \square_D c^{(1)} \varphi(\pi(C^{(2)}))} \\ &= c^{(3)} \otimes \overline{c^{(4)(2)} \varphi(\pi(c^{(4)(1)})) \otimes \varepsilon(c^{(1)}) c^{(2)}} = c^{(2)} \varphi(\pi(c^{(3)})) \otimes \overline{c^{(4)} \square_D c^{(1)}}. \end{aligned}$$

We use the definition of  $W$  which given in (3.12). Furthermore since  $C$  is a coalgebra on a field then  $c \in C^D$  implies  $c^{(1)} \otimes \pi(c^{(2)}) = c^{(2)} \otimes \pi(c^{(1)})$ . Therefore we have;

$$c^{(1)} \otimes c^{(2)} \otimes c^{(3)} \otimes \pi(c^{(4)}) = c^{(2)} \otimes c^{(3)} \otimes c^{(4)} \otimes \pi(c^{(1)}), \quad (3.17)$$

which shows

$$c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}} \in C \otimes C \square_D C.$$

The following computation shows that  $\blacktriangledown^{C^D}$  defines a coassociative coaction.

$$\begin{aligned} [\blacktriangledown^{C^D} \otimes \text{Id}_{(C \square_D C)_D}] \circ \blacktriangledown^{C^D}(c) &= (\blacktriangledown^{C^D} \otimes \text{Id}_{(C \square_D C)_D})(c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}}) \\ &= (c^{(3)} \otimes \overline{c^{(4)} \square_D c^{(2)}} \otimes \overline{c^{(5)} \square_D c^{(1)}}) = c^{(2)} \otimes \overline{c^{(3)(1)} \square_D c^{(1)(2)}} \otimes \overline{c^{(3)(2)} \square_D c^{(1)(1)}} \\ &= (\text{Id}_{C^D} \otimes \Delta)(c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}}) = (\text{Id}_{C^D} \otimes \Delta) \circ \blacktriangledown^{C^D}(c). \end{aligned}$$

The counitality of the coaction induced by  $\blacktriangledown^{C^D}$  can be shown as follows.

$$(\text{Id}_{C^D} \otimes \varepsilon) \circ \blacktriangledown^{C^D}(c) = (\text{Id}_{C^D} \otimes \varepsilon)(c^{(2)} \otimes \overline{c^{(3)} \square_D c^{(1)}}) = c^{(2)} \varepsilon(c^{(3)}) \varepsilon(c^{(1)}) = \text{Id}_{C^D}(c).$$

Therefore  $C^D$  is a right coassociative and counital  $(C \square_D C)_D$ -comodule.

The following computation shows that  $\nabla^{C^D}$  defines a coassociative  $H$ -coaction on  $C^D$ .

$$\begin{aligned} (\text{Id}_H \otimes \nabla^{C^D}) \circ \nabla^{C^D}(c) &= (\text{Id}_H \otimes \nabla^{C^D})(\kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)}) \\ &= \kappa(\overline{c^{(5)} \square_D c^{(1)}}) \otimes \kappa(\overline{c^{(4)} \square_D c^{(2)}}) \otimes c^{(3)} \\ &= \kappa(\overline{c^{(3)} \square_D c^{(1)}})^{(1)} \otimes \kappa(\overline{c^{(3)} \square_D c^{(1)}})^{(2)} \otimes c^{(2)} \\ &= (\Delta \otimes \text{Id}_H)(\kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)}) = (\Delta \otimes \text{Id}_H) \circ \nabla^{C^D}(c). \end{aligned}$$

We use the anti coalgebra map property of  $\kappa$  in the third equality. To prove the counitality of the coaction for all  $c \in C^D$  we have;

$$\begin{aligned} (\varepsilon \otimes \text{Id}_{C^D}) \circ \nabla^{C^D}(c) &= (\varepsilon \otimes \text{Id}_{C^D})(\kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)}) = \\ &= \varepsilon \circ \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)} = \varepsilon(c^{(3)}) \varepsilon(c^{(1)}) \otimes c^{(2)} = \text{id}_{C^D}(c). \end{aligned}$$

We use the anti coalgebra map property of  $\kappa$  in the third equality. Therefore  $C^D$  is a left  $H$ -comodule.  $\square$

**Lemma 3.7.** *Let  $C(D)^H$  be Hopf Galois coextension. Then  $C_D$  is a left  $(C \square_D C)_D$ -comodule and a right  $H$ -comodule by the following coactions,*

$$\nabla^{C_D} : C_D \longrightarrow (C \square_D C)_D \otimes C_D, \quad c \longmapsto \overline{c^{(1)} \square_D c^{(3)}} \otimes c^{(2)},$$

and

$$\nabla^{C_D} : C_D \longrightarrow C_D \otimes H, \quad c \longmapsto c^{(2)} \otimes \kappa(\overline{c^{(1)} \square_D c^{(3)}}). \quad (3.18)$$

*Proof.* Since  $C_D = \frac{C}{\overline{W}}$  is a coalgebra this coaction is well-defined. The following computation shows that this coaction is coassociative.

$$\begin{aligned} & (\Delta_{C \square_D C_D} \otimes \text{Id}_C) \circ \nabla^{C_D}(c) \\ &= \overline{(t^{(1)} \otimes t^{(3)})^{(1)}} \otimes \overline{(t^{(1)} \otimes t^{(3)})^{(2)}} \otimes t^{(2)} = t^{(1(1))} \otimes t^{(3(2))} \otimes t^{(1(2))} \otimes t^{(3(1))} \otimes t^{(2)} \\ &= t^{(1)} \otimes t^{(5)} \otimes t^{(2)} \otimes t^{(4)} \otimes t^{(3)} = t^{(1)} \otimes t^{(3)} \otimes t^{(2(1))} \otimes t^{(2(3))} \otimes t^{(2(2))} \\ &= (\text{Id}_{C \square_D C_D} \otimes \nabla_C) \circ \nabla_C. \end{aligned}$$

The counitality of the coaction is obvious. Similar to Lemma 3.6 we can show that the anti coalgebra map  $\kappa$  turns this coaction to a right coaction of  $H$  on  $C_D$ .  $\square$

Now we are ready to show that the Hopf Galois coextensions of coalgebras are the sources of stable anti Yetter-Drinfeld modules.

**Theorem 3.8.** *Let  $C(D)^H$  be a Hopf Galois coextension. Then  $C^D$  is a right-left SAYD module over  $H$  by the coaction  $\nabla^{C^D}$  defined in (3.18) and action  $\blacktriangleleft$  defined in Lemma 3.2.*

*Proof.* The AYD condition holds because for all  $c \in C^D$  and  $h \in H$ , we have

$$\begin{aligned} & S(h^{(3)})c_{\langle -1 \rangle} h^{(1)} \otimes c_{\langle 0 \rangle} \blacktriangleleft h^{(2)} = S(h^{(3)})\kappa(\overline{c^{(3)} \otimes c^{(1)}})h^{(1)} \otimes c^{(2)} \blacktriangleleft h^{(2)} \\ &= \kappa(\overline{c^{(3)} \blacktriangleleft h^{(3)} \otimes c^{(1)} h^{(1)} \otimes c^{(2)} \blacktriangleleft h^{(2)}}) = \kappa(\overline{c^{(3)} \blacktriangleleft h^{(3)} \otimes c^{(1)} \blacktriangleleft h^{(1)}}) \otimes c^{(2)} \blacktriangleleft h^{(2)} \\ &= \kappa(\overline{(c \blacktriangleleft h)^{(3)} \otimes (c \blacktriangleleft h)^{(1)}}) \otimes (c \blacktriangleleft h)^{(2)} = \nabla^{C^D}(c \blacktriangleleft h). \end{aligned}$$

We use Lemma 3.5(iii) in the second equality, Lemma 3.5(ii) in the third equality and right  $H$ -module coalgebra property of  $C$  in the fourth equality. The following computation shows the stability condition.

$$\begin{aligned} & c_{\langle 0 \rangle} \blacktriangleleft c_{\langle -1 \rangle} = c^{(2)} \blacktriangleleft \kappa(\overline{c^{(3)} \otimes c^{(1)}}) = \\ & c^{(2(1))} \blacktriangleleft \kappa(\overline{c^{(2(2))} \otimes c^{(1)}}) = \varepsilon(c^{(2)})c^{(1)} = c. \end{aligned}$$

The penultimate equality holds by Lemma 3.5(iv).  $\square$

The following statement which is proved in [HR2][Remark 5.4] will be used in the sequel subsection.

**Lemma 3.9.** *For any coalgebra coextension  $\pi : C \longrightarrow D$  we have;*

$$C^D \cong D \square_{D^e} C,$$

where the isomorphism is given by

$$\xi(d \square_{D^e} c) = \varepsilon_D(d)c, \quad \text{and} \quad \xi^{-1}(c) = \pi(c^{(1)}) \square_{D^e} c^{(2)}.$$



**3.3. Cohomology theories related to Hopf Galois coextensions.** In this subsection we show that the cyclic cohomology of the Hopf algebra involved in the coextensions with coefficients in the SAYD module  $C^D$  is isomorphic to the relative cyclic cohomology of coalgebra coextensions.

**Definition 3.10.** *Let  $D$  be a coalgebra and  $M$  and  $N$  be a  $D$ -bicomodule. The cyclic cotensor product of  $M$  and  $N$  is defined by*

$$M \widehat{\square}_D N := (M \square_D N) \square_{D^e} D \cong D \square_{D^e} (M \square_D N). \quad (3.19)$$

The cyclic cotensor product also can be defined for a finite number of arbitrary  $D$ -bicomodules  $M_1, \dots, M_n$  as follows;

$$(M_1 \widehat{\square}_D \dots \widehat{\square}_D M_n) := (M_1 \square_D \dots \square_D M_n) \square_{D^e} D. \quad (3.20)$$

One has

$$(M_1 \widehat{\square}_D \dots \widehat{\square}_D M_n) \cong (M_1 \square_D \dots \square_D M_n)^D. \quad (3.21)$$

Let  $\pi : C \rightarrow D$  be a coalgebra coextension and  $M$  be a  $C$ -bicomodule. Therefore  $M$  is a  $D$ -bicomodule by the following comodule structures;

$$c \mapsto \pi(c_{<-1>}) \otimes c_{<0>}, \quad c \mapsto c_{<0>} \otimes \pi(c_{<1>}). \quad (3.22)$$

For all  $n \in \mathbb{N}$  and  $0 \leq i \leq n$  we define cofaces

$$d_i : M \widehat{\square}_D C \widehat{\square}_D^{n-1} \longrightarrow M \widehat{\square}_D C \widehat{\square}_D^{n-1}, \quad (3.23)$$

which are given by

$$d_i(m \square_D c_1 \square_D \dots \square_D c_n) = \begin{cases} m_{<0>} \square_D m_{<1>} \square_D c_1 \square_D \dots \square_D c_n, & i = 0, \\ m \square_D c_1 \square_D \dots \square_D \Delta(c_i) \square_D \dots \square_D c_n, & 0 < i < n, \\ m_{<0>} \square_D c_1 \square_D \dots \square_D c_n \square_D m_{<-1>}, & i = n. \end{cases} \quad (3.24)$$

Also for  $0 \leq i \leq n-1$  we define the codegeneracies

$$s_i : M \widehat{\square}_D C \widehat{\square}_D^{n-1} \longrightarrow M \widehat{\square}_D C \widehat{\square}_D^{n-2}, \quad (3.25)$$

which are given by

$$s_i(m \square_D c_1 \square_D \dots \square_D c_n) := m \square_D c_1 \square_D \dots \square_D \varepsilon(c_{i+1}) \square_D \dots \square_D c_n. \quad (3.26)$$

One can easily check that for  $0 \leq i \leq n-1$ , the cofaces  $d_i$  are well-defined. To show that the last coface is well-defined, one notes that the condition

$$m \otimes c_1 \otimes \dots \otimes c_n \in (M \square C \square \dots \square C)^D,$$

implies

$$\pi(m_{<-1>}) \otimes m_{<0>} \otimes c_1 \otimes \dots \otimes c_n = \pi(c_n^{(2)}) \otimes m \otimes c_1 \otimes \dots \otimes c_{n-1} \otimes c_n^{(1)}. \quad (3.27)$$

Therefore the following computation proves that the last coface is well-defined.

$$\begin{aligned} & m_{<0>} \otimes c_1 \otimes \dots \otimes c_{n-1} \otimes c_n^{(1)} \otimes \pi(c_n^{(2)}) \otimes m_{<-1>} \\ &= m_{<0><0>} \otimes c_1 \otimes \dots \otimes c_{n-1} \otimes c_n \otimes \pi(m_{<-1>}) \otimes m_{<0><-1>} \\ &= m_{<0>} \otimes c_1 \otimes \dots \otimes c_{n-1} \otimes \pi(m_{<-1>}^{(1)}) \otimes m_{<-1>}^{(2)}. \end{aligned}$$

We use (3.27) in the first equality and the coassociativity of  $C$ -coaction of  $M$  on the second equality.

**Theorem 3.11.** *Let  $\pi : C \rightarrow D$  be a coalgebra coextension and  $M$  be a  $D$ -bicomodule.*

a) *The module  $C^n(C(D), M) = M \widehat{\square}_D C \widehat{\square}_{D^n}$  is a cosimplicial object with the cofaces and codegeneracies as defined in (3.24) and (3.25).*

b) *Let  $M = C$  and set*

$$t : C \widehat{\square}_D C \widehat{\square}_{D^n} \longrightarrow C \widehat{\square}_D C \widehat{\square}_{D^n},$$

which is given by

$$t_n(c_0 \otimes c_1 \otimes \dots \otimes c_n) = c_1 \otimes \dots \otimes c_n \otimes c_0. \quad (3.28)$$

Then  $(C^n(C(D), C), d_i, s_i, t_n)$  is a cocyclic module.

The cyclic cohomology of the preceding cocyclic module is called relative cyclic cohomology of coalgebra coextension  $C(D)$ . The Theorem 3.11 is the dual statement of [J-S][Theorem 1.5]. In fact

$$C^n(C(D), C) = D \square_{D^e} \underbrace{C \square_D \dots \square_D C}_{n+1 \text{ times}}. \quad (3.29)$$

For any Hopf Galois coextension  $C(D)^H$  the canonical map  $\beta : C \otimes H \rightarrow C \square_D C$  induces the following bijection,

$$\beta^n : C \otimes H^{\otimes n} \longrightarrow C^{\square_D(n+1)}$$

$$c \otimes h_1 \otimes \dots \otimes h_n \longmapsto c^{(1)} \otimes c^{(2)} \triangleleft h_1^{(1)} \otimes c^{(3)} \triangleleft h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n+1)} \triangleleft h_1^{(n)} \dots h_{n-1}^{(2)} h_n.$$

Now by applying  $D \square_{D^e} -$  on the both sides of the map  $\beta^n$  we obtain the following isomorphism of  $C$ -bicomodules;

$$\varphi : C^D \otimes H^{\otimes n} \cong D \square_{D^e} C \otimes H^{\otimes n} \longrightarrow D \square_{D^e} C \square_D C^{\square_D n} \cong C^{\square_D n+1},$$

which is given by

$$\begin{aligned} & c \otimes h_1 \otimes \dots \otimes h_n \longmapsto \\ & c^{(1)} \otimes c^{(2)} \triangleleft h_1^{(1)} \otimes c^{(3)} \triangleleft h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n)} \triangleleft h_1^{(n-1)} \dots h_{n-1}^{(1)} \otimes \\ & c^{(n+1)} \triangleleft h_1^{(n)} \dots h_{n-1}^{(2)} h_n. \end{aligned} \quad (3.30)$$

The inverse map

$$\varphi^{-1} : C^{\square_D n+1} \longrightarrow C^D \otimes H^{\otimes n},$$

is given by

$$\begin{aligned} & c_0 \otimes \dots \otimes c_n \longmapsto \\ & c_0^{(1)} \otimes \kappa(c_0^{(2)} \otimes c_1^{(1)}) \otimes \kappa(c_1^{(2)} \otimes c_2^{(1)}) \otimes \dots \otimes \kappa(c_{n-2}^{(2)} \otimes c_{n-1}^{(1)}) \otimes \\ & \kappa(c_{n-1}^{(2)} \otimes c_n). \end{aligned} \quad (3.31)$$

By Theorem 3.8, the subspace  $C^D$  is a right-left SAYD module over  $H$ . Using the cocyclic module (1.10), the Hopf cyclic cohomology of  $H$  with

coefficients in  $C^D$  is computed by the following cocyclic module;

$$\begin{aligned}
\delta_i(c \otimes \tilde{h}) &= h_1 \otimes \cdots \otimes h_i \otimes 1_H \otimes h_{i+1} \otimes \cdots \otimes h_n \otimes c, \quad 0 \leq i \leq n, \\
\delta_{n+1}(c \otimes \tilde{h}) &= h_1^{(1)} \otimes \cdots \otimes h_n^{(1)} \otimes S(h_1^{(2)} \cdots h_n^{(2)}) \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)}, \\
\sigma_i(c \otimes \tilde{h}) &= h_1 \otimes \cdots \otimes h_i h_{i+1} \otimes \cdots \otimes h_n \otimes c, \\
\sigma_n(c \otimes \tilde{h}) &= h_1 \otimes \cdots \otimes h_{n-1} \varepsilon(h_n) \otimes c, \\
\tau_n(c \otimes \tilde{h}) &= h_2^{(1)} \otimes \cdots \otimes h_n^{(1)} \otimes S(h_1^{(2)} \cdots h_n^{(2)}) \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)} \triangleleft h_1^{(1)},
\end{aligned} \tag{3.32}$$

where  $c \otimes \tilde{h} = c \otimes h_1 \otimes \cdots \otimes h_n$ . We denote the above cocyclic module by  $C^n(H, C^D)$ . Now using the isomorphism  $\varphi$  given in (3.30) we define the following map,

$$\psi : C^n(H, C^D) \longrightarrow C^n(C(D), C), \tag{3.33}$$

which is given by

$$\begin{aligned}
&h_1 \dots \otimes h_n \otimes c \longmapsto \\
&c^{(1)} \otimes c^{(2)} \triangleleft h_1^{(1)} \otimes c^{(3)} \triangleleft h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n)} \triangleleft h_1^{(n-1)} \dots h_{n-1}^{(1)} \otimes \\
&c^{(n+1)} \triangleleft h_1^{(n)} \dots h_{n-1}^{(2)} h_n,
\end{aligned} \tag{3.34}$$

with the inverse map;

$$\begin{aligned}
&c_0 \otimes \dots \otimes c_n \longmapsto \\
&\kappa(c_0^{(2)} \otimes c_1^{(1)}) \otimes \kappa(c_1^{(2)} \otimes c_2^{(1)}) \otimes \dots \otimes \kappa(c_{n-2}^{(2)} \otimes c_{n-1}^{(1)}) \otimes \\
&\kappa(c_{n-1}^{(2)} \otimes c_n) \otimes c_0^{(1)}.
\end{aligned} \tag{3.35}$$

Now we are ready to mention the main result.

**Theorem 3.12.** *Let  $C(D)^H$  be a Hopf Galois coextension. The map  $\psi$  introduced in (3.34) induces an isomorphism between the cocyclic modules  $C^n(H, C^D)$  and  $C^n(C(D), C)$  and we obtain*

$$HC^n(H, C^D) \cong HC^n(C(D), C). \tag{3.36}$$

*Proof.* It is straight forward and easy to prove that the map  $\psi$  commutes with all cofaces, except the last one, and also codegeneracies. Here we show

that  $\psi$  commutes with the cyclic operator and therefore with the last coface.

$$\begin{aligned}
 & t_n(\psi(h_1 \otimes \dots \otimes h_n \otimes c)) \\
 &= t_n(c^{(1)} \otimes c^{(2)} \triangleleft h_1^{(1)} \otimes c^{(3)} \triangleleft h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n)} \triangleleft h_1^{(n-1)} \dots h_{n-1}^{(1)} \otimes \\
 & \quad c^{(n+1)} \triangleleft h_1^{(n)} \dots h_{n-1}^{(2)} h_n) \\
 &= c^{(2)} \triangleleft h_1^{(1)} \otimes c^{(3)} \triangleleft h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n)} \triangleleft h_1^{(n-1)} \dots h_{n-1}^{(1)} \otimes \\
 & \quad c^{(n+1)} \triangleleft h_1^{(n)} \dots h_{n-1}^{(2)} h_n \otimes c^{(1)} \\
 &= c^{(2)} h_1^{(1)} \otimes c^{(3)} h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n+1)} h_1^{(n)} \dots h_{n-1}^{(1)} h_n \otimes \\
 & \quad c^{(n+2)} \triangleleft \kappa(\overline{c^{(n+3)} \square_D c^{(1)}}) \\
 &= c^{(2)} h_1^{(1)} \otimes c^{(3)} h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n+1)} h_1^{(n)} \dots h_n^{(1)} \otimes \\
 & \quad c^{(n+2)} h_1^{(n+1)} h_2^{(n)} \dots h_n^{(2)} S(h_n^{(3)}) \dots S(h_1^{(n+2)}) \kappa(\overline{c^{(n+3)} \square_D c^{(1)}}) \\
 &= c^{(2)} h_1^{(1)} \otimes c^{(3)} h_1^{(2)} h_2^{(1)} \otimes \dots \otimes c^{(n+1)} h_1^{(n)} \dots h_n^{(1)} \otimes \\
 & \quad c^{(n+2)} h_1^{(n+1)} h_2^{(n)} \dots h_n^{(2)} S(h_1^{(n+2)} \dots h_n^{(3)}) \kappa(\overline{c^{(n+3)} \square_D c^{(1)}}) \\
 &= (c^{(2)} h_1^{(1)})^{(1)} \otimes (c^{(2)} h_1^{(1)})^{(2)} h_2^{(1)} \otimes \dots \otimes (c^{(2)} h_1^{(1)})^{(n)} h_2^{(1)(n-1)} \dots h_n^{(1)(1)} \otimes \\
 & \quad (c^{(2)} h_1^{(1)})^{(n+1)} h_2^{(1)(n)} \dots h_n^{(1)(2)} S(h_1^{(2)} \dots h_n^{(2)}) \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \\
 &= \psi(h_2^{(1)} \otimes \dots \otimes h_n^{(1)} \otimes S(h_1^{(2)} \dots h_n^{(2)}) \kappa(\overline{c^{(3)} \square_D c^{(1)}}) \otimes c^{(2)} \triangleleft h_1^{(1)}) \\
 &= \psi\tau(h_1 \otimes \dots \otimes h_n \otimes c).
 \end{aligned}$$

We use the Lemma 3.5(iv) in third equality.  $\square$

We summarize the headlines of this section in the following remark which are the dual ones in Remark 2.5.

**Remark 3.13.** For any Hopf Galois coextension  $C(D)^H$ , the following statements hold.

- i) The relative cocyclic module of coalgebra coextensions is a subspace which is given by,

$$D \otimes_{D^e} \underbrace{C \square_D \dots \square_D C}_{n+1 \text{ times}} = C \widehat{\square}_D \dots \widehat{\square}_D C = (C \square_D \dots \square_D C)^D.$$

- ii) The objects  $C \otimes H$  and  $C \square_D C$  are isomorphic in the category of  ${}^C \mathcal{M}_H$ .
- iii) The quotient space  $C_D = \frac{C}{\overline{W}}$  is a coalgebra.
- iv) The quotient space  $(C \square_D C)_D$  is a coalgebra.
- v) The quotient space  $C_D$  is a left  $(C \square_D C)_D$ -comodule and a right  $H$ -comodule.
- vi) The subspace  $C^D$  is a right  $(C \square_D C)_D$ -comodule and a left  $H$ -comodule.
- vii) The subspace  $C^D \cong D \square_{D^e} C$  is a right-left SAYD module over  $H$ .
- viii)  $HC^*(H, C_D) \cong HC^*(C(D), C)$ .

## REFERENCES

- [Bál-Thesis] I. Bálint, *Quantum groupoid symmetry, with an application to low dimensional algebraic quantum field theory*, PhD thesis, Loránd Eötvös University, 154 pages, (2008).
- [BO] G. Böhm, *Galois theory for Hopf algebroids*, Ann. Univ. Ferrara Sez. VII (N.S.) 51, pp. 233–262. math.RA/0409513, (2005).
- [BS] G. Böhm, and D. Stefan, *(co)cyclic (co)homology of bialgebroids: An approach via (co)monads*, Commun. Math. Phys **282**, pp. 239–286, (2008).
- [BH] T. Brzezinski, and P. Hajac, *Coalgebra extensions and algebra coextensions of Galois type*, Commun. Algebra **27(3)**, pp. 1347–1368, (1999).
- [BM] T. Brzezinski, and S. Majid, *Coalgebra bundles*. Commun. Math. Phys., 191:pp. 467–492, (1998).
- [C-Book] A. Connes, *Noncommutative geometry*, Academic Press, (1994).
- [CM98] A. Connes and H. Moscovici, *Hopf algebras, cyclic cohomology and the transverse index theorem*, Comm. Math. Phys. **198**, pp. 199–246, (1998).
- [Dur] M. Durdevic, *Geometry of quantum principal bundles, I, II*, Comm. Math. Phys. 175 (1996), pp. 427–521; Rev. Math. Phys. 9, pp. 531–607, (1997).
- [Groth-SGA1] A. Grothendieck, Raynaud, Michéle (2003) [1971], *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 3, Paris: Société Mathématique de France, arXiv:math/0206203, ISBN 978-2-85629-141-2, MR2017446.
- [Ha] P. M. Hajac. *Strong connections on quantum principal bundles*, Commun. Math. Phys. 182, pp. 579–617, (1996).
- [HKRS1] P.M. Hajac, M. Khalkhali, B. Rangipour and Y. Sommerhäuser, *Stable anti-Yetter-Drinfeld modules*, C. R. Math. Acad. Sci. Paris **338**, pp. 587–590, (2004).
- [HKRS2] P.M. Hajac, M. Khalkhali, B. Rangipour and Y. Sommerhäuser, *Hopf-cyclic homology and cohomology with coefficients*, C. R. Math. Acad. Sci. Paris **338**, pp. 667–672, (2004).
- [HR] M. Hassanzadeh, and B. Rangipour, *Equivariant Hopf Galois extensions and Hopf cyclic cohomology*, To appear in Journal of noncommutative Geometry, 33 pages, (2010).
- [HR2] M. Hassanzadeh, and B. Rangipour, *Quantum Groupoids and their Hopf Cyclic Cohomology*, <http://arxiv.org/abs/1108.0030>, 54 pages, (2011).
- [J-S] P. Jara, and D. Stefan, *Hopf-cyclic homology and relative cyclic homology of Hopf-Galois extensions*, Proc. London Math. Soc. (3) **93**, pp. 138–174, (2006).
- [Kay] A. Kaygun, *The Universal Hopf cyclic theory*, Journal of Noncommutative Geometry, Vol. 2 , No. 3, pp. 333-351, (2008).
- [Kh] M. Khalkhali, *A short survey of cyclic cohomology*, in Quanta of Maths, Alain Connes’ festshrift, published by Clay Math Institute, and American Mathematical Society, ( arXiv:1008.1212), (2010).
- [KR02] M. Khalkhali, and B. Rangipour, *A new cyclic module for Hopf algebras*, K-Theory 27, pp. 111–131, (2002).
- [KR03] M. Khalkhali, and B. Rangipour, *A note on cyclic duality and Hopf algebras*, Comm. Algebra 33, no. 3, pp. 763–773, (2005) .
- [KT] HF. Kreimer, and M. Takeuchi, *Hopf algebras and Galois extensions of an algebra*, Indiana Univ. Math. J. 30, pp. 675–692, (1981) .
- [KZ] U. Kraehmer, and B. Zielinski, *On piecewise trivial Hopf-Galois extensions*, Czech. J. Phys. 56/10-11, pp. 1221–1226, (2006).
- [Lo] J.-L. Loday, *Cyclic Homology*. Springer-Verlag (1992).
- [J-S] P. Jara, and D. Stefan, *Hopf-cyclic homology and relative cyclic homology of Hopf-Galois extensions*, Proc. London Math. Soc. (3) **93**, pp. 138–174, (2006).
- [Q] D. Quillen, *Cyclic cohomology and algebra extensions*, KTheory, 3 , pp. 205–246, (1989).
- [Sch] P. Schauenburg, *Duals and doubles of quantum groupoids ( $\times_R$ -Hopf algebras)*, in: N. Andruskiewitsch, W.R. Ferrer-Santos and H.-J. Schneider (eds.) AMS Contemp. Math. **267**, AMS Providence, pp. 273–293, (2000).

- [Schn1] H.-J. Schneider, *Some remarks on exact sequences of quantum groups*. Comm. Algebra **21**, no. 9, pp. 3337–3357, (1993).
- [Schn2] H.-J. Schneider *Principal homogeneous spaces for arbitrary Hopf algebras*, Isr. J. Math., 72, pp. 167195, (1990).
- [SS] P. Schauenburg, and H .J. Schneider *On generalized Hopf Galois extensions*, Journal of Pure and Applied Algebra, Volume: 202, (2004) .
- [Ver] J. Vercruysse, *Galois theory for corings and comodules*, PhD-thesis, Vrije Universiteit Brussel, (2007).

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, IHES, BURES SUR YVETTE, FRANCE.,  
*E-mail address:* `mhassanz@ihes.fr`