

**Algebraic rational cells, equivariant intersection theory,
and Poincaré duality**

Richard Paul GONZALES VILCARROMERO



Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Avril 2014

IHES/M/14/15

ALGEBRAIC RATIONAL CELLS, EQUIVARIANT INTERSECTION THEORY, AND POINCARÉ DUALITY

RICHARD P. GONZALES *

ABSTRACT. We provide a notion of algebraic rational cell with applications to intersection theory on singular varieties with torus action. Based on this notion, we study the algebraic analogue of \mathbb{Q} -filtrable varieties: algebraic varieties where a torus acts with isolated fixed points, such that the associated Białynicki-Birula decomposition consists of algebraic rational cells. We show that the rational equivariant Chow group of any \mathbb{Q} -filtrable variety is freely generated by the cell closures. We apply this result to group embeddings, and more generally to spherical varieties. In view of the localization theorem for equivariant operational Chow rings, we get some conditions for Poincaré duality in this setting.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let \mathbb{k} be an algebraically closed field. The most commonly studied cell decompositions in algebraic geometry are those obtained by the method of Białynicki-Birula [B1]. If $\mathbb{G}_m \simeq \mathbb{k}^*$ acts on a smooth projective variety X with finitely many fixed points x_1, \dots, x_m , then $X = \bigsqcup X_i$, where

$$X_i = \{x \in X \mid \lim_{t \rightarrow 0} tx = x_i\}.$$

Moreover, the cells X_i are isomorphic to affine spaces; that is $X_i \simeq \mathbb{A}^{n_i}$, for every i . From this one concludes e.g. that the Chow groups of X are freely generated by the classes of the cell closures $\overline{X}_i \subseteq X$. This is quite notable, because the Chow groups of smooth varieties need not be finitely generated (consider e.g. a smooth projective curve of genus one). If $\mathbb{k} = \mathbb{C}$, then this decomposition implies that X has no singular cohomology in odd degrees, and that the cycle map $cl_X : A_*(X) \rightarrow H_*(X)$ is an isomorphism, to mention just a few interesting applications. The *BB-decomposition* makes sense even if X is singular, but the cells may no longer be so well-behaved.

In [G1] we study the BB-decompositions of possibly singular complex projective varieties, assuming that the cells are rationally smooth (i.e. *rational cells*). Recall that a complex algebraic variety X , of dimension n , is called *rationally smooth* if, for every $x \in X$,

$$H^m(X, X - \{x\}) = (0) \text{ if } m \neq 2n, \text{ and } H^{2n}(X, X - \{x\}) = \mathbb{Q}.$$

* Supported by the Institut des Hautes Études Scientifiques, the Max-Planck-Institut für Mathematik, and TÜBİTAK Project No. 112T233.

Such varieties satisfy Poincaré duality with rational coefficients. If X_i as above is a rational cell, then $\mathbb{P}(X_i) := (X_i \setminus \{x_i\})/\mathbb{G}_m$ is a rational cohomology complex projective space. Many important results on the equivariant cohomology of T -varieties admitting a BB-decomposition by rational cells are provided in [G1]; for instance, such varieties have no cohomology in odd degrees and their equivariant cohomology is freely generated by the classes of the cell closures.

The purpose of this paper is to provide analogues of such results, along with conditions for Poincaré duality, in the context of intersection theory for schemes with torus action. For this, we introduce the notion of *algebraic rational cell*. Concisely, let X be an affine \mathbb{G}_m -variety with an attractive fixed point x . Then X is an algebraic rational cell if $\mathbb{P}(X) := [X \setminus \{0\}]/\mathbb{G}_m$ satisfies

$$A_*(\mathbb{P}(X))_{\mathbb{Q}} \simeq A_*(\mathbb{P}^{n-1})_{\mathbb{Q}},$$

where $n = \dim(X)$. The definition applies to actions of higher dimensional tori as well (Definition 3.1). Algebraic rational cells are modelled after (topological) rational cells [G1], although the resulting objects are not equivalent. In what follows, we show that algebraic rational cells are a good substitute for the notion of affine space in the study of Chow groups of singular varieties. This has applications to embedding theory (Section 5), Poincaré duality (Section 6) and the topology of spherical varieties (Section 7). In addition, some links between our present approach and that of [G1] are built (Theorems 5.5, 5.9, and 7.3). The techniques are mostly algebraic, and no essential use of the cycle map is made, except in Section 7.

This article is organized as follows. Section 2 briefly reviews equivariant Chow groups and equivariant multiplicities at nondegenerate fixed points. The section concludes with some inequalities relating Chow groups and fixed point loci. In Section 3, we study the intersection-theoretical properties of algebraic rational cells (Proposition 3.3, Theorem 3.4, Corollary 3.8). Next, in Section 4, we introduce the concept of (algebraically) \mathbb{Q} -filtrable spaces: projective T -varieties with isolated fixed points, such that the associated BB-decomposition is filtrable, and consists of algebraic rational cells (Definition 4.1). The key result is given below (rational coefficients are understood).

Theorem (4.3, 4.4). *Let X be a \mathbb{Q} -filtrable T -variety. Then the T -equivariant Chow group of X is a free $A_T^*(pt)$ -module of rank $|X^T|$. In fact, it is freely generated by the classes of the closures of the cells $X_+(x_i, \lambda)$. Consequently, $A_*(X)$ is also freely generated by the classes of the cell closures $\overline{X_+(x_i, \lambda)}$.*

Having developed the theoretical framework for the study of \mathbb{Q} -filtrable varieties, we devote the last three sections to examples and applications. In Section 5 we apply the theory to group embeddings. Let M be a reductive monoid with zero, unit group G and maximal torus $T \subset G$. Recall that $T \times T$ acts on M with 0 as an attractive fixed point. Moreover, M has finitely many $T \times T$ -invariant curves, and they are indexed by the rank-one

elements of the Renner monoid \mathcal{R}_1 . In general, $\dim M \leq |\mathcal{R}_1|$. Similar remarks apply to the affine toric variety $\overline{T} \subset M$. In this case, the T -invariant curves are indexed by the poset $E_1(\overline{T})$ of rank-one idempotents of \overline{T} . Thus $\dim \overline{T} \leq |E_1(\overline{T})|$. In this context, one of our main results states that reductive monoids which are algebraic rational cells are characterized in the same way as rationally smooth monoids. This adds to Renner's list of equivalences from [R2] and [R3].

Theorem (5.5). *Let M be a reductive monoid with zero and unit group G . Then the following are equivalent.*

- (a) $M \sim_0 \prod_i M_{n_i}(\mathbb{k})$.
- (b) If T is a maximal torus of G , then $\dim T = |E_1(\overline{T})|$.
- (c) $\overline{T} \sim_0 \mathbb{A}^n$.
- (d) $(\overline{T}, 0)$ is an algebraic rational cell.
- (e) $(M, 0)$ is an algebraic rational cell.
- (f) $\dim M = |\mathcal{R}_1|$.

As in [R2], a reductive monoid M with zero element is called *quasismooth* if, for any minimal non-zero idempotent $e \in E(M)$, M_e satisfies the conditions of Theorem 5.5. The second main result of Section 5 concerns projective group embeddings $\mathbb{P}_\epsilon(M)$. The outcome is an extension of [G1, Theorem 7.4] to equivariant Chow groups.

Theorem (5.9, 5.10). *Let M be a reductive monoid with zero. If M is quasismooth, then the projective group embedding $\mathbb{P}_\epsilon(M)$ is (algebraically) \mathbb{Q} -filtrable. Furthermore, if $\mathbb{k} = \mathbb{C}$, then the corresponding equivariant cycle map is a natural isomorphism.*

In Section 6 we pair our ‘‘homological’’ theory with a natural ‘‘cohomology’’ theory on singular T -varieties: equivariant *operational* Chow groups. The motivation for this is Brion's characterization of Poincaré duality (in equivariant cohomology), in terms of Betti numbers and equivariant multiplicities [Br5]. In the setup of operational Chow groups, there is also an equivariant Poincaré duality map $\mathcal{P}_T : \text{op}A_T^k(X) \rightarrow A_{n-k}^T(X)$, $z \mapsto z \cap [X]$, for any T -scheme X of pure dimension n . This is an isomorphism if X is smooth. Combining Section 4 with [G3] yields some conditions for Poincaré duality on \mathbb{Q} -filtrable T -linear varieties. This is an extension of [Br5, Theorem 4.1] to operational Chow groups.

Theorem (6.8). *Let X be a complete equidimensional T -variety with isolated fixed points. Suppose that (a) X is \mathbb{Q} -filtrable, and (b) X satisfies the strong T -equivariant Kronecker duality. Then the following conditions are equivalent.*

- (i) X satisfies Poincaré duality.
- (ii) X satisfies T -equivariant Poincaré duality.
- (iii) The Chow homology Betti numbers of X satisfy $b_q(X) = b_{n-q}(X)$ for $0 \leq q \leq n$, and all equivariant multiplicities are nonzero.

If any of these conditions holds, then all equivariant multiplicities are in fact inverses of polynomial functions.

Put in perspective, the previous theorem asserts that \mathbb{Q} -filtrable projective T -linear varieties resemble the equivariantly formal spaces of [GKM], from the viewpoint of equivariant Chow cohomology.

Finally, in the last section of this article, we compare the two notions of \mathbb{Q} -filtrable varieties, the algebraic one (Section 4) and the topological one [G1]. We do so in the case of complex spherical varieties. Let G be a connected reductive group. Recall that a normal G -variety X is called *spherical* if a Borel subgroup B of G has a dense orbit in X . Then it is known that G and B have finitely many orbits in X . It follows that X contains only finitely many fixed points of a maximal torus $T \subset B$, see e.g. [Ti]. These features make spherical varieties especially suitable for applying the techniques of this paper. The main result of Section 7 is as follows.

Theorem (7.2, 7.3). *Let X be a spherical G -variety with an attractive T -fixed point x . Let X_x be the unique open affine T -stable neighborhood of x . If X is rationally smooth at x , then (X_x, x) is an algebraic rational cell. More generally, if X has a topological \mathbb{Q} -filtration, then this filtration is also an algebraic \mathbb{Q} -filtration. In this case, the cycle maps $cl_X : A_*(X) \rightarrow H_*(X)$ and $cl_X^T : A_*^T(X) \rightarrow H_*^T(X)$ are isomorphisms; and there is an isomorphism of free S -modules $\text{op}A_T^*(X) \simeq H_T^*(X)$.*

Acknowledgments. The research in this paper was done during my visits to the Institute des Hautes Études Scientifiques (IHES) and the Max-Planck-Institut für Mathematik (MPIM). I am deeply grateful to both institutions for their support, outstanding hospitality, and excellent working conditions. A very special thank you goes to Michel Brion for the productive meeting we had at IHES, from which this paper received much inspiration. I would also like to thank the support that I received, as a postdoctoral fellow, from Sabancı Üniversitesi and the Scientific and Technological Research Council of Turkey (TÜBİTAK), Project No. 112T233.

2. PRELIMINARIES

Conventions and notation. Throughout this paper, we work over an algebraically closed field \mathbb{k} of arbitrary characteristic (unless stated otherwise). All schemes and algebraic groups are assumed to be defined over \mathbb{k} . By a scheme we mean a separated scheme of finite type. A variety is a reduced scheme. Observe that varieties need not be irreducible. A subvariety is a closed subscheme which is a variety. A point on a scheme will always be a closed point.

We denote by T an algebraic torus. A scheme X provided with an algebraic action of T is called a T -scheme. For a T -scheme X , we denote by X^T the fixed point subscheme and by $i_T : X^T \rightarrow X$ the natural inclusion. If H is a closed subgroup of T , we similarly denote by $i_H : X^H \rightarrow X$ the inclusion of the fixed point subscheme. When comparing X^T and X^H we write $i_{T,H} : X^T \rightarrow X^H$ for the natural (T -equivariant) inclusion. For a scheme

X , the dimension of the local ring of X at x is denoted $\dim_x X$. We denote by Δ the character group of T , and by S the symmetric algebra over \mathbb{Q} of the abelian group Δ . We denote by \mathcal{Q} the quotient field of S . Equivariant Chow groups and equivariant operational Chow groups are considered with rational coefficients.

In this paper, torus actions are assumed to be *locally linear*, i.e. the schemes we consider are covered by invariant affine open subsets upon which the action is linear. This assumption is fulfilled e.g. by T -stable subschemes of normal T -schemes [Su].

2.1. The Bialynicki-Birula decomposition. The results in this subsection are due to Bialynicki-Birula [B1, B2] (in the smooth case) and Konarski [Ko] (in the general case). For our purposes, it suffices to consider the case of torus actions with isolated fixed points.

Let T be an algebraic torus. Let X be a T -scheme with isolated fixed points. Then X^T is finite and we write $X^T = \{x_1, \dots, x_m\}$. Recall that a one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow T$ is called *generic* if $X^{\mathbb{G}_m} = X^T$, where \mathbb{G}_m acts on X via λ . Generic one-parameter subgroups always exist, due to local linearity of the action. Now fix a generic one-parameter subgroup λ of T . For each i , define the subset

$$X_+(x_i, \lambda) = \{x \in X \mid \lim_{t \rightarrow 0} \lambda(t) \cdot x = x_i\}.$$

Then $X_+(x_i, \lambda)$ is a locally closed T -invariant subscheme of X . The (disjoint) union of the $X_+(x_i, \lambda)$'s might not cover all of X , but when it does (e.g., when X is complete), the decomposition $\{X_+(x_i, \lambda)\}_{i=1}^m$ is called the Bialynicki-Birula decomposition, or *BB-decomposition*, of X associated to λ . Each $X_+(x_i, \lambda)$ is called a *cell* of the decomposition. Usually the *BB-decomposition* of a complete T -scheme is not a Whitney stratification; that is, it may happen that the closure of a cell is not a union of cells, even when the scheme is assumed to be smooth. For instance, see [B2, Example 1].

Definition 2.1. Let X be a T -scheme with finitely many fixed points. Let $\{X_+(x_i, \lambda)\}_{i=1}^m$ be the BB-decomposition associated to some generic one-parameter subgroup λ of T . The decomposition $\{X_+(x_i, \lambda)\}$ is said to be *filtrable* if there exists a finite increasing sequence $\Sigma_0 \subset \Sigma_1 \subset \dots \subset \Sigma_m$ of T -invariant closed subschemes of X such that:

- a) $\Sigma_0 = \emptyset$, $\Sigma_m = X$,
- b) $\Sigma_j \setminus \Sigma_{j-1}$ is a cell of the decomposition $\{X_+(x_i, \lambda)\}$, for each $j = 1, \dots, m$.

We will refer to Σ_j as the *j -th filtered piece* of X . In this context, it is common to say that X is *filtrable*. If, moreover, the cells $X_+(x_i, \lambda)$ are isomorphic to affine spaces \mathbb{A}^{n_i} , then X is called *T -cellular*.

Theorem 2.2 ([B1], [B2]). *Let X be a complete T -scheme with isolated fixed points, and let λ be a generic one-parameter subgroup. If X admits an ample T -linearized invertible sheaf, then the associated BB-decomposition $\{X_+(x_i, \lambda)\}$ is filtrable. Further, if X is smooth, then X is T -cellular. \square*

2.2. Review of equivariant Chow groups. Localization theorem.

Let X be a T -scheme of dimension n (not necessarily equidimensional). Let V be a finite dimensional T -module, and let $U \subset V$ be an invariant open subset such that a principal bundle quotient $U \rightarrow U/T$ exists. Then T acts freely on $X \times U$ and the quotient scheme $X_T := (X \times U)/T$ exists. Following Edidin and Graham [EG], we define the i -th *equivariant Chow group* $A_i^T(X)$ by $A_i^T(X) := A_{i+\dim U - \dim T}(X)$, if $V \setminus U$ has codimension more than $n - i$. Such pair (V, U) always exist, and the definition is independent of the choice of (V, U) , see [EG]. Finally, set $A_*^T(X) = \bigoplus_i A_i^T(X)$. If X is a T -scheme, and $Y \subset X$ is a T -stable closed subscheme, then Y defines a class $[Y]$ in $A_*^T(X)$. If X is smooth, then so is X_T , and $A_*^T(X)$ admits an intersection pairing; in this case, denote by $A_T^*(X)$ the corresponding ring graded by codimension. The equivariant Chow ring $A_T^*(pt)$ identifies to S , and $A_*^T(X)$ is a S -module, where Δ acts on $A_*^T(X)$ by homogeneous maps of degree -1 . This module structure is induced by pullback through the flat map $p_{X,T} : X_T \rightarrow U/G$. Restriction to a fiber of $p_{X,T}$ gives $i^* : A_*^T(X) \rightarrow A_*(X)$. If X is complete, we denote by $\int_X(\alpha)$ the proper pushforward to a point of a class $\alpha \in A_*^T(X)$. See [EG] for details.

Next we state Brion's description [Br1] of the equivariant Chow groups in terms of invariant cycles. It also shows how to recover the usual Chow groups from equivariant ones.

Theorem 2.3. *Let X be a T -scheme. Then the S -module $A_*^T(X)$ is defined by generators $[Y]$ where Y is an invariant irreducible subvariety of X and relations $[\text{div}_Y(f)] - \chi[Y]$ where f is a rational function on Y which is an eigenvector of T of weight χ . Moreover, the map $A_*^T(X) \rightarrow A_*(X)$ vanishes on $\Delta A_*^T(X)$, and it induces an isomorphism*

$$A_*^T(X)/\Delta A_*^T(X) \rightarrow A_*(X).$$

□

The following is the localization theorem for equivariant Chow groups [Br1, Corollary 2.3.2] (cf. [G3, Proposition 2.15]).

Theorem 2.4. *Let X be a T -scheme, let $H \subset T$ be a closed subgroup, and let $i_H : X^H \rightarrow X$ be the inclusion of the fixed point subscheme. Then the induced morphism of equivariant Chow groups*

$$i_{H*} : A_*^T(X^H) \rightarrow A_*^T(X)$$

becomes an isomorphism after inverting finitely many characters of T that restrict non-trivially to H . □

Let X be a T -scheme. In many situations, Theorems 2.3 and 2.4 combined yield a relation between the dimensions of the \mathbb{Q} -vector spaces $A_*(X)$ and $A_*(X^T)$.

Lemma 2.5. *Let X be a T -scheme. If $A_*(X)$ is a finite-dimensional \mathbb{Q} -vector space, then the inequality $\dim_{\mathbb{Q}} A_*(X^T) \leq \dim_{\mathbb{Q}} A_*(X)$ holds. Furthermore, $\dim_{\mathbb{Q}} A_*(X^T) = \dim_{\mathbb{Q}} A_*(X)$ if and only if the S -module $A_*^T(X)$ is free.*

Proof. The degrees in $A_*^T(X)$ are at most the dimension of X , so by the graded Nakayama lemma [E, Exercise 4.6], the S -module $A_*^T(X)$ is finitely generated. The content of the corollary is now deduced from applying Lemma 2.6 and Remark 2.7 below to $M = A_*^T(X)$, taking into account that $\dim_{S/\mathfrak{m}}(M/\mathfrak{m}M) = \dim_{\mathbb{Q}}(A_*(X))$ (Theorem 2.3), $\dim_{\mathbb{Q}}(M \otimes_S \mathbb{Q}) = \dim_{\mathbb{Q}}(A_*^T(X^T) \otimes \mathbb{Q})$ (Theorem 2.4), and observing that this corresponds to $\dim_{\mathbb{Q}} A_*(X^T)$, since $A_*^T(X^T) = A_*(X^T) \otimes_{\mathbb{Q}} S$. \square

Lemma 2.6. *Let S be a Noetherian positively graded ring such that S_0 is a field (e.g. $S = A_T^*(pt)$). Let \mathfrak{m} be the unique graded maximal ideal and suppose M is a non-zero finitely generated, graded, S -module. Suppose further that S is an integral domain. Then M is a free S -module if and only if*

$$\dim_{S/\mathfrak{m}}(M/\mathfrak{m}M) = \dim_{\mathbb{Q}}(M \otimes_S \mathbb{Q}),$$

where \mathbb{Q} is the quotient field of S .

Proof. If M is free, then clearly the equation above holds. Conversely, denote by n the common value of the two sides of the equation above. By the graded Nakayama lemma, M has a system $\{x_1, \dots, x_n\}$ of homogeneous generators. Now the elements $x_j \otimes 1$, generate the vector space $M \otimes_S \mathbb{Q}$ over \mathbb{Q} . But as by hypothesis this space is of dimension n over \mathbb{Q} , the elements $x_j \otimes 1$ are linearly independent over \mathbb{Q} . It follows that the x_j are linearly independent over S and so they form a basis of M . \square

Remark 2.7. The proof of Lemma 2.6 shows that that if M is a finitely generated, graded, S -module, then the inequality

$$\dim_{S/\mathfrak{m}}(M/\mathfrak{m}M) \geq \dim_{\mathbb{Q}}(M \otimes_S \mathbb{Q})$$

holds, as we can refine the generating set $\{x_j \otimes 1\}$ to get a basis of $M \otimes \mathbb{Q}$.

An important class of schemes to which Lemma 2.5 applies is the class of T -linear schemes. These schemes have been studied in [Jann], [J], [JK], [To], [FMSS], and [G3]. Briefly, a T -linear scheme is a T -scheme that can be obtained by an inductive procedure starting with a finite dimensional T -representation, in such a way that the complement of a T -linear scheme equivariantly embedded in affine space is also a T -linear scheme, and any T -scheme which can be stratified as a finite disjoint union of T -linear schemes is a T -linear scheme. See [JK] or [G3] for details. It is known that if X is a T -linear scheme, then $A_*^T(X)_{\mathbb{Z}}$ is a finitely generated $S_{\mathbb{Z}}$ -module, and $A_*(X)_{\mathbb{Z}}$ is a finitely generated abelian group (e.g. [G3, Lemma 2.11]). Below are the concrete examples we are interested in. For a proof of items (i)-(iii), see [JK, Proposition 3.6], for item (iv) see [G3, Theorem 2.5].

Theorem 2.8. *Let T be an algebraic torus. Then the following hold:*

- (i) *A T -cellular scheme is T -linear.*
- (ii) *Every T -scheme with finitely many T -orbits is T -linear. In particular, a toric variety with dense torus T is T -linear.*
- (iii) *Assume $\text{char}(\mathbb{k}) = 0$. Let B be a connected solvable linear algebraic group with maximal torus T . Let X be a B -scheme. If B acts on X with finitely many orbits, then X is T -linear. In particular, spherical varieties are T -linear. \square*

2.3. Nondegenerate fixed points and equivariant multiplicities. Let X be a T -scheme. A fixed point $x \in X$ is called *nondegenerate* if all weights of T in the tangent space $T_x X$ are non-zero. A fixed point in a nonsingular T -variety is nondegenerate if and only if it is isolated. To study possibly singular schemes, Brion developed a notion of *equivariant multiplicity* at nondegenerate fixed points [Br1]. The main features of this concept are outlined below.

Theorem 2.9 ([Br1, Theorem 4.2]). *Let X be a T -scheme with an action of T , let $x \in X$ be a nondegenerate fixed point and let χ_1, \dots, χ_n be the weights of $T_x X$ (counted with multiplicity).*

- (i) *There exists a unique S -linear map*

$$e_{x,X} : A_*^T(X) \longrightarrow \frac{1}{\chi_1 \cdots \chi_n} S$$

such that $e_{x,X}[x] = 1$ and that $e_{x,X}[Y] = 0$ for any T -invariant irreducible subvariety $Y \subset X$ which does not contain x .

- (ii) *For any T -invariant irreducible subvariety $Y \subset X$, the rational function $e_{x,X}[Y]$ is homogeneous of degree $-\dim(Y)$ and it coincides with $e_{x,Y}[Y]$.*

- (iii) *The point is nonsingular in X if and only if $e_x[X] = \frac{1}{\chi_1 \cdots \chi_n}$. \square*

For any T -invariant irreducible subvariety $Y \subset X$, we set $e_{x,X}[Y] := e_x[Y]$, and we call $e_x[Y]$ the *equivariant multiplicity of Y at x* .

Proposition 2.10 ([Br1, Corollary 4.2]). *Let X be a T -scheme such that all fixed points in X are nondegenerate, and let $\alpha \in A_*^T(X)$. Then we have in $A_*^T(X) \otimes_S \mathcal{Q}$:*

$$\alpha = \sum_{x \in X^T} e_x(\alpha)[x].$$

\square

Next we consider a special class of nondegenerate fixed points. Let X be a T -variety. Call a fix point $x \in X$ *attractive* if all weights in the tangent space $T_x X$ are contained in some open half-space of $\Delta_{\mathbb{R}} = \Delta \otimes_{\mathbb{Z}} \mathbb{R}$, that is, some one-parameter subgroup of T acts on $T_x X$ with positive weights only. Below is a characterization.

Theorem 2.11 ([Br3, Proposition A2]). *Let X be a T -variety with a fixed point x . The following conditions are equivalent:*

- (i) x is attractive.
- (ii) *There exists a one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow T$ such that, for all y in a neighborhood of x , we have $\lim_{t \rightarrow 0} \lambda(t)y = x$.*

Moreover, if (i) or (ii) holds, then x admits a unique open affine T -stable neighborhood in X , denoted X_x , and X_x admits a closed equivariant embedding into $T_x X$. \square

Let X be a T -variety with an attractive fixed point x . Denote by χ_1, \dots, χ_n the weights of $T_x X$. Let Δ^* be the lattice of one-parameter subgroups of T , and let $\Delta_{\mathbb{R}}^*$ be the associated real vector space. Notice that the one-parameter subgroups λ satisfying Theorem 2.11 (ii) form a rational polyhedral cone $\sigma_x \subset \Delta_{\mathbb{R}}^*$ with non-empty interior σ_x^0 , by setting

$$\sigma_x := \{\lambda \in \Delta_{\mathbb{R}}^* \mid \langle \lambda, \chi_i \rangle \geq 0 \text{ for } 1 \leq i \leq n\}.$$

It follows from Theorem 2.11 that X_x equals $X_+(x, \lambda)$ for any $\lambda \in \sigma_x^0$.

Proposition 2.12 ([Br1, Proposition 4.4]). *Notation being as above, the rational function $e_x[X]$, viewed as a rational function on $\Delta_{\mathbb{R}}^*$, is defined at λ and its value is the multiplicity of the algebra of regular functions on X_x graded via the action of λ . In particular, $e_x[X]$ is non-zero.* \square

2.4. Local study. Some inequalities relating Chow groups and fixed point loci. Let X be an affine T -variety with an attractive fixed point x . It follows from Proposition 2.12 that $X = X_+(x, \lambda)$ for any $\lambda \in \sigma_x^0$. Also, $\{x\}$ is the unique closed T -orbit in X , and X admits a closed T -equivariant embedding into $T_x X$. Observe that $\dim_x X = \dim X$, because x is contained in every irreducible component of X .

Choose $\lambda \in \sigma_x^0$. Then all the weights of the \mathbb{G}_m -action on $T_x X$ via λ are positive. Hence the geometric quotient

$$\mathbb{P}_\lambda(X) := (X \setminus \{x\})/\mathbb{G}_m$$

exists and is a projective variety. In fact, it is a closed subvariety of the weighted projective space $\mathbb{P}(T_x X)$. On the other hand, by [Br3, Proposition A3], there exists a \mathbb{G}_m -module V and a finite equivariant surjective morphism $\pi : X \rightarrow V$ such that $\pi^{-1}(0) = \{x\}$ (as a set). This allows to estimate the size of the Chow groups of $\mathbb{P}_\lambda(X)$ in various cases.

Lemma 2.13. *In the situation above, π induces a surjection*

$$\pi_* : A_k(\mathbb{P}_\lambda(X)) \rightarrow A_k(\mathbb{P}(V))$$

for all $k \geq 0$. Consequently, $A_k(\mathbb{P}_\lambda(X)) \neq 0$ if $0 \leq k \leq \dim(X)$, and $A_k(\mathbb{P}_\lambda(X)) = 0$ otherwise. \square

Remark 2.14. Clearly, $\pi_* : A_*(X) \rightarrow A_*(V)$ is also surjective. Observe that if X is equidimensional and d is the degree of π , then

$$e'_x[X] = d \cdot e'_0[V],$$

where $e'_x[X]$ (resp. $e'_0[V]$) is the \mathbb{G}_m -equivariant multiplicity of X at x (resp. of V at 0) [Br1, Proposition 4.3].

Now assume that $\dim_{\mathbb{Q}} A_*(\mathbb{P}_{\lambda}(X)) < \infty$. We record below a few elementary inequalities:

- (1) $\dim A_*(\mathbb{P}_{\lambda}(X)) \geq \dim A_*(\mathbb{P}_{\lambda}(X)^T)$, by Lemma 2.5.
- (2) Notice that $\mathbb{P}_{\lambda}(X)^T = \bigsqcup_H \mathbb{P}_{\lambda}(X^H)$, where the union runs over all codimension-one subtori of T . In fact, by linearity of the action, there is only a finite collection of codimension-one subtori, say H_1, \dots, H_r , for which $X^{H_j} \neq X^T$. Thus

$$\dim_{\mathbb{Q}} A_*(\mathbb{P}(X)^T) = \sum_{H_j} \dim_{\mathbb{Q}} A_*(\mathbb{P}(X^{H_j})).$$

- (3) Let H_j be as in (2). We may also assume that x is an attractive fixed point of X^{H_j} , for the action of $\mathbb{G}_m \simeq T/H_j$. Hence, as in Lemma 2.13, there is a T -equivariant finite surjective map $\pi_j : X^{H_j} \rightarrow V_j$, where V_j is some T -module with a trivial action of H_j . Thus $\dim_{\mathbb{Q}} A_*(\mathbb{P}(X^{H_j})) \geq \dim_{\mathbb{Q}} A_*(\mathbb{P}(V_j))$, which in turn yields

$$\sum_{j=1}^r \dim_{\mathbb{Q}} A_*(\mathbb{P}(X^{H_j})) \geq \sum_{j=1}^r \dim_{\mathbb{Q}} A_*(\mathbb{P}(V_j)).$$

Equality holds if and only if the π_j 's induce isomorphisms on the Chow groups.

- (4) Since each $\mathbb{P}(V_j)$ in (3) is a projective space, we get

$$\sum_{j=1}^r \dim_{\mathbb{Q}} A_*(\mathbb{P}(V_j)) = \sum_{j=1}^r \dim V_j = \sum_{j=1}^r \dim X^{H_j},$$

where the last equality stems from the fact that each π_j is finite and surjective.

- (5) Because x is an attractive fixed point, we have

$$\sum_{j=1}^r \dim X^{H_j} \geq \dim X,$$

by [Br3, Theorem 1.4]. The equality holds if and only if there exists a T -equivariant finite surjective map $\pi : X \rightarrow V$, where V is a T -module with zero as an attractive fixed point. This is done by synchronizing the maps π_j from (3), cf. [Br3, Proof of Theorem 1.2].

Combining items (1) to (5), we obtain the chain of inequalities

$$\dim_{\mathbb{Q}} A_*(\mathbb{P}_{\lambda}(X)) \geq \sum_{j=1}^r \dim_{\mathbb{Q}} A_*(\mathbb{P}(X^{H_j})) \geq \sum_{j=1}^r \dim_{\mathbb{Q}} A_*(\mathbb{P}(V_j)) = \sum_{j=1}^r \dim X^{H_j} \geq \dim X.$$

Proposition 2.15. *Notation as above, if $\dim_{\mathbb{Q}} A_*(\mathbb{P}_{\lambda}(X)) = \dim X$, then the chain displayed above is a chain of equalities, and the maps π_j and π from (3) and (5) induce isomorphisms*

$$\begin{aligned}\pi_{j*} : A_k(\mathbb{P}(X^{H_j})) &\xrightarrow{\sim} A_k(\mathbb{P}(V_j)), \\ \pi_* : A_k(\mathbb{P}(X)) &\xrightarrow{\sim} A_k(\mathbb{P}(V)),\end{aligned}$$

for all j and k . □

Put in perspective, this result is our motivation for the material in the next section.

3. ALGEBRAIC RATIONAL CELLS

This section is devoted to the study of our main technical tool: algebraic rational cells.

We thank M. Brion for leading us to the following definition.

Definition 3.1. Let X be an affine T -variety with an attractive fixed point x , and let $n = \dim X$. We say that (X, x) , or simply X , is an *algebraic rational cell* if and only if, for some $\lambda \in \sigma_x^0$, we have

$$A_k(\mathbb{P}_{\lambda}(X)) = \begin{cases} \mathbb{Q} & \text{if } 0 \leq k \leq n-1, \\ 0 & \text{otherwise.} \end{cases}$$

We abbreviate this condition by writing $A_*(\mathbb{P}_{\lambda}(X)) \simeq A_*(\mathbb{P}^{n-1})$.

Algebraic rational cells are such T -varieties for which Proposition 2.15 holds. In principle, Definition 3.1 depends on a particular choice of $\lambda \in \sigma_x^0$. But, as we shall see next, it is independent of λ : if $A_*(\mathbb{P}_{\lambda}(X)) \simeq A_*(\mathbb{P}^{n-1})$ holds for some $\lambda \in \sigma_x^0$, then it holds for all $\lambda \in \sigma_x^0$.

Lemma 3.2. *Let X be an affine T -variety with an attractive fixed point x , and let $n = \dim X$. Then (X, x) is an algebraic rational cell if and only if*

$$A_k(X) = \begin{cases} \mathbb{Q} & \text{if } k = n \\ 0 & \text{if } k \neq n. \end{cases}$$

In particular, if (X, x) is an algebraic rational cell, then it is irreducible.

Proof. Let \mathbb{G}_m act on X via λ . Recall that we have a short exact sequence

$$0 \rightarrow A_*^{\mathbb{G}_m}(x) \rightarrow A_*^{\mathbb{G}_m}(X) \rightarrow A_*^{\mathbb{G}_m}(X \setminus \{x\}) \rightarrow 0,$$

which stems from the localization theorem (Theorem 2.4). As in Lemma 2.13, there exists a \mathbb{G}_m -equivariant finite surjective map $\pi : X \rightarrow \mathbb{A}^n$ such that $\pi^{-1}(0) = x$. This map induces the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & A_*^{\mathbb{G}_m}(x) & \xrightarrow{i_*} & A_*^{\mathbb{G}_m}(X) & \xrightarrow{j_*} & A_*^{\mathbb{G}_m}(X \setminus \{x\}) \longrightarrow 0 \\ & & \downarrow \pi_* & & \downarrow \pi_* & & \downarrow \pi_* \\ 0 & \longrightarrow & A_*^{\mathbb{G}_m}(0) & \xrightarrow{i_*} & A_*^{\mathbb{G}_m}(\mathbb{A}^n) & \xrightarrow{j_*} & A_*^{\mathbb{G}_m}(\mathbb{A}^n \setminus \{0\}) \longrightarrow 0 \end{array}$$

where all vertical maps are surjective. Moreover, the vertical map on the right represents

$$\pi_* : A_*(\mathbb{P}(X)) \rightarrow A_*(\mathbb{P}^{n-1}),$$

because $A_*^{\mathbb{G}^m}(X \setminus \{x\}) \simeq A_*(\mathbb{P}(X))$, by [EG, Theorem 3]. Since the left vertical map is clearly an isomorphism, it follows that the right vertical map is an isomorphism if and only if so is the middle one. But, in view of Lemma 2.5, the latter happens if and only if $A_*(X) \simeq A_*(\mathbb{A}^n) \simeq \mathbb{Q}$. This yields the first assertion of the lemma.

For the second assertion, observe that the condition $A_*(X) \simeq \mathbb{Q}$ implies that X has a unique irreducible component, say Y , of dimension $n = \dim X$. (Note that all irreducible components of X contain the fixed point x .) To finish the argument, let us suppose, by contradiction, that X has another irreducible component of dimension $e < n$. It suffices to consider the case $X = Y \cup Z$, with Z an irreducible component of dimension $e < n$. Now the sequence

$$A_e(Y) \rightarrow A_e(X) \rightarrow A_e(Z \setminus Y) \rightarrow 0$$

yields $A_e(X) \neq 0$, because $A_e(Z \setminus Y) = \mathbb{Q}$ (as $Z \setminus Y$ is open and Z is irreducible). So $\dim_{\mathbb{Q}} A_*(X) \geq 2$. But this is a contradiction. \square

The previous lemma hints to a more general structural property of algebraic rational cells with respect to the T -action.

Proposition 3.3. *Let X be an affine T -variety with an attractive fixed point x , and let $\lambda \in \sigma_x^0$. Let $n = \dim X$. Then the following conditions are equivalent.*

- (i) $A_*(\mathbb{P}_\lambda(X)) \simeq A_*(\mathbb{P}^{n-1})$.
- (ii) $A_*(X) \simeq A_*(\mathbb{A}^n)$.
- (iii) $A_*^T(X) \simeq A_*^T(pt) = S$.
- (iv) $A_*^T(\mathbb{P}_\lambda(X)) \simeq A_*^T(\mathbb{P}^{n-1}) \otimes_{\mathbb{Q}} S$

Proof. The equivalence (i) \Leftrightarrow (ii) follows from Lemma 3.2.

The equivalence (ii) \Leftrightarrow (iii) follows from Lemma 2.5.

The implication (iv) \Rightarrow (i) is deduced from Theorem 2.3 and Lemma 2.13.

Finally, we dispose of the direction (i) \Rightarrow (iv). Recall that (i) yields the existence of a T -equivariant finite surjective morphism $\pi : X \rightarrow V$, such that the induced map $\pi_* : A_*(\mathbb{P}_\lambda(X)) \rightarrow A_*(\mathbb{P}_\lambda(V))$ is an isomorphism (Proposition 2.15). By the graded Nakayama lemma, the corresponding map $\widetilde{\pi}_* : A_*^T(\mathbb{P}_\lambda(X)) \rightarrow A_*^T(\mathbb{P}_\lambda(V))$ is surjective. We claim that $\widetilde{\pi}_*$ is also injective (hence an isomorphism). Indeed, choose a basis z_1, \dots, z_n of $A_*(\mathbb{P}_\lambda(V))$. Now identify that basis with a basis of $A_*(\mathbb{P}_\lambda(X))$, via π_* , and lift it to a generating system of the S -module $A_*^T(\mathbb{P}_\lambda(X))$. This generating system is a basis, since its image under $\widetilde{\pi}_*$ is a basis of $A_*^T(\mathbb{P}_\lambda(V))$. \square

Next, we exhibit some additional features of algebraic rational cells. The result is an algebraic counterpart of [Br2, Theorem 18].

Theorem 3.4. *Let X be an affine T -variety with an attractive fixed point x . If (X, x) is an algebraic rational cell, then the following equivalent conditions hold:*

- (i) *For each subtorus $H \subset T$ of codimension one, (X^H, x) is an algebraic rational cell, and $\dim X = \sum_H \dim X^H$ (sum over all subtori of codimension one).*
- (ii) *For each subtorus $H \subset T$ of codimension one, (X^H, x) is an algebraic rational cell, and*

$$e_x[X] = d \prod_H e_x[X^H],$$

where d is a positive rational number. If moreover each X^H is smooth, then d is an integer.

Proof. There is only a finite collection of codimension one subtori, say H_1, \dots, H_r , for which $X^{H_j} \neq X^T$. Thus statement (i) follows at once from Proposition 2.15.

It only remains to show that, in the situation at hand, the equality $\dim X = \sum_{H_j} \dim X^{H_j}$ is equivalent to $e_x[X] = d \prod_{H_j} e_x[X^{H_j}]$. Nevertheless, in light of Proposition 2.15, this equivalence is obtained arguing exactly as in [Br2, Theorem 18]. Indeed, since X and all the X^{H_j} 's are irreducible (Lemma 3.2), assuming $e_x[X] = d \prod_{H_j} e_x[X^{H_j}]$ yields $\dim X = \sum_{H_j} \dim X^{H_j}$, by Theorem 2.9 (ii). Conversely, if the latter equation holds, there exists a T -equivariant finite surjective map $\pi : X \rightarrow V$ (Proposition 2.15). Thus $e_x[X] = de_0[V]$, where $d = \deg \pi$. In turn, $de_0[V] = d \prod_{H_j} e_0[V_j]$, because V is a T -module (Theorem 2.9 (iii)). Finally, the last expression identifies to $\frac{d}{\prod_j d_j} \prod_j e_x[X^{H_j}]$, where $d_j = \deg \pi_j$, and π_j is as in Proposition 2.15. \square

In general, it is not true that properties (i) or (ii) of Theorem 3.4 characterize algebraic rational cells. Here is an example, cf. [Br3, Remark 1.4].

Example 3.5. Let X be the hypersurface of \mathbb{A}^5 with equation $x^2 + yz + xtw = 0$. Note that X is irreducible, with singular locus $x = y = z = tw = 0$, a union of two lines meeting at the origin. Now consider the $\mathbb{G}_m \times \mathbb{G}_m$ -action on \mathbb{A}^5 given by $(u, v) \cdot (x, y, z, t, w) := (u^2v^2x, u^3vy, uv^3z, u^2t, v^2w)$. Then the origin of \mathbb{A}^5 is an attractive fixed point, X is T -stable of dimension four and X contains four closed irreducible T -stable curves, namely, the coordinate lines except for the x -axis. So X satisfies condition (i) of Theorem 3.4. Nevertheless, $(X, 0)$ is not an algebraic rational cell. To see this, consider the \mathbb{G}_m -action on \mathbb{A}^5 given by $u \cdot (x, y, z, t, w) := (x, uy, u^{-1}z, t, w)$. Then X is \mathbb{G}_m -stable and $X^{\mathbb{G}_m}$ is defined by $y = z = x^2 + xtw = 0$. Thus $X^{\mathbb{G}_m}$ is reducible at the origin. In fact $A_*(X^{\mathbb{G}_m}) = \mathbb{Q} \oplus \mathbb{Q}$ (as it consists of the union of two copies of \mathbb{A}^2). Thus $\dim_{\mathbb{Q}} A_*(X^{\mathbb{G}_m}) = 2$, and so $\dim_{\mathbb{Q}} A_*(X) \geq 2$, by Lemma 2.5. Therefore, in view of Lemma 3.2, $(X, 0)$ is not an algebraic rational cell.

Example 3.6 (Smooth rational cells). Let X be an affine T -variety with an attractive fixed point x . If X is smooth at x , then $X \simeq T_x X$, T -equivariantly. Thus $(X, x) \simeq (T_x X, 0)$ is an algebraic rational cell. This agrees with the fact that $\mathbb{P}_\lambda(T_x X)$ is a weighted projective space.

Remark 3.7. Let $\mathbb{k} = \mathbb{C}$. In general, there is no immediate relation between algebraic rational cells and rational cells. Indeed, if (X, x) is an algebraic rational cell, then using Proposition 2.15 one checks that the cycle map $cl : A_*(\mathbb{P}_\lambda(X)) \rightarrow H_{2*}(\mathbb{P}_\lambda(X))$ is injective. But then $H_{\text{odd}}(\mathbb{P}_\lambda(X))$ could be non-zero. Conversely, if (X, x) is a rational cell, then $\mathbb{P}_\lambda(X)$ is a rational cohomology complex projective space, and one checks that the cycle map is surjective. In some important cases, however, stronger connections exist (Sections 5 and 7).

We conclude this section by computing equivariant multiplicities of algebraic rational cells. Recall that a character χ of T is called *singular* if $X^{\ker(\chi)} \neq X^T$.

Corollary 3.8. *Let X be an irreducible T -variety with attractive fixed point x . Let X_x be the unique open affine T -stable neighborhood of x . If (X_x, x) is an algebraic rational cell, then the following hold:*

(i) $e_x[X]$ is the inverse of a polynomial. In fact,

$$e_x[X] = \frac{d}{\chi_1 \cdots \chi_r},$$

where the χ_i 's are singular characters, and d is a positive rational number.

(ii) Additionally, if the number of closed irreducible T -stable curves through x is finite, say $\ell(x)$, then $\dim X = \ell(x)$. Furthermore, if $\chi_1, \dots, \chi_{\ell(x)}$ are the characters associated to the closed irreducible T -invariant curves through x , then

$$e_x[X] = \frac{d'}{\chi_1 \cdots \chi_{\ell(x)}},$$

where d' is a positive rational number.

Proof. Replacing X by X_x we may assume that X is affine. Then (i) follows at once from Theorem 3.4 and its proof. As for (ii) simply use Theorem 3.4 and Corollary 3.8, to adapt the argument of [Br3, Corollary 1.4.2] and [G1, Corollary 5.6]. \square

In general, if X is an affine T -variety with attractive fixed point x , and $\ell(x)$ as above is finite, then $\dim_x X \leq \ell(x)$ [Br3, Corollary 1.4.2].

4. \mathbb{Q} -FILTRABLE VARIETIES AND EQUIVARIANT CHOW GROUPS

We aim at an inductive description of the equivariant Chow groups of filtrable T -varieties in the case when the cells are all algebraic rational cells. Our findings provide purely algebraic analogues of the topological results of [G1].

Definition 4.1. Let X be a T -variety. We say that X is \mathbb{Q} -filtrable if the following hold:

- (1) the fixed point set X^T is finite, and
- (2) there exists a generic one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow T$ for which the associated BB -decomposition of X is filtrable (Definition 2.1) and consists of T -invariant *algebraic rational cells*.

Observe that the fixed points $x_j \in X^T$ need not be attractive in X , but they are so in their particular algebraic rational cells $X_+(x_j, \lambda)$. The following technical result will be of importance in the sequel.

Lemma 4.2. *If (X, x) is an algebraic rational cell, then the equivariant multiplicity morphism $e_{X,x} : A_*^T(X) \rightarrow \mathcal{Q}$ is injective.*

Proof. By [Br1, Proposition 4.1] that the map $i_* : A_*^T(x) \rightarrow A_*^T(X)$ is injective. Moreover, the image of i_* contains $\chi_1 \cdots \chi_n A_*^T(X)$, where χ_i are the T -weights of $T_x X$. Next, recall that e_x is defined as follows: given $\alpha \in A_*^T(X)$, we can form the product $\chi_1 \cdots \chi_n \alpha$. Thus, there exists $\beta \in S$ such that $i_*(\beta) = \chi_1 \cdots \chi_n \alpha$. Now let $e_x(\alpha) = \frac{\beta}{\chi_1 \cdots \chi_n}$. Since $A_*^T(X)$ is S -free (Proposition 3.3), it is clear from the construction that e_x is injective. \square

Let X be a \mathbb{Q} -filtrable T -variety. Then, by assumption, there is a closed algebraic rational cell $F = X_+(x_1, \lambda)$ (using the order of fixed points induced by the filtration, cf. Definition 2.1). Moreover $U = X \setminus F$ is also \mathbb{Q} -filtrable. We now proceed to describe $A_*^T(X)$ in terms of $A_*^T(F)$ and $A_*^T(U)$. Let $j_F : F \rightarrow X$ and $j_U : U \rightarrow X$ denote the inclusion maps.

Proposition 4.3. *Notation being as above, the maps $j_{F*} : A_*^T(F) \rightarrow A_*^T(X)$ and $j_U^* : A_*^T(X) \rightarrow A_*^T(U)$ fit into the exact sequence*

$$0 \rightarrow A_*^T(F) \rightarrow A_*^T(X) \rightarrow A_*^T(U) \rightarrow 0.$$

Proof. It is well-known that the sequence

$$A_*^T(F) \xrightarrow{j_{F*}} A_*^T(X) \xrightarrow{j_U^*} A_*^T(U) \longrightarrow 0$$

is exact. Thus it suffices to show that j_{F*} is injective. But this follows easily from the factorization $e_{x,F} = e_{x,X} \circ j_{F*}$. Indeed, since $e_{x,F}$ is injective (Lemma 4.2), so is j_{F*} . \square

Arguing by induction on the length of the filtration leads to the following result.

Corollary 4.4. *Let X be a \mathbb{Q} -filtrable T -variety. Then the T -equivariant Chow group of X is a free module of rank $|X^T|$. In fact, it is freely generated by the classes of the closures of the cells $X_+(x_i, \lambda)$. Consequently, $A_*(X)$ is also freely generated by the classes of the cell closures $\overline{X_+(x_i, \lambda)}$. \square*

If X is a \mathbb{Q} -filtrable variety, then each filtered piece Σ_i is also \mathbb{Q} -filtrable, and so Corollary 4.4 applies at each step of the filtration. Our approach, based on equivariant multiplicities, is more flexible than the general approach which compares (equivariant) Chow groups with (equivariant) homology via the (equivariant) cycle map. This flexibility will be illustrated in the next sections.

5. APPLICATIONS TO EMBEDDING THEORY

We now furnish our theory with its first set of examples: \mathbb{Q} -filtrable embeddings of reductive groups. We show that the notion of algebraic rational cell is well adapted to the study of group embeddings.

Further notation. We denote by G a connected reductive linear algebraic group with Borel subgroup B and maximal torus $T \subset B$. We denote by W the Weyl group of (G, T) .

An affine algebraic monoid M is called *reductive* if it is irreducible, normal, and its unit group is a reductive algebraic group. See [R1] for details. Let M be a reductive monoid with zero and unit group G . We denote by $E(M)$ the idempotent set of M , that is, $E(M) = \{e \in M \mid e^2 = e\}$. Likewise, we denote by $E(\overline{T})$ the idempotent set of the associated affine torus embedding \overline{T} . One defines a partial order on $E(\overline{T})$ by declaring $f \leq e$ if and only if $fe = f$. Denote by $\Lambda \subset E(\overline{T})$, the cross section lattice of M . The Renner monoid $\mathcal{R} \subset M$ is a finite monoid whose group of units is W and contains $E(\overline{T})$ as idempotent set. Any $x \in \mathcal{R}$ can be written as $x = fu$, where $f \in E(\overline{T})$ and $u \in W$. Given $e \in E(\overline{T})$, we write $C_W(e)$ for the centralizer of e in W . Denote by \mathcal{R}_k the set of elements of rank k in \mathcal{R} , that is, $\mathcal{R}_k = \{x \in \mathcal{R} \mid \dim Tx = k\}$. Analogously, one has $\Lambda_k \subset \Lambda$ and $E_k \subset E(\overline{T})$. For $e \in E(M)$, set $M_e := \{g \in G \mid ge = eg = e\}$. Then M_e is an irreducible, normal reductive monoid with e as its zero element [Br4].

5.1. Group embeddings. A normal irreducible variety X is called an *embedding* of G , or a *group embedding*, if X is a $G \times G$ -variety containing an open orbit isomorphic to G . Due to the Bruhat decomposition, group embeddings are spherical $G \times G$ -varieties. Substantial information about the topology of a group embedding is obtained by restricting one's attention to the induced action of $T \times T$. When $G = B = T$, we get back the notion of toric varieties. Group embeddings are classified as follows.

(I) *Affine case:* Let M be a reductive monoid with unit group G . Then $G \times G$ -acts naturally on M via $(g, h) \cdot x = gxh^{-1}$. The orbit of the identity is $G \times G / \Delta(G) \simeq G$. Thus M is an affine embedding of G . Remarkably, by a result of Rittatore [Ri], reductive monoids are exactly the affine embeddings of reductive groups.

(II) *Projective case:* Let M be a reductive monoid with zero and unit group G . Then there exists a central one-parameter subgroup $\epsilon : \mathbb{G}_m^* \rightarrow T$, with

image Z , such that $\lim_{t \rightarrow 0} \epsilon(t) = 0$. Moreover, the quotient space

$$\mathbb{P}_\epsilon(M) := (M \setminus \{0\})/Z$$

is a normal projective variety on which $G \times G$ acts via $(g, h) \cdot [x] = [gxh^{-1}]$. Hence, $\mathbb{P}_\epsilon(M)$ is a normal projective embedding of the quotient group G/Z . These varieties were introduced by Renner in his study of algebraic monoids ([R2], [R3]). Notably, normal projective embeddings of connected reductive groups are exactly the projectivizations of normal algebraic monoids [Ti].

5.2. Algebraic monoids and algebraic rational cells.

Lemma 5.1. *Let $\varphi : L \rightarrow M$ be a finite dominant morphism of normal, reductive monoids. Then φ is a quotient map; specifically,*

$$L/\mu \simeq M,$$

where $\mu = \ker(\varphi|_{G_L})$ and G_L is the unit group of L .

Proof. Because μ is a finite and normal subgroup of the connected reductive group G_L , it is central. Hence $\mu \subset T_L$ (for the center of G_L is the intersection of all its maximal tori). It follows that the induced map $\tilde{\varphi} : L/\mu \rightarrow M$ is bijective and birational. But M is normal, so $\tilde{\varphi}$ is an isomorphism. \square

The following is crucial for our purposes.

Lemma 5.2 ([Gr, Lemma 1]). *Let G be a connected linear algebraic group. Let X be a G -variety. Then the action of G on $A_*(X)$ is trivial.* \square

Corollary 5.3. *Let $\varphi : L \rightarrow M$ be a finite dominant morphism of normal algebraic monoids. Then φ induces an isomorphism of (rational) Chow groups, namely, $A_*(L) \simeq A_*(M)$.*

Proof. By Lemma 5.1 and [F], Example 1.7.6, we have $(A_*(L))^\mu \simeq A_*(M)$. Now, since the action of μ on $A_*(L)$ comes induced from the action of G on $A_*(L)$ we have $(A_*(L))^{G_L} \subset (A_*(L))^\mu$. But, by Lemma 5.2, we have $(A_*(L))^{G_L} = A_*(L)$. We conclude that $(A_*(L))^\mu = A_*(L)$. \square

Let M and N be reductive monoids. Following Renner [R2], we write $M \sim_0 N$ if there is a reductive monoid L and finite dominant morphisms $L \rightarrow M$ and $L \rightarrow N$ of algebraic monoids. One checks that this gives rise to an equivalence relation. The following basic result, a consequence of Corollary 5.3, states that rational Chow groups are an invariant of the equivalence classes.

Corollary 5.4. *Let M and N be reductive monoids. If $M \sim_0 N$, then $A_*(M) \simeq A_*(N)$.* \square

Now let M be a reductive monoid with zero and unit group G . Recall that $T \times T$ acts on M via $(s, t) \cdot x = txs^{-1}$ and 0 is the unique attractive fixed point for this action. Recall that the number of closed irreducible $T \times T$ -invariant curves in M is finite (all of them passing through 0), and it equals $|\mathcal{R}_1|$. So

each closed $T \times T$ -curve of M can be written as $\overline{T x T}$, where $x \in \mathcal{R}_1$. (Indeed, they correspond to the $T \times T$ -fixed points of $\mathbb{P}_\epsilon(M)$ [G5, Theorem 3.1].) It follows that $\dim M \leq |\mathcal{R}_1|$. Similarly, \overline{T} is an affine T -variety with 0 as its unique attractive fixed point and with finitely many T -stable curves. The number of these curves equals $|E_1|$ and so $\dim T \leq |E_1|$. Next we provide combinatorial criteria for showing when M is an algebraic rational cell (for the $T \times T$ -action). This adds to the list of equivalences from [R2] and [R3].

Theorem 5.5. *Let M be a reductive monoid with zero and unit group G . Then the following are equivalent.*

- (a) $M \sim_0 \prod_i M_{n_i}(\mathbb{k})$.
- (b) If T is a maximal torus of G , then $\dim T = |E_1(\overline{T})|$.
- (c) $\overline{T} \sim_0 \mathbb{A}^n$.
- (d) $(\overline{T}, 0)$ is an algebraic rational cell.
- (e) $(M, 0)$ is an algebraic rational cell.
- (f) $\dim M = |\mathcal{R}_1|$.

Proof. The equivalence of (a), (b) and (c) is proven in [R2, Theorem 2.1] (no use of rational smoothness is made there). The implication (c) \Rightarrow (d) follows from Corollary 5.4 and Proposition 3.3. On the other hand, condition (d) implies (b) because of Corollary 3.8 and the fact that $|E_1(\overline{T})|$ is the number of T -invariant curves of \overline{T} passing through 0. Hence conditions (a), (b), (c) and (d) are all equivalent.

Certainly (a) implies (e), by Corollary 5.4 and Proposition 3.3. In turn, (e) yields (f) due to Corollary 3.8 and the fact that the number of closed irreducible $T \times T$ -curves in M equals $|\mathcal{R}_1|$. So to conclude the proof it suffices to show that (f) implies (b). For this we argue as follows.

Assume (f) and recall that each closed $T \times T$ -curve in M can be written as $\overline{T x T}$, with $x \in \mathcal{R}_1$. Moreover, if we write $x = ew$, with $e \in E_1$ and $w \in W$, then $T \times T$ acts on $\overline{T x T}$ through the character $(\lambda_e, \lambda_e(\text{int}(w)))$, where $\lambda_e : T \rightarrow \epsilon T \simeq \mathbb{k}^*$ is the character sending t to et .

Now, for each $x = ew \in \mathcal{R}_1$, we can find a finite equivariant surjective map $\pi_x : \overline{T x T} \rightarrow \mathbb{k}_x$. Here, $T \times T$ -acts on $\mathbb{k}_x \simeq \mathbb{k}$ via $(\lambda_e, \lambda_e(\text{int}(w)))$. Since $\overline{T x T}$ is $T \times T$ -invariant and closed in M , we can extend π_x to an equivariant morphism $\pi_x : M \rightarrow \mathbb{k}_x$. Sincronizing efforts via the product map, we obtain a $T \times T$ -equivariant map

$$\pi : M \rightarrow V = \prod_{x \in \mathcal{R}_1} \mathbb{k}_x, \quad m \mapsto (\pi_x(m))_{x \in \mathcal{R}_1}.$$

By construction, π is finite, and given that $\dim M = |\mathcal{R}_1|$, it is also surjective.

Let $\Delta T \subset T \times T$ be the diagonal torus. We know that the fixed point set $M^{\Delta T}$ equals \overline{T} . Let us look at the restriction map

$$\pi : \overline{T} \rightarrow V^{\Delta T}.$$

We claim that $\dim V^{\Delta T} = |E_1(\bar{T})|$. Indeed, it is clear that for $e \in E_1(\bar{T})$, the invariant affine line $\mathbb{k}_e \subset V$ is fixed by ΔT , since $\lambda_e(t)e\lambda_e(t^{-1}) = e$. Hence

$$\prod_{e \in E_1} \mathbb{k}_e \subset V^{\Delta T}.$$

Thus,

$$|E_1(\bar{T})| = \dim \prod_{e \in E_1(\bar{T})} \mathbb{k}_e \leq \dim V^{\Delta(T)} = \dim T.$$

But, in general, $\dim T \leq |E_1(\bar{T})|$. Hence $\dim T = |E_1(\bar{T})|$. As this is condition (b), the proof is now complete. \square

Remark 5.6. Let M be a reductive monoid with zero. Theorem 5.5 gives a converse to Corollary 3.8: $(M, 0)$ is an algebraic rational cell if and only if $\dim M = |\mathcal{R}_1|$.

Theorem 5.5 and [R3, Theorem 2.4] immediately give the following. Notice that the cycle map is not needed in the proof.

Corollary 5.7. *Let $\mathbb{k} = \mathbb{C}$. Let M be a reductive monoid with zero, and let \bar{T} be the associated affine toric variety. Then M (resp. \bar{T}) is rationally smooth if and only if M (resp. \bar{T}) is an algebraic rational cell. \square*

5.3. \mathbb{Q} -filtrable projective group embeddings. We start by recalling [R2, Definition 2.2].

Definition 5.8. A reductive monoid M with zero element is called *quasi-smooth* if, for any minimal non-zero idempotent $e \in E(M)$, M_e satisfies the conditions of Theorem 5.5.

In other words, M is quasismooth if and only if M_e is an algebraic rational cell, for any minimal non-zero idempotent $e \in E(M)$.

Now consider the projective group embedding $\mathbb{P}_e(M) = (M \setminus \{0\})/Z$ (as in Section 5.1). It is worth noting that M is quasismooth if and only if $\mathbb{P}_e(M)$ is rationally smooth. Indeed, when $\mathbb{k} = \mathbb{C}$, this is the content of [R3, Theorem 2.5]. In the general case, one has to first adapt the results of [Br3], replacing rational cohomology by l -adic cohomology, and then argue as in [R3, Theorems 2.4 and 2.5].

Next is the second main result of this section. It is an extension of [G1, Theorem 7.4] to equivariant Chow groups.

Theorem 5.9. *Let M be a reductive monoid with zero. If M is quasismooth, then the projective group embedding $\mathbb{P}_e(M)$ is \mathbb{Q} -filtrable (as in Section 4).*

Proof. The strategy is to adapt the proof of [G1, Theorem 7.4] in light of Proposition 3.3 and Theorem 5.5. Recall that, by [R2, Theorem 3.4], $\mathbb{P}_e(M)$ comes equipped with a BB-decomposition

$$\mathbb{P}_e(M) = \bigsqcup_{r \in \mathcal{R}_1} C_r,$$

where \mathcal{R}_1 identifies to $\mathbb{P}_\epsilon(M)^{T \times T}$. (In fact these cells are $B \times B$ -invariant, where B is a Borel subgroup of G .) Given that $\mathbb{P}_\epsilon(M)$ is normal, projective, and \mathcal{R}_1 is finite, this BB-decomposition is filtrable (Theorem 2.2). So we just need to show that these cells are algebraic rational cells. Furthermore, since the C_r are affine $T \times T$ -varieties with an attractive fixed point $[r]$, Proposition 3.3 reduces the proof to showing that $A_*(C_r) \simeq \mathbb{Q}$.

Bearing this in mind, we delve a bit further into the structure of these cells. By [R2, Lemma 4.6 and Theorem 4.7], each C_r equals

$$U_1 \times C_r^* \times U_2,$$

where the U_i are affine spaces. Moreover, writing $r \in \mathcal{R}_1$ as $r = ew$, with $e \in E_1(\overline{T})$ and $w \in W$, yields $C_r^* = C_e^*w$. Hence, by the Kunnet formula (which holds, because the U_i 's are affine spaces), we are further reduced to showing that $A_*(C_e^*) \simeq \mathbb{Q}$, for $e \in E_1(\overline{T})$.

Now we call the reader's attention to [R2, Theorem 5.1]. It states that if M is quasismooth, then

$$C_e^* = f_e M(e)/Z,$$

for some unique $f_e \in E(\overline{T})$, where $M(e) = M_e Z$, and M_e is reductive monoid with e as its zero. By hypothesis, we know that M_e is an algebraic rational cell, that is, $A_*(M_e) \simeq \mathbb{Q}$. Since $M(e)/Z$ is a reductive monoid with $[e]$ as its zero, and $M_e \sim_0 M(e)/Z$, Corollary 5.4 yields $A_*(M(e)/Z) \simeq \mathbb{Q}$. Now, by [Br4, Lemma 1.1.1], one can find a one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow T$, with image S , such that $\lambda(0) = f$ and

$$f_e M(e)/Z = (M(e)/Z)^S.$$

That is, $f_e M(e)/Z$ is the S -fixed point set of $M(e)/Z$. But now we invoke Lemma 2.5 to get

$$\dim_{\mathbb{Q}} A_*((M(e)/Z)^S) \leq \dim_{\mathbb{Q}} A_*(M(e)/Z) = 1.$$

Hence, *a fortiori*

$$\dim_{\mathbb{Q}} A_*((M(e)/Z)^S) = \dim_{\mathbb{Q}} A_*(f_e M(e)/Z) = \dim_{\mathbb{Q}} A_*(C_e^*) = 1.$$

This shows that $A_*(C_e^*) = \mathbb{Q}$, concluding the argument. \square

It is well-known that for projective simplicial toric varieties (equivalently, rationally smooth projective toric varieties) the equivariant cycle map is an isomorphism over \mathbb{Q} . Below we extend this result to all rationally smooth projective group embeddings.

Corollary 5.10. *Let $\mathbb{k} = \mathbb{C}$. If M is a quasismooth monoid with zero, then the equivariant cycle map*

$$cl_{\mathbb{P}_\epsilon(M)}^T : A_*^T(\mathbb{P}_\epsilon(M)) \rightarrow H_*^T(\mathbb{P}_\epsilon(M))$$

is an isomorphism of free S -modules. Moreover, the usual cycle map

$$cl_{\mathbb{P}_\epsilon(M)} : A_*(\mathbb{P}_\epsilon(M)) \rightarrow H_*(\mathbb{P}_\epsilon(M))$$

is an isomorphism of \mathbb{Q} -vector spaces.

Proof. By [G1, Theorem 7.4] $\mathbb{P}_\epsilon(M)$ has no homology in odd degrees, and each cell is rationally smooth, so $H_{*,c}(C_r) \simeq \mathbb{Q}$ and $H_{*,c}^T(C_r) \simeq S$. Now Theorem 5.9 implies that the cycle maps $cl_{C_r} : A_*(C_r) \rightarrow H_{*,c}(C_r)$ and $cl_{C_r}^T : A_*^T(C_r) \rightarrow H_{*,c}^T(C_r)$ are isomorphisms. Arguing by induction on the length of the filtration concludes the proof. \square

In [R2] and [R3], Renner has computed the H -polynomial of a quasi-smooth monoid. This polynomial counts the number of algebraic rational cells (of each dimension) that appear in the BB-decomposition of $\mathbb{P}_\epsilon(M)$. In particular, when $\mathbb{P}_\epsilon(M)$ is simple (i.e. it contains a unique $G \times G$ -orbit), [R2] shows that the number of cells of dimension k equals the number of cells of dimension $n - k$, where $n = \dim \mathbb{P}_\epsilon(M)$. By Corollary 4.4, this yields

$$\dim_{\mathbb{Q}} A_k(\mathbb{P}_\epsilon(M)) = \dim_{\mathbb{Q}} A_{n-k}(\mathbb{P}_\epsilon(M)).$$

Over the complex numbers, this is equivalent to the fact that $\mathbb{P}_\epsilon(M)$ satisfies Poincaré duality for rational singular cohomology. We shall see next that this phenomenon also amounts to Poincaré duality in operational Chow groups.

6. APPLICATIONS TO EQUIVARIANT POINCARÉ DUALITY

The goal is to show that \mathbb{Q} -filtrable T -linear varieties are analogues of the equivariantly formal spaces of Goresky, Kottwitz, MacPherson [GKM], from the viewpoint of equivariant operational Chow groups. In this section we assume $\text{char}(\mathbb{k}) = 0$.

6.1. Equivariant Chow cohomology. Localization theorem. Let X be a T -scheme. The i -th T -equivariant operational Chow group of X , denoted $\text{op}A_T^i(X)$, is defined as follows: an element $c \in \text{op}A_T^i(X)$ is a collection of homomorphisms $c_f^{(m)} : A_m^T(Y) \rightarrow A_{m-i}^T(Y)$, written $z \mapsto f^*c \cap z$, for every T -map $f : Y \rightarrow X$ and all integers m . (The underlying category is the category of T -schemes.) These homomorphisms must satisfy certain compatibility conditions, see [F, Chapter 17] and [EG] for details. For any X , the ring structure on $\text{op}A_T^*(X) := \bigoplus_i \text{op}A_T^i(X)$ is given by composition of such homomorphisms. The ring $\text{op}A_T^*(X)$ is graded, and $\text{op}A_T^i(X)$ can be non-zero for any $i \geq 0$. The basic properties we need are summarized below.

- (i) Cup products $\text{op}A_T^p(X) \otimes \text{op}A_T^q(X) \rightarrow \text{op}A_T^{p+q}(X)$, $a \otimes b \mapsto a \cup b$, making $\text{op}A_T^*(X)$ into a graded associative commutative ring.
- (ii) Contravariant graded ring maps $f^* : \text{op}A_T^i(X) \rightarrow \text{op}A_T^i(Y)$ for arbitrary equivariant morphisms $f : Y \rightarrow X$.
- (iii) Cap products $\text{op}A_T^i(X) \otimes A_m^T(X) \rightarrow A_{m-i}^T(X)$, $c \otimes z \mapsto c \cap z$, making $A_m^T(X)$ into an $\text{op}A_T^*(X)$ -module and satisfying the projection formula.
- (iv) For any T -scheme X of pure dimension n , there is an *equivariant Poincaré duality map*:

$$\mathcal{P}_T : \text{op}A_T^k(X) \rightarrow A_{n-k}^T(X), \quad z \mapsto z \cap [X].$$

If X is nonsingular, then \mathcal{P}_T is an isomorphism, and the ring structure on $\mathrm{op}A_T^*(X)$ is that determined by intersection products of cycles on the mixed spaces X_T . In particular, by (iii) and (iv), $\mathrm{op}A_T^*(X)$ is a graded S -algebra. We say that X *satisfies equivariant Poincaré duality* if \mathcal{P}_T is an isomorphism. Similar remarks apply to the usual (non-equivariant) Poincaré duality map (denoted \mathcal{P}).

Let X be a T -scheme, and let (V, U) be a pair as in Section 2.2. By [EG, Corollary 2], there is an isomorphism $\mathrm{op}A_T^j(X) \simeq \mathrm{op}A^j(X \times U/T)$, provided $V \setminus U$ has codimension more than j . Thus there is a canonical map $i^* : \mathrm{op}A_T^*(X) \rightarrow \mathrm{op}A^*(X)$ induced by restriction to a fiber of $p_{X,T} : X_T \rightarrow U/T$. But, unlike the case of equivariant Chow groups, this map is not surjective in general, and its kernel is not necessarily generated in degree one, not even for toric varieties [KP]. This becomes an issue when trying to translate results from equivariant to non-equivariant Chow cohomology. Nevertheless, for certain \mathbb{Q} -filtrable varieties the map i^* is surjective. Before presenting them, let us recall a definition from [G3].

Definition 6.1. Let X be a complete T -variety. We say that X satisfies *T -equivariant Kronecker duality* if the following conditions hold:

- (i) $A_*^T(X)$ is a finitely generated S -module.
- (ii) The equivariant Kronecker duality map

$$\mathcal{K}_T : \mathrm{op}A_T^*(X) \longrightarrow \mathrm{Hom}_S(A_*^T(X), A_*^T(pt)), \quad \alpha \mapsto (\beta \mapsto \int_X (\beta \cap \alpha)).$$

is an isomorphism of S -modules.

If, in addition, the map \mathcal{K} is also an isomorphism, then we say that X satisfies the *strong T -equivariant Kronecker duality*.

Example 6.2. By the results of [FMSS], [To] and [G3], strong T -equivariant duality varieties include: nonsingular projective T -cellular schemes, complete G -spherical varieties and more generally projective T -linear schemes equivariantly embedded in projective space.

The next result makes \mathbb{Q} -filtrations relevant to the study of T -linear varieties. The proof is an easy adaptation of [G3, Corollary 3.9].

Proposition 6.3. *Let X be a complete T -scheme. If X satisfies the strong T -equivariant Kronecker duality and $A_*^T(X)$ is S -free, then the map*

$$\mathrm{op}A_T^*(X)/\Delta\mathrm{op}A_T^*(X) \rightarrow \mathrm{op}A^*(X)$$

is an isomorphism, where Δ is the character group of T . □

Now we state the localization theorem for equivariant Chow cohomology. It is applicable to possibly singular complete T -schemes, regardless of whether or not $\mathrm{op}A_T^*(X)$ is a free S -module.

Theorem 6.4 ([G3, Theorem 2.28]). *Let X be a complete T -scheme and let $i_T : X^T \rightarrow X$ be the inclusion of the fixed point subscheme. Then the pull-back map*

$$i_T^* : \mathrm{op}A_T^*(X) \rightarrow \mathrm{op}A_T^*(X^T)$$

is injective, and its image is exactly the intersection of the images of

$$i_{T,H}^* : \mathrm{op}A_T^*(X^H) \rightarrow \mathrm{op}A_T^*(X^T),$$

where H runs over all subtori of codimension one in T . \square

Theorem 6.4 makes equivariant Chow cohomology more computable. For instance, a version of GKM theory also holds [G3, Theorem 2.31], and there is a description of the equivariant operational Chow groups of spherical varieties [G3, Section 4], which generalizes [Br1, Theorem 7.3]. In particular, combining Theorem 5.9, Corollary 5.10, Example 6.2 and [G3, Theorem 2.31] yields the following.

Corollary 6.5. *Let M be a reductive monoid with zero. If M is quasi-smooth, then $\mathrm{op}A_{T \times T}^*(\mathbb{P}_\epsilon(M))$ is a free S -module, and it is isomorphic, as an S -algebra, to the ring of piecewise polynomial functions $PP_{T \times T}(\mathbb{P}_\epsilon(M))$ associated to the GKM graph of $\mathbb{P}_\epsilon(M)$. Furthermore, if $\mathbb{k} = \mathbb{C}$, then $\mathrm{op}A_{T \times T}^*(\mathbb{P}_\epsilon(M)) \simeq H_{T \times T}^*(\mathbb{P}_\epsilon(M))$.* \square

For a description of the GKM graph of $\mathbb{P}_\epsilon(M)$ see [G2].

6.2. Equivariant Poincaré duality and Chow homology Betti numbers. Let X be a projective T -variety of pure dimension n . Suppose that X is \mathbb{Q} -filtrable. Then, by Corollary 4.4, $A_*^T(X)$ is a free S -module and $A_*(X)$ is a free \mathbb{Q} -vector space. Now set $b_k := A_k(X)$, and call it the k -th *Chow homology Betti number* of X . It follows from Corollary 4.4 that b_k equals the number of k -dimensional algebraic rational cells. When X is smooth, these cells are actually affine spaces, and we get $b_k = b_{n-k}$ [B1, Corollary 1]. Moreover, Poincaré duality holds, and all equivariant multiplicities are non-zero (Theorem 2.10). In the singular case, this is not necessarily true, and our motivation for this section is to determine in which cases the identity $b_k = b_{n-k}$ holds. Is this equivalent to Poincaré duality for the Chow cohomology of X ? Could it be studied via equivariant multiplicities? Notice that these multiplicities play a fundamental role in Sections 3 and 4. In equivariant cohomology these questions have been answered in [Br5]. Below we provide some analogues of the results of [Br5] in equivariant Chow cohomology. Our methods rely on Theorem 6.4, Proposition 6.3 and the notion of algebraic rational cells. No comparison via the cycle map is needed.

A first approximation to Poincaré duality via equivariant multiplicities is given next. For the corresponding statement in equivariant cohomology, see [Br5, Theorem 4.1].

Lemma 6.6. *Let X be a complete equidimensional T -scheme with finitely many fixed points. If all equivariant multiplicities are non-zero, then the equivariant Poincaré duality map is injective for all $q \in \mathbb{Z}$.*

Proof. In view of Theorem 6.4, the argument is the same as that of [Br5, Theorem 4.1]. We include it for convenience. Let $\alpha \in \text{op}A_T^*(X)$ and suppose that $\alpha \cap [X] = 0$, then

$$\int_X (\alpha \cup \beta) \cap [X] = 0$$

for all $\beta \in A_T^*(X)$. Thus, in Q_S , we have

$$\sum_{x \in X^T} \alpha_x \beta_x e_T(x, X) = 0.$$

By the localization theorem, this identity holds for all sequences $(\beta_x)_{x \in X^T}$ in Q_S . Since, by assumption, no $e_x[X]$ vanishes, we must have $\alpha_x = 0$ for all $x \in X^T$. Thus $\alpha = 0$ (for $i_T^* : \text{op}A_T^*(X) \rightarrow \text{op}A_T^*(X^T)$ is injective). \square

Remark 6.7. Lemma 6.6 applies to: (i) projective nonsingular T -varieties with isolated fixed points, for then the equivariant multiplicities are all inverses of polynomials (Theorem 2.9); (ii) Schubert varieties and toric varieties, as they have only attractive fixed points, so Theorem 2.12 implies that the corresponding equivariant multiplicities are non-zero; (iii) simple projective embeddings of a connected reductive group G , as they have only one closed $G \times G$ -orbit, and $W \times W$ -acts transitively on the $T \times T$ -fixed points (least one of these is attractive, hence so are all).

We now combine our previous results to produce a criterion for Poincaré duality. In equivariant cohomology, this is done in [Br5, Theorem 4.1].

Theorem 6.8. *Let X be a complete equidimensional T -variety with isolated fixed points. Suppose that*

- (a) X is \mathbb{Q} -filtrable, and
- (b) X satisfies the strong T -equivariant Kronecker duality.

Then the following conditions are equivalent.

- (i) X satisfies Poincaré duality.
- (ii) X satisfies T -equivariant Poincaré duality.
- (iii) The Chow homology Betti numbers of X satisfy $b_q(X) = b_{n-q}(X)$ for $0 \leq q \leq n$, and all equivariant multiplicities are nonzero.

If any of these conditions holds, then all equivariant multiplicities are in fact inverses of polynomial functions.

Proof. Assumptions (a) and (b), together with the graded Nakayama lemma, imply readily the equivalence of (i) and (ii).

(ii) \Rightarrow (iii) It only remains to show that all the equivariant multiplicities are nonzero. For this, let $\{[\overline{W}_1], \dots, [\overline{W}_m]\}$ be the basis of $A_*^T(X)$ consisting of the closures of the algebraic rational cells. Fix $j \in \{1, \dots, m\}$, and let x_j be the unique attractive fixed point of W_j . By (ii) there is a unique $\alpha \in \text{op}A_T^*(X)$ such that

$$\alpha \cap [X] - [\overline{W}_j] = 0.$$

But then, arguing as in the proof of Proposition 6.6, the identity

$$\sum_{x_i \in X^T} \beta_{x_i} (\alpha_{x_i} e_{x_i}[X] - e_{x_i}[\overline{W}_j]) = 0$$

holds for all sequences $(\beta_{x_i})_{x_i \in X^T}$ in Q_S . In particular, we have

$$\alpha_{x_j} e_{x_j}[X] - e_{x_j}[\overline{W}_j] = 0.$$

Now, since x_j is an attractive fixed point of \overline{W}_j , it follows that $e_{x_j}[\overline{W}_j] \neq 0$. This yields $\alpha_{x_j} \neq 0$ and $e_{x_j}[X] \neq 0$, so that $e_{x_j}[X]$ is the inverse of a polynomial. Indeed, $e_{x_j}[\overline{W}_j] = \frac{d}{\prod_{s=1}^{\ell} \chi_s}$ (Corollary 3.8). and $\alpha_{x_j} \in S$.

(iii) \Rightarrow (i) In view of Proposition 6.6, it remains to show that

$$\mathcal{P}_T : \text{op}A_T^q(X) \rightarrow A_{n-q}^T(X)$$

is surjective for all $q \in \mathbb{Z}$. For this, it suffices to show that the dimension of $\text{op}A_T^q(X)$ equals that of $A_{n-q}^T(X)$. But this follows from the assumption on the Chow homology Betti numbers combined with the isomorphisms

$$\text{op}A_T^*(X) \simeq \text{op}A^*(X) \otimes_{\mathbb{Q}} S \quad \text{and} \quad A_*^T(X) \simeq A_*(X) \otimes_{\mathbb{Q}} S,$$

where the first one is granted by Proposition 6.3. \square

Remark 6.9. It is worth noting that Kronecker duality does not imply Poincaré duality. For instance, consider the following example from [FMSS, p. 184]. Let X be the closure of a generic torus orbit in the Grassmannian $G(2, 4)$. Then X is a toric variety with Chow homology groups $\mathbb{Z}, \mathbb{Z}, \mathbb{Z}^5$, and \mathbb{Z} in dimensions 0, 1, 2, and 3. By Kronecker duality, the Chow cohomology groups are $\mathbb{Z}, \mathbb{Z}, \mathbb{Z}^5$ and \mathbb{Z} in codimensions 0, 1, 2, and 3. Clearly, the Poincaré duality maps $A^k \rightarrow A_{3-k}$ are not isomorphisms.

It stems from Theorem 6.8 and its proof that the class of \mathbb{Q} -filtrable varieties satisfying the strong T -equivariant Kronecker duality do resemble the equivariantly formal spaces of [GKM]. To push the analogy even further, here is a result to the account that a version of the Morse inequalities holds for these varieties. For the analogous result in equivariant cohomology, see [Br5, Theorem 4.2]. To simplify notation, a T -linear variety is called *special* if it satisfies the strong T -equivariant Kronecker duality (see Example 5.2).

Theorem 6.10. *Let X be a projective T -linear variety of pure dimension n with isolated fixed points. If X is special, \mathbb{Q} -filtrable, and all equivariant multiplicities are nonzero, then the following inequalities hold for the Chow homology Betti numbers:*

$$b_q + b_{q-1} + \dots + b_0 \leq b_{n-q} + b_{n-q+1} + \dots + b_n$$

for $0 \leq q \leq n$;

$$2b_1 + 4b_2 + \dots + 2nb_n \geq n\chi(X),$$

where $\chi(X) = b_0 + b_1 + \dots + b_n$ is the Euler characteristic, i.e. the number of algebraic rational cells of X . In fact, $\chi(X) = |X^T|$. Finally, X satisfies Poincaré duality if and only if

$$2b_1 + 4b_2 + \dots + 2nb_n = n\chi(X).$$

Proof. The proof is an easy adaptation of [Br5, Theorem 4.2], with a few changes. First, notice that X is a special T -linear variety. Hence, it is \mathbb{G}_m -linear, where \mathbb{G}_m acts on X via the generic one-parameter subgroup $\lambda : \mathbb{G}_m \rightarrow T$ chosen to obtain the \mathbb{Q} -filtration. In other words, X also satisfies the strong \mathbb{G}_m -equivariant Kronecker duality. Now, by Proposition 6.3, we have $\text{op}A_{\mathbb{G}_m}^*(X) \simeq \text{op}A^*(X) \otimes_{\mathbb{Q}} \mathbb{Q}[t]$ and $A_{\mathbb{G}_m}^*(X) \simeq A_*(X) \otimes_{\mathbb{Q}} \mathbb{Q}[t]$, as graded vector spaces, where t is an indeterminate of degree 1. On the other hand, since the $e_x[X]$ are nonzero, the same holds for $e'_x[X]$, the \mathbb{G}_m -equivariant multiplicity at x , by [Br1, Lemma 4.5]. It follows that the map

$$\mathcal{P}_{\mathbb{G}_m} : \text{op}A_{\mathbb{G}_m}^q(X) \rightarrow A_{n-q}^{\mathbb{G}_m}(X), \quad z \mapsto z \cap [X],$$

is injective for all $q \in \mathbb{Z}$. In view of these results, Brion's argument from [Br5, Proof of Theorem 4.2] applies verbatim, yielding the result. \square

Example 6.11. Let M be a quasismooth monoid, and consider the associated projective embedding $X = \mathbb{P}_\varepsilon(M)$. Suppose that X has a unique closed $G \times G$ -orbit. By the calculations of [R2] we get $b_k = b_{n-k}$. Since all the $T \times T$ -fixed points in X are attractive, then by Theorem 6.8, X satisfies Poincaré duality for Chow cohomology. Over the complex numbers this reflects the fact that X is rationally smooth. However, notice that the cycle map was not needed in our arguments.

7. CONNECTIONS WITH TOPOLOGY: SPHERICAL VARIETIES

In this section we work over the complex numbers. The aim is to relate the results of this paper with those of [G1] in the case of spherical varieties.

Let G be a connected reductive linear algebraic group with Borel subgroup B and maximal torus $T \subset B$. Recall that given a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$, we can define

$$G(\lambda) = \{g \in G \mid \lambda(t)g\lambda(t)^{-1} \text{ has a limit as } t \rightarrow 0\}.$$

It is well-known that $G(\lambda)$ is a parabolic subgroup of G with unipotent radical $R_u G(\lambda) = \{g \in G \mid \lim_{t \rightarrow 0} \lambda(t)g\lambda(t)^{-1} = 1\}$. Moreover, the centralizer $C_G(\lambda)$ of the image of λ is connected, and the product morphism $R_u G(\lambda) \times C_G(\lambda) \rightarrow G(\lambda)$ is an isomorphism of varieties. Also, the parabolic subgroups $G(\lambda)$ and $G(-\lambda)$ are opposite. Finally, $G(\lambda) = B$ if and only if λ lies in the interior of the Weyl chamber associated with B . See e.g. [Sp, Theorem 13.4.2].

The following result is a particular case of [To, Theorem 3].

Theorem 7.1. *Let Γ be connected solvable linear algebraic group. For any Γ -variety Y with a finite number of orbits, the natural map*

$$A_i(Y) \otimes \mathbb{Q} \longrightarrow W_{-2i}H_{2i}^{BM}(Y, \mathbb{Q}),$$

from the Chow groups into the smallest subspace of Borel-Moore homology with respect to the weight filtration is an isomorphism. \square

Next we show that algebraic rational cells are naturally found on rationally smooth spherical varieties.

Theorem 7.2. *Let X be a G -spherical variety with an attractive T -fixed point x . Let X_x be the unique open affine T -stable neighborhood of x . If X is rationally smooth at x , then (X_x, x) is an algebraic rational cell.*

Proof. Because x is attractive, we may choose λ such that $X_x = X_+(x, \lambda)$ and $G(\lambda) = B$. Since X is rationally smooth at x , so is the open subset X_x . Moreover, X_x is rationally smooth everywhere, and so X_x is a rational cell [G1, Definition 3.4]. By Theorem 7.1 we have

$$A_i(X_x) \simeq W_{-2i}H_{2i}^{BM}(X_x, \mathbb{Q}) \simeq H_c^{2i}(X_x, \mathbb{Q}) = \begin{cases} \mathbb{Q} & \text{if } i = \dim_{\mathbb{C}} X_x \\ 0 & \text{otherwise,} \end{cases}$$

where the last two identifications follow from the fact that X_x is a cone over a rational cohomology sphere [G1, Corollary 3.11]. \square

Let X be a G -spherical variety. Recall that X^T is finite. For convenience, we use the following nomenclature. We say that

- (a) X has an *algebraic* \mathbb{Q} -filtration, if it satisfies Definition 4.1 for some generic one-parameter subgroup λ of T .
- (b) X has a *topological* \mathbb{Q} -filtration, if there exists a generic one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ for which the associated BB-decomposition of X is filtrable, and consists of *rational cells* [G1].

Theorem 7.3. *Let X be a G -spherical variety. If X has a topological \mathbb{Q} -filtration, then this filtration is also an algebraic \mathbb{Q} -filtration.*

Proof. Let $X^T = \{x_1, \dots, x_m\}$. By assumption, there exists a generic one-parameter subgroup such that $X^\lambda = X^T$, and the cells $X_j := X_+(x_j, \lambda)$ are rational cells. Consider the parabolic subgroup $G(\lambda)$. We claim that the cells X_j are invariant under $G(\lambda)$. Indeed, $G(\lambda) = R_u(\lambda) \times C_G(\lambda)$, and $C_G(\lambda)$, being connected, fixes each $x_j \in X^\lambda$. Now let $x \in X_j$ and write $g \in G(\lambda)$ as $g = uh$, with $u \in R_u(\lambda)$ and $h \in C_G(\lambda)$. Then

$$\lambda(t)g \cdot x = \lambda(t)uh\lambda(t)^{-1}\lambda(t) \cdot x = \lambda(t)u\lambda(t)^{-1}h\lambda(t) \cdot x.$$

Taking limits at 0 gives the claim. Because X is spherical, it contains only finitely many orbits of any Borel subgroup of G . Therefore, a Borel subgroup of $G(\lambda)$ has finitely many orbits in X_j . Applying Theorem 7.1 to each X_j yields

$$A_i(X_j) \simeq W_{-2i}H_{2i}^{BM}(X_j, \mathbb{Q}) \simeq H_c^{2i}(X_j, \mathbb{Q}) = \begin{cases} \mathbb{Q} & \text{if } i = \dim_{\mathbb{C}} X_j \\ 0 & \text{otherwise,} \end{cases}$$

noting that X_j is a cone over a rational cohomology sphere [G1, Corollary 3.11]. Therefore, by Lemma 3.2, the cells X_j are algebraic rational cells. This concludes the proof. \square

Remark 7.4. Let X be a G -spherical variety, and let λ be a generic one-parameter subgroup. Then the argument above shows that the cells $X_+(x_j, \lambda)$ are T' -linear varieties, where $T' \subset G(\lambda)$ is a maximal torus of G .

Arguing by induction on the length of the filtration, using the fact that a T -variety with a topological \mathbb{Q} -filtration has no (co)homology in odd degrees, gives immediately the following.

Corollary 7.5. *Let X be a spherical G -variety with a topological \mathbb{Q} -filtration, say $\emptyset = \Sigma_0 \subset \Sigma_1 \subset \dots \subset \Sigma_m = X$. Then, for every j , both cycle maps, $cl_{\Sigma_j} : A_*(\Sigma_j) \rightarrow H_*(\Sigma_j)$ and $cl_{\Sigma_j}^T : A_*^T(\Sigma_j) \rightarrow H_*^T(\Sigma_j)$ are isomorphisms. Moreover, $opA_T^*(\Sigma_j) \simeq H_T^*(\Sigma_j)$, as free S -modules.* \square

We should remark that Theorem 7.3 provides another proof of Theorem 5.9. However, in the case of group embeddings, the approach taken in Section 5 is more intrinsic, for it uses the rich structure of the Chow groups and the fine combinatorial structure of algebraic monoids. Notice that the results of Section 5 are independent of Theorem 7.1. This shows how the notion of algebraic rational cells is well adapted to embedding theory, and opens the way for further work in this direction. For instance, group embeddings have resolution of singularities in arbitrary characteristic, so many results from Section 6 hold for projective group embeddings in arbitrary characteristic (as they do in the particular case of toric varieties). This will be pursued in [G5].

Finally, observe that, when looking for concrete examples, topological \mathbb{Q} -filtrations are slightly more approachable, for they are built using the classical topology of a complex variety, and could be obtained e.g. via Hamiltonian actions and the methods of [Br3]. Our Theorem 7.3 guarantees that the topological knowledge thus acquired gets transformed into algebraic information about the Chow groups. This provides examples of singular spherical varieties for which the cycle map is an isomorphism (e.g. rationally smooth group embeddings). It is worth noting, however, that the study of algebraically \mathbb{Q} -filtrable varieties can be carried out intrinsically, via equivariant intersection theory.

REFERENCES

- [B1] Bialynicki-Birula, A. *Some theorems on actions of algebraic groups*. The Annals of Mathematics, 2nd Ser., Vol 98, No. 3, Nov. 1973, pp. 480-497.
- [B2] Bialynicki-Birula, A. *Some properties of the decompositions of algebraic varieties determined by actions of a torus*. Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 24 (1976), no. 9, 667-674.
- [Bo] Borel, A. *Linear Algebraic Groups*. Third Edition, Springer-Verlag.
- [Br1] Brion, M. *Equivariant Chow groups for torus actions*. Transf. Groups, vol. 2, No. 3, 1997, pp. 225-267.

- [Br2] Brion, M. *Equivariant cohomology and equivariant intersection theory*. Notes by Alvaro Rittatore. NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Representation theories and algebraic geometry (Montreal, PQ, 1997), 1–37, Kluwer Acad. Publ., Dordrecht, 1998.
- [Br3] Brion, M. *Rational smoothness and fixed points of torus actions*. Transf. Groups, Vol. 4, No. 2-3, 1999, pp. 127-156.
- [Br4] Brion, M. *Local structure of algebraic monoids*. Mosc. Math. J. 8 (2008), no. 4, 647-666, 846.
- [Br5] Brion, M. *Poincaré duality and equivariant cohomology*. Michigan Math. J. 48, 2000, pp. 77-92.
- [D] Danilov, V. *The geometry of toric varieties*. Uspekhi Mat. Nauk 33 (1978), no. 2(200), 85-134, 247.
- [EG] Edidin, D., Graham, W. *Equivariant Intersection Theory*. Invent. math. 131, 595-634 (1998).
- [E] Eisenbud, D. *Commutative Algebra with a View Toward Algebraic Geometry*. Springer-Verlag.
- [F] Fulton, W. *Intersection Theory*. Springer-Verlag, Berlin, 1984.
- [FM] Fulton, W., MacPherson, R. *Categorical framework for the study of singular spaces*. Mem. Amer. Math. Soc. 31 (1981), no. 243.
- [FMSS] Fulton, W., MacPherson, R., Sottile, F., Sturmfels, B. *Intersection theory on spherical varieties*. J. of Algebraic Geometry, 4 (1994), 181-193.
- [G1] Gonzales, R. *Rational smoothness, cellular decompositions and GKM theory*. Geometry and Topology, Vol. 18, No. 1 (2014), 291-326.
- [G2] Gonzales, R. *Equivariant cohomology of rationally smooth group embeddings*. Submitted.
- [G3] Gonzales, R. *Equivariant operational Chow rings of T-linear schemes*. arXiv:1304.3645v3.
- [G4] Gonzales, R. *Localization in equivariant operational K-theory and the Chang-Skjelbred property*. arXiv:1403.4412.
- [G5] Gonzales, R. *Equivariant intersection theory on group embeddings*. Preprint.
- [GKM] Goresky, M., Kottwitz, R., MacPherson, R. *Equivariant cohomology, Koszul duality, and the localization theorem*. Invent. math. 131, 25-83 (1998)
- [Gr] Grothendieck, A. *Torsion homologique et sections rationnelles*. Seminaire Claude Chevalley, 3 (1958), Expose. No. 5.
- [Jann] Jannsen, U. *Mixed motives and algebraic K-theory*. Lecture Notes in Mathematics 1400, Springer-Verlag, New York, 1990.
- [J] Joshua, R. *Algebraic K-theory and higher Chow groups of linear varieties*. Math. Proc. Cambridge Phil. Soc., 130, pp. 37-60.
- [JK] Joshua, R.; Krishna, A. *Higher K-theory of toric stacks*. <http://arxiv.org/abs/1210.1057>.
- [KP] Katz, E.; Payne, S. *Piecewise polynomials, Minkowski weights, and localization on toric varieties*. Algebra Number Theory 2 (2008), no. 2, 135-155.
- [Ki] Kimura, S. *Fractional intersection and bivariant theory*. Comm. in Alg. 20:1, pp. 285-302.
- [Ko] Konarski, J. *Decompositions of normal algebraic varieties determined by an action of a one-dimensional torus*. Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 26 (1978), no. 4, 295-300.
- [R1] Renner, L. *Linear algebraic monoids*. Encyclopedia of Math. Sciences, vol. 134, Springer, Berlin (2005).
- [R2] Renner, L. *The H-polynomial of a semisimple monoid*. J. Alg. 319 (2008), 360-376.
- [R3] Renner, L. *Rationally smooth algebraic monoids*. Semigroup Forum 78 (2009), 384-395.

- [Ri] Rittatore, A. *Algebraic monoids and group embeddings*. Transformation Groups, Vol. 3, No. 4, 1998, pp. 375-396.
- [Sp] Springer, T. *Linear Algebraic Groups*. Birkhäuser Classics. 2nd Edition.
- [Su] Sumihiro, H. *Equivariant completion*. J. Math. Kyoto University 14 (1974), 1-28.
- [Ti] Timashev, D. *Homogeneous spaces and equivariant embeddings*. Encyc. Math. Sci. 138, Springer.
- [To] Totaro, B. *Chow groups, Chow cohomology and linear varieties*. Preprint.
- [V] Vistoli, A. *Alexander duality in intersection theory*. Compositio Math., tome 70, n.3 (1989), p. 199-225.

INSTITUT DES HAUTES ÉTUDES SCIENTIFIQUES, 35 ROUTE DE CHARTRES,, F-91440
BURES-SUR-YVETTE, FRANCE

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: `rgonzalesv@ihes.fr`