

Galois equivariance of critical values of L -functions for unitary groups

Lucio GUERBEROFF and Jie LIN



Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Avril 2017

IHES/M/17/04

GALOIS EQUIVARIANCE OF CRITICAL VALUES OF L -FUNCTIONS FOR UNITARY GROUPS

LUCIO GUERBEROFF AND JIE LIN

ABSTRACT. The goal of this paper is to provide a refinement of a formula proved by the first author which expresses some critical values of automorphic L -functions on unitary groups as Petersson norms of automorphic forms. Here we provide a Galois equivariant version of the formula. We also give some applications to special values of automorphic representations of $GL_n \times GL_1$. We show that our results are compatible with Deligne's conjecture.

CONTENTS

1. Introduction	2
1.1. Organization of the paper	3
Acknowledgements	3
Notation and conventions	3
2. Automorphic representations	4
2.1. Unitary groups, Shimura varieties and conjugation	4
2.2. Base change and conjugation	5
2.3. Conjugation of cohomological cuspidal representations	7
2.4. The standard L -function and the motivic normalization	9
2.5. Algebraic Hecke characters	9
3. The doubling method, conjugation and the main theorem	10
3.1. Basic assumptions	10
3.2. The double hermitian space	10
3.3. Petersson norms and CM periods	12
3.4. Eisenstein series and zeta integrals	14
3.5. Modified periods	18
3.6. The main theorem	19
3.7. A refinement	20
4. Applications to general linear groups	21
4.1. Transfer from similitude unitary groups to unitary groups	21
4.2. Base change for unitary groups	22
4.3. Special values of representations of general linear group	23
5. Motivic interpretation	25
5.1. The Deligne conjecture	25
5.2. Deligne conjecture for tensor product of motives	26
5.3. Compatibility of the main results with the Deligne conjecture	27
References	28

2010 *Mathematics Subject Classification*. 11F67 (Primary) 11F70, 11G18, 11R39, 22E55 (Secondary).

1. INTRODUCTION

In the present paper we provide a Galois equivariant version of a formula for the critical values of L -functions of cohomological automorphic representations of unitary groups. Such formula expresses the critical values in terms of Petersson norms of holomorphic automorphic forms, and was proved by Harris ([Har97]) when the base field is \mathbb{Q} , and by the first author when the base field is a general totally real field ([Gue16]). To state the main theorem, we need to introduce some notation. Let F/F^+ be a CM extension, and let G be a similitude unitary group attached to an n -dimensional hermitian vector space over F . Fix a CM type Φ for F/F^+ , and let (r_τ, s_τ) be the signature of G . Let π be a cohomological, cuspidal automorphic representation of $G(\mathbb{A})$. The weight of π can be parametrized by a tuple of integers $((a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in \Phi}; a_0)$. We let ψ be an algebraic Hecke character of F , with infinity type $(m_\tau)_{\tau: F \hookrightarrow \mathbb{C}}$. Under some additional hypotheses on π , it is proved in Theorem 4.5.1 of [Gue16] that

$$(1) \quad L^S \left(m - \frac{n-1}{2}, \pi \otimes \psi, \text{St} \right) \sim (2\pi i)^{[F^+:\mathbb{Q}](mn-n(n-1)/2)-2a_0} D_{F^+}^{\lfloor \frac{n+1}{2} \rfloor / 2} P(\psi) Q^{\text{hol}}(\pi)$$

for certain integers $m > n$ satisfying an inequality determined by the signatures of G and the weight of π . In this expression, \sim means that the elements on each side, which belong to $E(\pi, \psi) \otimes \mathbb{C}$, differ by an element of $E(\pi, \psi) \otimes F^{\text{Gal}}$. Here $E(\pi, \psi) = E(\pi) \otimes E(\psi)$, where $E(\pi)$ and $E(\psi)$ are certain number fields explicitly attached to π and ψ , and F^{Gal} is the Galois closure of F in \mathbb{C} . The element $P(\psi)$ is an explicit expression involving CM periods attached to ψ , and $Q^{\text{hol}}(\pi)$ is an automorphic quadratic period, which is basically given as the Petersson norm of an arithmetic holomorphic vector in π . It turns out that, up to multiplication by an element in $E(\pi, \psi) \otimes F^{\text{Gal}}$, the product $(2\pi i)^{-2a_0} P(\psi) Q^{\text{hol}}(\pi)$ can be seen as the inverse of a Petersson norm of an arithmetic vector in $\pi \otimes \psi$ contributing to antiholomorphic cohomology. In this paper, we will consider a Galois equivariant version of formula (1) when using these inverse Petersson norms, which we denote by $Q(\pi, \psi)$ in this introduction, for fixed choices of arithmetic vectors; we refer the reader to Subsection 3.3 for more details. Galois equivariance means that we obtain a formula up to factors in $E(\pi, \psi)$ instead of $E(\pi, \psi) \otimes F^{\text{Gal}}$. We also incorporate an auxiliary algebraic Hecke character α , which will provide useful for applications. The infinity type of α will be assumed to be given by an integer κ at all places of Φ , and by 0 at places outside Φ .

The formula (1) is proved using the doubling method, and it relies on a detailed analysis of certain global and local zeta integrals. In this paper, we study the action of $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ on these zeta integrals. The global and the finite zeta integrals are not hard to analyze, but the archimedean zeta integral is subtler. This integral depends on certain choices that will not be explicated in this introduction, but most importantly, it depends on π , ψ , α and the integer m . We denote it by $Z_\infty(m; \pi, \psi, \alpha)$ here. Garrett proved in [Gar08] that $Z_\infty(m; \pi, \psi, \alpha)$ is non-zero and belongs to F^{Gal} , so it

doesn't appear in (1), but at the moment we must include it in our Galois equivariant formulation.

Besides the archimedean integral, there is another factor that needs to be added to (1) to obtain a Galois equivariant version, which is not originally visible since it belongs to F^{Gal} . To the quadratic extension F/F^+ , there is attached a quadratic character ε_F of $\text{Gal}(\bar{F}/F^+)$ and an Artin motive $[\varepsilon_F]$ over F^+ . We let $\delta[\varepsilon_F]$ be the period of this motive, an element of \mathbb{C}^\times well defined up to multiplication by an element in \mathbb{Q}^\times . It can also be seen as $c^-[\varepsilon_L]$. When $F^+ = \mathbb{Q}$, it can be explicitly written down as a classical Gauss sum. In any case, $\delta[\varepsilon_F] \in F^{\text{Gal}}$.

We then define

$$\Omega(m; \pi, \psi, \alpha) = \frac{Q(\pi, \psi, \alpha)}{Z_\infty(m; \pi, \psi, \alpha)}.$$

We can define $L^*(s, \pi \otimes \psi, \text{St}, \alpha) \in E(\pi, \psi, \alpha) \otimes \mathbb{C}$ to be the collection of standard L -functions of ${}^\sigma\pi \otimes {}^\sigma\psi$, twisted by ${}^\sigma\alpha$, for $\sigma : E(\pi, \psi, \alpha) \hookrightarrow \mathbb{C}$. The automorphic representation ${}^\sigma\pi$ and the Hecke characters ${}^\sigma\psi$ and ${}^\sigma\alpha$ are obtained from π , ψ and α by conjugation by σ (see Subsections 2.3 and 2.5 for details). We can similarly define $\Omega^*(m; \pi, \psi, \alpha) \in E(\pi, \psi, \alpha) \otimes \mathbb{C}$. The main result of this paper is the following.

Theorem 1. *Keep the assumptions as above. Let $m > n - \frac{n}{2}$ be an integer satisfying inequality (3.2.1). Then*

$$L^{*,S} \left(m - \frac{n}{2}, \pi \otimes \psi, \text{St}, \alpha \right) \sim_{E(\pi, \psi, \alpha)} (2\pi i)^{[F^+:\mathbb{Q}](mn-n(n-1)/2)} D_{F^+}^{-\lfloor \frac{n}{2} \rfloor / 2} \delta[\varepsilon_F]^{\lfloor \frac{n}{2} \rfloor} \Omega^*(m; \pi, \psi, \alpha).$$

The presence of m in the element $\Omega^*(m; \pi, \psi, \alpha)$ is, as we explained above, due to the difficulty in analyzing the Galois action on the archimedean zeta integrals. If we all factors in $E(\pi, \psi, \alpha) \otimes F^{\text{Gal}}$, then we can replace $\Omega^*(m; \pi, \psi, \alpha)$ with the period $(2\pi i)^{-2a_0} P(\psi; \alpha) Q^{\text{hol}}(\pi)$, which becomes formula (1) when α is trivial. In any case, we can at least stress that the dependence on m of $\Omega(m; \pi, \psi, \alpha) \in E(\pi, \psi, \alpha) \otimes \mathbb{C}$ disappears if we see it modulo $E(\pi, \psi, \alpha) \otimes F^{\text{Gal}}$.

1.1. Organization of the paper. Section 2. Section 3. Section ???. Section ???.

Acknowledgements. The authors would like to thank Michael Harris for his numerous suggestions and comments. The first author would also like to thank Fabian Januszewski for several useful conversations.

Notation and conventions. We fix an algebraic closure \mathbb{C} of \mathbb{R} , a choice of $i = \sqrt{-1}$, and we let $\bar{\mathbb{Q}}$ denote the algebraic closure of \mathbb{Q} in \mathbb{C} . We let $c \in \text{Gal}(\mathbb{C}/\mathbb{R})$ denote complex conjugation on \mathbb{C} , and we use the same letter to denote its restriction to $\bar{\mathbb{Q}}$. Sometimes we also write $c(z) = \bar{z}$ for $z \in \mathbb{C}$. We let $\Gamma_{\mathbb{Q}} = \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. For a number field K , we let \mathbb{A}_K and $\mathbb{A}_{K,f}$ denote the rings of adèles and finite adèles of K respectively. When $K = \mathbb{Q}$, we write $\mathbb{A} = \mathbb{A}_{\mathbb{Q}}$ and $\mathbb{A}_f = \mathbb{A}_{\mathbb{Q},f}$.

A CM field L is a totally imaginary quadratic extension of a totally real field K . A CM type Φ for L/K is a choice of one of the two possible extensions to L of each embedding of K .

All vector spaces will be finite-dimensional except otherwise stated. By a variety over a field K we will mean a geometrically reduced scheme of finite type over K . We let $\mathbb{S} = R_{\mathbb{C}/\mathbb{R}}\mathbb{G}_{m,\mathbb{C}}$. We denote by c the complex conjugation map on \mathbb{S} , so for any \mathbb{R} -algebra A , this is $c \otimes_{\mathbb{R}} 1_A : (\mathbb{C} \otimes_{\mathbb{R}} A)^\times \rightarrow (\mathbb{C} \otimes_{\mathbb{R}} A)^\times$. We usually also denote it by $z \mapsto \bar{z}$, and on complex points it should not be confused with the other complex conjugation on $\mathbb{S}(\mathbb{C}) = (\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C})^\times$ on the second factor.

A tensor product without a subscript between \mathbb{Q} -vector spaces will always mean tensor product over \mathbb{Q} . For any number field K , we denote by $J_K = \text{Hom}(K, \mathbb{C})$. For $\sigma \in J_K$, we let $\bar{\sigma} = c\sigma$. Let E and K be number fields, and $\sigma \in J_K$. If $\alpha, \beta \in E \otimes \mathbb{C}$, we write $\alpha \sim_{E \otimes K, \sigma} \beta$ if either $\beta = 0$ or if $\beta \in (E \otimes \mathbb{C})^\times$ and $\alpha/\beta \in (E \otimes K)^\times$, viewed as a subset of $(E \otimes \mathbb{C})^\times$ via σ . When $K = \mathbb{Q}$ and $\sigma = 1$, we simply write $a \sim_E b$. There is a natural isomorphism $E \otimes \mathbb{C} \simeq \prod_{\varphi \in J_E} \mathbb{C}$ given by $e \otimes z \mapsto (\varphi(e)z)_\varphi$ for $e \in E$ and $z \in \mathbb{C}$. Under this identification, we denote an element $\alpha \in E \otimes \mathbb{C}$ by $(\alpha_\varphi)_{\varphi \in J_E}$.

We choose Haar measures on local and adelic points of unitary groups as in the Introduction of [Har97].

2. AUTOMORPHIC REPRESENTATIONS

In this section we recall some basic facts about cohomological representation of a unitary group and their conjugation by $\text{Aut}(\mathbb{C})$.

2.1. Unitary groups, Shimura varieties and conjugation. Let V be a hermitian space of dimension n over F with respect to F/F^+ . We let U be the (restriction of scalars from F^+ to \mathbb{Q} of the) unitary group associated to V , and we let G be the associated similitude unitary group with rational similitude factors. To be more precise, U and G are reductive algebraic groups over \mathbb{Q} , such that for any \mathbb{Q} -algebra A , the A -points are given as

$$U(A) = \{g \in \text{Aut}_{F \otimes A}(V \otimes A) : gg^* = \text{Id}\}$$

and

$$G(A) = \{g \in \text{Aut}_{F \otimes A}(V \otimes A) : gg^* = \mu(g)\text{Id} \text{ with } \mu(g) \in A^\times\},$$

where we write g^* for the adjoint of g with respect to the hermitian form.

We fix once and for all a CM type Φ for F/F^+ . Attached to G and Φ is a Shimura variety which we denote by $S = \text{Sh}(G, X)$. The choice of Φ and an orthogonal basis of V determine a choice of CM point $x \in X$, which will be fixed throughout the paper. We let $K_x \subset G_{\mathbb{R}}$ be the centralizer of x . For each compact open subgroup $K \subset G(\mathbb{A}_f)$, S_K will be the corresponding Shimura variety at level K . We also let $E(G, X)$ be the reflex field of S . For each $\tau \in J_F$, we let $(r_\tau, s_\tau) = (r_\tau(V), s_\tau(V))$ be the signature of V at the place τ . We can write the group $G_{\mathbb{R}}$ as

$$(2.1.1) \quad G_{\mathbb{R}} \cong G \left(\prod_{\tau \in \Phi} \text{GU}(r_\tau, s_\tau) \right),$$

which is defined to be the set of tuples $(g_\tau)_{\tau \in \Phi}$ that have the same similitude factor.

We will parametrize irreducible representations of $G_{\mathbb{C}}$ and of $K_{x,\mathbb{C}}$ by their highest weights, and we will use the conventions used in [Gue16], 3.3. Thus, an irreducible representation of $G_{\mathbb{C}}$ (resp. $K_{x,\mathbb{C}}$) will be given by a highest weight $\mu \in \Lambda^+$ (resp. $\lambda \in \Lambda_c^+$). The corresponding representations will be denoted by W_μ (resp. V_λ). All these parameters can be written as tuples

$$((a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in \Phi}; a_0)$$

where each $a_{\tau,i}$ and a_0 are integers, and $a_{\tau,1} \geq \dots \geq a_{\tau,n}$ for each $\tau \in \Phi$ in the case of irreducible representations of $G_{\mathbb{C}}$. In the case of representations of $K_{x,\mathbb{C}}$, the condition is that $a_{\tau,1} \geq \dots \geq a_{\tau,r_\tau}$ and $a_{\tau,r_\tau+1} \geq \dots \geq a_{\tau,n}$ for every $\tau \in \Phi$.

Let $\sigma \in \text{Aut}(\mathbb{C})$. We let $({}^\sigma G, {}^\sigma X)$ be the conjugate Shimura datum with respect to the automorphism σ and the CM point $x \in X$ (see [Mil90], II.4, for details). It follows from [MS10], Theorem 1.3, that the group ${}^\sigma G$ can be realized as the unitary group attached to another n -dimensional hermitian space ${}^\sigma V$, whose signatures at infinity are obtained by permutation from those of G . More precisely,

$$(r_\tau({}^\sigma V), s_\tau({}^\sigma V)) = (r_{\sigma\tau}(V), s_{\sigma\tau}(V))$$

for any $\tau \in J_F$. The local invariants of ${}^\sigma V$ at finite places are the same as those of V , and we identify ${}^\sigma G(\mathbb{A}_f)$ with $G(\mathbb{A}_f)$ without further mention.

We can also conjugate automorphic vector bundles, as in [Mil90]. The CM point $x \in X$ will be fixed throughout, and all conjugations will be with respect to this fixed point. For any $\sigma \in \text{Aut}(\mathbb{C})$, we have a CM point ${}^\sigma x \in {}^\sigma X$, and we let ${}^\sigma \Lambda^+$ and ${}^\sigma \Lambda_c^+$ denote the corresponding set of dominant weights for the groups ${}^\sigma G$ and $K_{{}^\sigma x} \subset {}^\sigma G_{\mathbb{R}}$. When x needs to be specified, we will denote Λ_c^+ by $\Lambda_{c,x}^+$. Suppose that \mathcal{E}_λ is a fully decomposed automorphic vector bundle over $S_{\mathbb{C}}$, associated with the irreducible representation of $K_{x,\mathbb{C}}$ with highest weight λ . Then $\mathcal{E}_\lambda \times_{\mathbb{C},\sigma} \mathbb{C}$ is a vector bundle over $S_{\mathbb{C}} \times_{\mathbb{C},\sigma} \mathbb{C}$, and identifying the latter with ${}^\sigma S_{\mathbb{C}}$, we get an automorphic vector bundle ${}^\sigma \mathcal{E}_\lambda$ over ${}^\sigma S_{\mathbb{C}}$. It is fully decomposed, associated with an irreducible representation of $K_{{}^\sigma x,\mathbb{C}}$ whose highest weight we denote by ${}^\sigma \lambda \in {}^\sigma \Lambda_c^+$.

2.2. Base change and conjugation. Let E/F be an unramified quadratic extension of local non-archimedean fields. Let G be a reductive group over F . Let P be a minimal parabolic subgroup of $G(F)$ and M be a corresponding Levi factor.

Let χ be an unramified character of M . We may regard it as a representation of P . The unitarily parabolic induction is defined as

$$i_P^G(\chi) = \{\phi : G(F) \rightarrow \mathbb{C} \text{ continuous} : \phi(pg) = \delta_P^{1/2}(p)\chi(p)\phi(g), p \in P, g \in G(F)\}$$

where δ_P is the modulus character of P (see [Min11]).

The unitarily parabolic induction gives rise to a surjective map from the set of unramified characters of M to the set of isomorphism classes of unramified representations of $G(F)$. Two unramified characters induce the same $G(F)$ -representation if and only if they are equivalent under the action of the Weyl group.

2.2.1. *Conjugation of representations.* Let $\sigma \in \text{Aut}(\mathbb{C})$. Let V be a complex representation of $G(F)$. We let ${}^\sigma V = V \otimes_{\mathbb{C}, \sigma} \mathbb{C}$, with $G(F)$ acting on the first factor.

Let χ be as before. If $\phi \in i_P^G(\chi)$, we have $(\sigma \circ \phi)(pg) = \sigma(\delta_P^{1/2}(p))^\sigma \chi(g)(\sigma \circ \phi)(g)$ for any $p \in P$ and $g \in G(F)$.

We define the character T_σ on P by

$$T_\sigma(p) = \frac{\sigma(\delta_P^{1/2}(p))}{\delta_P^{1/2}(p)}.$$

It is easy to see that $\sigma(i_P^G(\chi)) \simeq i_P^G(T_\sigma * \chi)$.

2.2.2. *Local base change.* Let E/F be an unramified quadratic extension of local non-archimedean fields. If we look at the dual side, we can construct a base change map which sends unramified representations of $G(F)$ to unramified representations of $G(E)$. Our aim is to study the commutativity of the local base change and the conjugation by an element of $\text{Aut}(\mathbb{C})$ for quasi-split unitary groups.

We write P_E for a minimal parabolic subgroup of $G(E)$, and M_E for a corresponding Levi factor. The base change map induces a map from the set of equivalence classes of characters of M (under the action of the Weyl group) to those of $M(E)$. We write $[\chi]$ for the equivalence class of χ . We take χ_E a character in the equivalence class of the image of $[\chi]$.

We can define the character $T_{E, \sigma}$ of P_E in a similar fashion as T_σ . The commutativity of local base change with conjugation by $\sigma \in \text{Aut}(\mathbb{C})$ can be stated as:

$$(2.2.1) \quad [T_{\sigma, E} * \chi_E] = [(T_\sigma * \chi)_E]$$

2.2.3. *Commutativity for quasi-split unitary groups.* We now prove that (2.2.1) is true for quasi-split unitary groups. Let n be an integer. We assume that $n = 2m$ is even for simplicity.

We take U to be the quasi-split unitary group of rank n with respect to E/F defined over F . Choosing a proper basis, we may identify $U(F)$ with

$$\left\{ X \in GL_n(E) : {}^t \bar{X} \begin{pmatrix} 0 & I_m \\ -I_m & 0 \end{pmatrix} X = \begin{pmatrix} 0 & I_m \\ -I_m & 0 \end{pmatrix} \right\}.$$

We let P be the minimal parabolic given as the intersection of $U(F)$ with the set of upper triangular matrices in $GL_n(E)$.

Let P_0 be the algebraic group defined over F consisting of upper triangular matrices in $GL_m(E)$. Let S be the algebraic group defined over F such that $S(F) = \{X \in M_m(E) : {}^t \bar{X} = X\}$.

The parabolic group P consists of elements of the form

$$\begin{pmatrix} g & gX \\ 0 & {}^t \bar{g}^{-1} \end{pmatrix},$$

where $g \in P_0(F)$ and $X \in S(F)$.

Let $d_l g$ (resp. $d_r g$) be a left (resp. right) invariant Haar measure on $P_0(F)$ and dX be a (left and right) invariant Haar measure on $S(F)$. We may assume that $d_r g = \delta_{P_0}^{-1} d_l(g)$.

Let $\begin{pmatrix} A & AB \\ 0 & {}^t\bar{A}^{-1} \end{pmatrix} \in P$. We have that

$$\begin{pmatrix} A & AB \\ 0 & {}^t\bar{A}^{-1} \end{pmatrix} \begin{pmatrix} g & gX \\ 0 & {}^t\bar{g}^{-1} \end{pmatrix} = \begin{pmatrix} Ag & Ag(X + g^{-1}B{}^t\bar{g}^{-1}) \\ 0 & {}^tAg^{-1} \end{pmatrix}$$

and

$$\begin{pmatrix} g & gX \\ 0 & {}^t\bar{g}^{-1} \end{pmatrix} \begin{pmatrix} A & AB \\ 0 & {}^t\bar{A}^{-1} \end{pmatrix} = \begin{pmatrix} gA & gA(B + A^{-1}X{}^t\bar{A}^{-1}) \\ 0 & {}^tgA^{-1} \end{pmatrix}$$

It is easy to verify that $d_l g dX$ is a left invariant Haar measure and

$$|\det(g)|_F^{2m} d_r g dX$$

is a right invariant Haar measure on P . We obtain that

$$\delta_P \begin{pmatrix} g & X \\ 0 & {}^t\bar{g}^{-1} \end{pmatrix} = \delta_{P_0}(g) |\det(g) \det(\bar{g})|_F^m = \delta_{P_0}(g) |\det(g)|_E^m.$$

The last equation is due to the fact that E/F is unramified. In the following, we write $|\cdot|$ for the absolute value in E .

We write the diagonal of g as (g_1, \dots, g_m) . Then

$$\delta_{P_0}^{1/2}(g) = |g_1|^{\frac{m-1}{2}} |g_2|^{\frac{m-3}{2}} \cdots |g_m|^{-\frac{m-1}{2}}.$$

Therefore, $\delta_P^{1/2}(g) = |g_1|^{\frac{2m-1}{2}} |g_2|^{\frac{2m-3}{2}} \cdots |g_m|^{\frac{1}{2}}$.

We now consider $U(E) \cong GL_n(E)$. We take P_E to be the minimal parabolic subgroup of $U(E)$ consisting of upper triangular matrices. Let $p_E \in U(E)$ with diagonal (g_1, \dots, g_{2m}) . By Theorem 4.1 of [Min11], we have that

$$\chi_E(p_E) = \chi((g_1, \dots, g_m)) \chi((g_{m+1}, \dots, g_{2m}))^{-1}$$

for any character χ . (Here we consider the first case in Theorem 4.1 of *op. cit.*. The proof for the second case is similar.)

We can see easily that $(T_\sigma * \chi)_E = (T_\sigma)_E * \chi_E$. Thus, to show (2.2.1), it is enough to show that $(T_\sigma)_E = T_{\sigma,E}$. In fact, both sides map p_E to

$$\left(\frac{\sigma(|g_1|)}{|g_1|} \right)^{\frac{2m-1}{2}} \left(\frac{\sigma(|g_2|)}{|g_2|} \right)^{\frac{2m-3}{2}} \cdots \left(\frac{\sigma(|g_{2m}|)}{|g_{2m}|} \right)^{-\frac{2m-1}{2}}.$$

2.2.4. Global base change. If we already know that global base change exists, then the commutativity follows from local base change case and strong multiplicity one for GL_n . For example, see Theorem 1.4 of [GR14] for the Jacquet-Langlands transfer.

2.3. Conjugation of cohomological cuspidal representations. From now on, we let $\pi = \pi_\infty \otimes \pi_f$ be an automorphic representation of $G(\mathbb{A})$. We will assume that π satisfies the following list of hypotheses.

Hypothesis 2.3.1. (1) π occurs in the discrete spectrum.

(2) π is cohomological with respect to some irreducible representation $W = W_\mu$ of $G_{\mathbb{C}}$, with $\mu \in \Lambda^+$.

(3) The representation W is defined over \mathbb{Q} .

(4) π_∞ is essentially tempered.

Remark 2.3.1. By Theorem 4.3 of [Wal84], Hypotheses 2.3.1 (1) and (4) imply that π is cuspidal. This can also be deduced by assuming that π_v is tempered at some finite place, as in Proposition 4.10 of [Clo93].

Remark 2.3.2. Hypothesis (3) is assumed mostly for simplicity of notation.

Remark 2.3.3. If μ is regular, in the sense that $a_{\tau,i} > a_{\tau,i+1}$ for every $\tau \in \Phi$ and every $i = 1, \dots, n-1$, then 2.3.1 (1) and (2) imply (4) (see Prop. 4.2 and 5.2 of [LS04] and Prop. 2.2 of [Sch94]).

Under these hypotheses, π_∞ is a discrete series representation that belongs to the L -packet whose infinitesimal character is that of W^\vee .

By Theorem 4.4.1 of [BHR94], the field of definition $\mathbb{Q}(\pi_f)$ of the isomorphism class of π_f is a subfield of a CM field. There is a finite extension $E_0(\pi)$ of $\mathbb{Q}(\pi_f)$, which can also be taken to be a CM field, such that π_f has a model $\pi_{f,0}$ over $E_0(\pi)$ (see Corollary 2.13 of [Har13b]). We let ${}^\sigma\pi_f = \pi_f \otimes_{\mathbb{C},\sigma} \mathbb{C} \cong \pi_{f,0} \otimes_{E_0(\pi),\sigma} \mathbb{C}$. We can also define the conjugate ${}^\sigma\pi_\infty$, a discrete series representation of ${}^\sigma G(\mathbb{R})$, as in (2.19) of [Har13b] (see also [BHR94], 4.2).

We will make the following assumption throughout the paper: there exists an automorphic representation ${}^\sigma\pi$ of ${}^\sigma G(\mathbb{A})$ satisfying Hypotheses 2.3.1 such that $({}^\sigma\pi)_f \cong \pi_f$ (recall that we are identifying ${}^\sigma G(\mathbb{A}_f) \cong G(\mathbb{A}_f)$).

Remark 2.3.4. One of the main results of [BHR94] (Theorem 4.2.3) guarantees the existence of such ${}^\sigma\pi$ when the Harish-Chandra parameter of π_∞ is far enough from the walls. Moreover, under these conditions, we have that $({}^\sigma\pi)_\infty \cong {}^\sigma\pi_\infty$. In [Har13b], 4.3, further conditions under which ${}^\sigma\pi$ is shown to exist are discussed. A particular case of this is when the infinitesimal character of π_∞ is regular and π is not a CAP representation. This last condition is expected to be true for tempered representations. See also Corollary 2.14 of [Har13b].

Remark 2.3.5. Under the above assumptions, ${}^\sigma\pi$ is cohomological of a certain weight $\gamma\mu \in \gamma\Lambda^+$. This parameter can be worked out explicitly as follows. We let π_0 be a constituent of the restriction of π to $U(\mathbb{A}) \subset G(\mathbb{A})$. This is cohomological of weight $((a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in \Phi})$. By a theorem of Labesse ([Lab11]), there exists an automorphic representation $BC(\pi_0)$ (resp. $BC({}^\sigma\pi_0)$) of $GL_n(\mathbb{A}_F)$, which is the base change of π_0 (resp. ${}^\sigma\pi_0$). Moreover, $BC(\pi_0)$ is cohomological of weight

$$\tilde{\mu} = (a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in J_F},$$

where $a_{\tau,i} = -a_{\bar{\tau},n+1-i}$ if $\tau \notin \Phi$. Then the conjugate ${}^\sigma BC(\pi_0)$, as defined for example in [GH16], 2.6, is cohomological of weight

$${}^\sigma\tilde{\mu} = (a_{\sigma\tau,1}, \dots, a_{\sigma\tau,n})_{\tau \in J_F}$$

(see [GH16], Proposition 2.4). On the other hand, in the last subsection we proved that ${}^\sigma BC(\pi_0) \cong BC({}^\sigma\pi_0)$. It follows from the same reasoning as above that ${}^\sigma\pi$ is cohomological of weight

$${}^\sigma\mu = ((a_{\sigma\tau,1}, \dots, a_{\sigma\tau,n})_{\tau \in \Phi}; a_0).$$

2.4. The standard L -function and the motivic normalization. Let π be as above. As in [Har97], 2.7, we can define the standard L -function of π as $L^S(s, \pi, \text{St}) = L^S(s, BC(\pi_0), \text{St})$. Here St refers to the standard representation of the L -group of GL_n over L , π_0 is an irreducible constituent of the restriction of π to $U(\mathbb{A})$, and $BC(\pi_0)$ is the base change of π_0 to an irreducible admissible representation of $\text{GL}_n(\mathbb{A}_L^S)$, for a big enough finite set of places S of L . The base change is defined locally at archimedean places, at split places, and at places of K where the local unitary group U_v and $\pi_{0,v}$ are unramified. Under our assumptions, it is known that $BC(\pi_0)$ is the restriction to $\text{GL}_n(\mathbb{A}_L^S)$ of an automorphic representation Π of $\text{GL}_n(\mathbb{A}_L)$, so we can actually define $L(s, \pi, \text{St})$ at all places as $L(s, \Pi, \text{St})$. We define the motivic normalization by

$$L^{\text{mot}, S}(s, \pi, \text{St}) = L^S\left(s - \frac{n-1}{2}, \pi, \text{St}\right).$$

More generally, if α is an algebraic Hecke character of F , we define

$$L^S(s, \pi, \text{St}, \alpha) = L(s, BC(\pi_0), \text{St}, \alpha),$$

the twisted L -function. The motivic normalization is defined similarly. We define

$$L^{*, \text{mot}, S}(s, \pi, \text{St}, \alpha) = \left(L^{\text{mot}, S}(s, {}^\sigma\pi, \text{St}, {}^\sigma\alpha)\right)_{\sigma \in \text{Aut}(\mathbb{C})}.$$

2.5. Algebraic Hecke characters. Let ψ be an algebraic Hecke character of F , of infinity type $(m_\tau)_{\tau \in J_F}$. Recall that this means that

$$\psi : \mathbb{A}_F^\times / F^\times \rightarrow \mathbb{C}^\times$$

is continuous, and for each embedding $\tau \in J_F$, we have

$$\psi(x) = \tau(x)^{-m_\tau} \bar{\tau}(x)^{-m_{\bar{\tau}}} \quad (x \in F_w^\times).$$

Here w is the infinite place of F determined by τ . We let $\mathbb{Q}(\psi)$ be the field generated over \mathbb{Q} by the values of ψ on $\mathbb{A}_{F,f}^\times$. Then $\mathbb{Q}(\psi)$ is either \mathbb{Q} or a CM field. If $\sigma \in \text{Aut}(\mathbb{C})$, we define ${}^\sigma\psi$ to be the algebraic Hecke character whose values on $\mathbb{A}_{F,f}^\times$ are obtained from those of ψ by applying σ , and whose infinity type is $(m_{\sigma^{-1}\tau})_{\tau \in J_F}$.

We need to fix the following notation. Suppose that α_0 is an algebraic Hecke character of F^+ of finite order, and $\sigma \in J_{F^+}$. Then

$$\delta_\sigma[\alpha_0] \in (\mathbb{Q}(\alpha_0) \otimes \mathbb{C})^\times$$

is the δ -period of the Artin motive $[\alpha_0]$. This is a motive over F^+ with coefficients in $\mathbb{Q}(\alpha_0)$. We also let

$$\delta[\alpha_0] = \delta_1(\text{Res}_{F^+/\mathbb{Q}}[\alpha_0]) \in (\mathbb{Q}(\alpha_0) \otimes \mathbb{C})^\times$$

be the period of the motive $\text{Res}_{F^+/\mathbb{Q}}[\alpha_0]$ obtained from $[\alpha_0]$ by restriction of scalars from F^+ to \mathbb{Q} . It is proved in [Yos94] that

$$\delta[\alpha_0] \sim_{\mathbb{Q}(\alpha_0)} D_{F^+}^{\frac{1}{2}} \prod_{\sigma \in J_{F^+}} \delta_\sigma[\alpha_0].$$

Suppose now that α is an algebraic Hecke character of F of weight w . Then we can write

$$\alpha|_{\mathbb{A}_{F^+}^\times} = \alpha_0 \| \cdot \|_{\mathbb{A}_{F^+}}^{-w},$$

where α_0 is a finite order algebraic Hecke character of F^+ . We define

$$G(\alpha) = \prod_{\sigma \in J_{F^+}} \delta_\sigma[\alpha_0] \in (\mathbb{Q}(\alpha_0) \otimes \mathbb{C})^\times.$$

We can then write

$$(2.5.1) \quad \delta[\alpha_0] \sim_{\mathbb{Q}(\alpha_0)} D_{F^+}^{\frac{1}{2}} G(\alpha).$$

For each embedding $\rho \in J_{\mathbb{Q}(\alpha_0)}$, we let $G(\alpha)_\rho \in \mathbb{C}^\times$ be its ρ -component.

3. THE DOUBLING METHOD, CONJUGATION AND THE MAIN THEOREM

3.1. Basic assumptions. In this section, we briefly recall the doubling method used to obtain the main formula of [Gue16], and explain how it behaves under Galois conjugation. We fix once and for all a cuspidal automorphic representation π of $G(\mathbb{A})$, satisfying all the previous hypotheses. In particular, π is cohomological of type $\mu = ((a_{\tau,1}, \dots, a_{\tau,n})_{\tau \in \Phi}; a_0)$, with $W = W_\mu$ defined over \mathbb{Q} . We also assume that $\pi^\vee \cong \pi \otimes \|\nu\|^{2a_0}$, that π contributes to antiholomorphic cohomology, and that

$$(3.1.1) \quad \dim_{\mathbb{C}} \text{Hom}_{\mathbb{C}[G(\mathbb{A}_f)]} \left(\sigma \pi_f, H_c^d(\sigma S_{\mathbb{C}}, \sigma \mathcal{E}_\mu) \right) \leq 1$$

for all $\sigma \in \text{Aut}(\mathbb{C})$. This is part of Arthur's multiplicity conjectures for unitary groups, a proof of which is expected to appear in the near future. We refer the reader to [KMSW14] and their forthcoming sequels for more details.

We fix a CM type Φ for F/F^+ , and an algebraic Hecke character ψ of F with infinity type $(m_\tau)_{\tau \in J_F}$. We let $\Lambda = \Lambda(\mu; \psi) \in \Lambda_c^+$ be the parameter

$$\Lambda = ((b_{\tau,1}, \dots, b_{\tau,n})_{\tau \in \Phi}; b_0),$$

where

$$b_{\tau,i} = \begin{cases} a_{\tau,s_\tau+i} + m_{\bar{\tau}} - m_\tau - s_\tau & \text{if } 1 \leq i \leq r_\tau, \\ a_{\tau,i-r_\tau} + m_{\bar{\tau}} - m_\tau + r_\tau & \text{if } r_\tau + 1 \leq i \leq n, \end{cases}$$

and $b_0 = a_0 - n \sum_{\tau \in \Phi} m_{\bar{\tau}}$ (this was denoted by $\Lambda(\mu; \eta^{-1})$ in [Gue16]). We similarly define ${}^\sigma \Lambda = \Lambda({}^\sigma \mu; {}^\sigma \psi)$ for $\sigma \in \text{Aut}(\mathbb{C})$.

3.2. The double hermitian space. Given our hermitian space V , we let $-V$ be the hermitian space whose underlying F -vector space is V , but whose hermitian form is multiplied by -1 . Its associated Shimura conjugacy class will be denoted by X^- . We let $2V = V \oplus -V$, and $(G^{(2)}, X^{(2)})$ be the Shimura datum attached to $2V$. The choice of our CM point $x \in X$ gives rise to fixed CM points $x^- = \bar{x} \in X^-$ and $x^{(2)} \in X^{(2)}$. The reflex field of $(G^{(2)}, X^{(2)})$ is \mathbb{Q} , and hence we can identify $({}^\sigma G^{(2)}, {}^\sigma X^{(2)}) = (G^{(2)}, X^{(2)})$ for any $\sigma \in \text{Aut}(\mathbb{C})$. We let $S^{(2)}$ be the associated Shimura variety.

We also let $G^\sharp \subset G \times G$ be the subgroup of pairs with the same similitude factor, and we let $x^\sharp : \mathbb{S} \rightarrow G_{\mathbb{R}}^\sharp$ be the map (x, x^-) . The corresponding Shimura datum will be denoted by (G^\sharp, X^\sharp) , and the Shimura variety by S^\sharp . There is a natural embedding

$$i : (G^\sharp, X^\sharp) \rightarrow (G^{(2)}, X^{(2)})$$

of Shimura data, which induces a closed embedding of Shimura varieties $i : S^\sharp \rightarrow S^{(2)}$.

For $\sigma \in \text{Aut}(\mathbb{C})$, we let $(\sigma G)^\sharp \subset \sigma G \times \sigma G$ be the group defined in a similar fashion but using σV and σG instead of V and G . Using the definition of the twisting, given for example in [Mil90], it is easy to see that we can naturally identify $(\sigma G)^\sharp$ with $\sigma(G^\sharp)$ as a subgroup of $\sigma G \times \sigma G$. We let $\sigma_i : \sigma G^\sharp \hookrightarrow G^{(2)}$ be the inclusion defined above for σV .

Keep in mind that our CM point $x \in X$ is fixed, and this in turn gives choices of CM points $\sigma x \in \sigma X$ for each σ . We fix these CM points, as well as their variant $x^{(2)}$, x^\sharp , σx^\sharp for varying $\sigma \in \text{Aut}(\mathbb{C})$. We will parametrize fully decomposed automorphic vector bundles over the corresponding Shimura varieties by irreducible representations of the corresponding groups $K_{x^{(2)}, \mathbb{C}}$, $K_{x^\sharp, \mathbb{C}}$, $K_{\sigma x, \mathbb{C}}$. We have the following identifications:

$$\begin{aligned} K_{x^\sharp, \mathbb{C}} &\cong \left(\prod_{\tau \in \Phi} GL_{r_\tau, \mathbb{C}} \times GL_{s_\tau, \mathbb{C}} \times GL_{s_\tau, \mathbb{C}} \times GL_{r_\tau, \mathbb{C}} \right) \times \mathbb{G}_m, \mathbb{C}, \\ K_{\sigma x^\sharp, \mathbb{C}} &\cong \left(\prod_{\tau \in \Phi} GL_{r_{\sigma\tau}, \mathbb{C}} \times GL_{s_{\sigma\tau}, \mathbb{C}} \times GL_{s_{\sigma\tau}, \mathbb{C}} \times GL_{r_{\sigma\tau}, \mathbb{C}} \right) \times \mathbb{G}_m, \mathbb{C}, \\ K_{x^{(2)}, \mathbb{C}} &\cong \left(\prod_{\tau \in \Phi} GL_{n, \mathbb{C}} \times GL_{n, \mathbb{C}} \right) \times \mathbb{G}_m, \mathbb{C}. \end{aligned}$$

For

$$\lambda = ((\lambda_{\tau,1}, \dots, \lambda_{\tau,n})_{\tau \in \Phi}; \lambda_0) \in \Lambda_{c,x}^+$$

and

$$\lambda^- = ((\lambda_{\tau,1}^-, \dots, \lambda_{\tau,n}^-)_{\tau \in \Phi}; \lambda_0^-) \in \Lambda_{c,x^-}^+$$

we let

$$(\lambda, \lambda^-)^\sharp = ((\lambda_{\tau,1}, \dots, \lambda_{\tau,n}, \lambda_{\tau,1}^-, \dots, \lambda_{\tau,n}^-)_{\tau \in \Phi}; \lambda_0 + \lambda_0^-) \in \Lambda_{c,x^\sharp}^+.$$

Let

$$\lambda^* = ((-\lambda_{\tau,n}, \dots, -\lambda_{\tau,1})_{\tau \in \Phi}; -\lambda_0) \in \Lambda_{c,x^-}^+.$$

For an integers κ , we define $\lambda^\sharp[\kappa] \in \Lambda_{c,x^\sharp}^+$ as $\lambda^\sharp[\kappa] = (\lambda, \lambda^* \otimes \det^{-\kappa})^\sharp \otimes \nu^\kappa$. Explicitly,

$$\lambda^\sharp[\kappa] = ((\lambda_{\tau,1}, \dots, \lambda_{\tau,n}, -\lambda_{\tau,n} - \kappa, \dots, -\lambda_{\tau,1} - \kappa)_{\tau \in \Phi}; 0).$$

For any pair of integers (m, k) , we let $\mathcal{E}_{m,\kappa}$ be the fully decomposed automorphic line bundle over $S_{\mathbb{C}}^{(2)}$ corresponding to the one-dimensional irreducible representation of $K_{x^{(2)}, \mathbb{C}}$ given by

$$((g_\tau, g'_\tau)_{\tau \in \Phi}; z) \mapsto \prod_{\tau \in \Phi} \det(g_\tau)^{-m-\kappa} \det(g'_\tau)^m.$$

It is easy to see that this line bundle $\mathcal{E}_{m,\kappa}$ has a canonical model over \mathbb{Q} . Its highest weight is parametrized by

$$((-m - \kappa, \dots, -m - \kappa, m, \dots, m)_{\tau \in \Phi}; 0).$$

Recall that Λ was defined in the previous subsection. We then obtain an element $\Lambda^\sharp[\kappa] \in \Lambda_{c,x^\sharp}^+$ as above. The corresponding irreducible representation of $K_{x^\sharp, \mathbb{C}}$ defines an automorphic vector bundle $\mathcal{E}_{\Lambda^\sharp[\kappa]}$ over $S_{\mathbb{C}}^\sharp$. Its conjugate

${}^\sigma \mathcal{E}_{\Lambda^\sharp[\kappa]}$, as an automorphic vector bundle over ${}^\sigma S_{\mathbb{C}}^\sharp$, can be identified with $\mathcal{E}_{\sigma \Lambda^\sharp[\kappa]}$.

Remark 3.2.1. We correct here a simple misprint of [Gue16], Section 4.5, where an element denoted by $\Lambda^\sharp(\ell)$ was used, with $\ell = n \sum_{\tau \in \Phi} m_\tau - m_{\bar{\tau}}$. The correct element to use is $\Lambda^\sharp(0)$ (that is, with $\ell = 0$). Indeed, the only purpose of ℓ was to make sure that the parameter $(\mu + \mu(\eta), \mu^\vee - \mu(\eta))^\sharp$ equals the Serre dual of $\Lambda^\sharp(\ell)$. The computation of these parameters actually shows that the last integer, corresponding to the similitude factor, must be 0 instead of ℓ in both cases, so there is no need to introduce the integer ℓ , which has no influence on the rest of the proof. Also, note that the $\Lambda^\sharp(0)$ of [Gue16] is what we call $\Lambda^\sharp[0]$ here. In this paper we give a slightly more general version of the results for any integer κ .

Let $m \in \mathbb{Z}$ satisfy the inequalities

$$(3.2.1) \quad \frac{n - \kappa}{2} \leq m \leq \min\{-a_{\tau, s_\tau + 1} + s_\tau + m_\tau - m_{\bar{\tau}} - \kappa, a_{\tau, s_\tau} + r_\tau + m_{\bar{\tau}} - m_\tau\}_{\tau \in \Phi}.$$

By Proposition 4.2.1 of [Gue16], there exist non-zero differential operators

$$(3.2.2) \quad \Delta_{m, \kappa} = \Delta_{m, \kappa}(\Lambda) : \mathcal{E}_{m, \kappa}|_{S_{\mathbb{C}}^\sharp} \rightarrow \mathcal{E}_{\Lambda^\sharp[\kappa]},$$

which are moreover rational over the relevant reflex fields (all of these are contained in L^{Gal}). In *op. cit.*, κ was taken to be zero, but the proof for any κ is completely similar.

If $\sigma \in \text{Aut}(\mathbb{C})$, then m also satisfies (3.2.1) for the conjugate Shimura data, and the corresponding differential operator

$${}^\sigma \Delta_{m, \kappa} = \Delta_{m, \kappa}({}^\sigma \Lambda) : \mathcal{E}_{m, \kappa}|_{{}^\sigma S_{\mathbb{C}}^\sharp} \rightarrow \mathcal{E}_{\sigma \Lambda^\sharp[\kappa]}$$

is the conjugate of (3.2.2) under σ .

3.3. Petersson norms and CM periods. We recall now the definition of certain CM periods attached to ψ that appear in our critical value formula. The determinant defines a map $\det : G \rightarrow T^F = \text{Res}_{F/\mathbb{Q}} \mathbb{G}_{m, F}$, and thus we have a morphism $\det \circ x : \mathbb{S} \rightarrow (T^F)_{\mathbb{R}}$. The pair $(T^F, \det \circ x)$ is a Shimura datum defining a zero dimensional Shimura variety, and the point $\det \circ x$ is a CM point. Recall that $\mathbb{Q}(\psi)$ is the field generated over \mathbb{Q} by the values of ψ on $\mathbb{A}_{F, f}^\times$. Also, let $E(\mu) \supset E(G, X)$ be the reflex field of the automorphic vector bundle \mathcal{E}_μ over S . Define $E(\psi) = E(\mu)E(T^F, \det \circ x)\mathbb{Q}(\psi)$. The infinity type of ψ can be seen as an algebraic character of T^F , and the corresponding automorphic vector bundle \mathcal{E}_ψ has a canonical model over $E(\psi)$. Note that $E(T^F, \det \circ x) \supset E(G, X)$.

Attached to the CM point $\det \circ x$ there is a CM period

$$p(\psi; \det \circ x) \in \mathbb{C}^\times,$$

defined in [HK91] (see also [Har93]). For every $\sigma \in \text{Aut}(\mathbb{C})$, the conjugate Shimura datum is canonically identified with $(T^F, \det \circ (\sigma x))$ (this is clear from the definitions), so we can define as well a CM period $p(\sigma \psi; \det \circ (\sigma x)) \in \mathbb{C}^\times$. If $\sigma \in \text{Aut}(\mathbb{C}/E(\psi))$, then this coincides with $p(\psi; \det \circ x)$, and this

allows us to define $p({}^\rho\psi; \det \circ({}^\rho x))$ for any $\rho \in J_{E(\psi)}$ by extending ρ to an element of $\text{Aut}(\mathbb{C})$. We let

$$p^*(\psi; \det \circ x) = (p({}^\rho\psi; \det \circ({}^\rho x)))_{\rho \in J_{E(\psi)}},$$

viewed as an element of $(E(\psi) \otimes \mathbb{C})^\times$. We also define

$$P(\psi) = P(\psi; x) = p(\psi; \det \circ x)p(\psi^{-1}; \det \circ \bar{x})$$

and

$$P^*(\psi) = P^*(\psi; x) = p^*(\psi; \det \circ x)p^*(\psi^{-1}; \det \circ \bar{x}).$$

Note that this depends on the choice of the CM point x , but we will ignore x for simplicity of notation. If α is another algebraic Hecke character of F , we let

$$P^*(\psi; \alpha) = P^*(\psi; \alpha; x) = p^*(\psi; \det \circ x)p^*(\psi^{-1}\alpha^{-1}; \det \circ \bar{x}) \in E(\psi, \alpha) \otimes \mathbb{C},$$

where $E(\psi, \alpha) = E(\psi)\mathbb{Q}(\alpha)$.

As in [Gue16], 3.10, we let s_ψ be an automorphic form that contributes to $H_1^0(S(\det \circ x)_{\mathbb{C}}, \mathcal{E}_\psi)$, which is rational over $E(\psi)$. Similarly, we let f_0 be an automorphic form in π , contributing to $H_1^d(S_{\mathbb{C}}, \mathcal{E}_\mu)$, rational over $E(\mu)$. We can then form an automorphic form $f = f_0 \otimes s_\psi$ on $\pi \otimes \psi$, and a corresponding non-zero $G(\mathbb{A}_f)$ -equivariant map $\gamma : \pi_{f,0} \otimes E(\psi) \rightarrow H_1^d(S_{E(\psi)}, \mathcal{E})$, where \mathcal{E} is the automorphic vector bundle over $S_{\mathbb{C}}$ obtained by pulling back \mathcal{E}_ψ and taking the tensor product with \mathcal{E}_μ . Concretely, \mathcal{E} is attached to the irreducible representation of $K_{x,\mathbb{C}}$ whose highest weight is $\mu + \mu(\psi)$, where

$$\mu(\psi) = \left((m_\tau - m_{\bar{\tau}}, \dots, m_\tau - m_{\bar{\tau}})_{\tau \in \Phi}; n \sum_{\tau \in \Phi} m_{\bar{\tau}} \right).$$

(see [Gue16], 4.5).

We now let α be another algebraic Hecke character of F , whose infinity type is given at each place $\tau \in \Phi$ by an integer $-\kappa$ (the same for all $\tau \in \Phi$), and at each place $\tau \notin \Phi$ by 0. As similar construction as above, using $\pi^\vee \otimes \alpha^{-1}$ and ψ^{-1} instead of π and ψ , gives rise to elements $s_{\psi^{-1}}$, f'_0 and f' , which in turn are associated with a map γ' to coherent cohomology in degree d of the conjugate Shimura variety ${}^c S_{\mathbb{C}}$. See [Gue16], 4.5, for details. The maps γ and γ' define via cup product and pullback to $S_{\mathbb{C}}^\# \hookrightarrow S_{\mathbb{C}} \times {}^c S_{\mathbb{C}}$, an element $(\gamma, \gamma')^\#$ that contributes to

$$H_1^{2d}(S_{\mathbb{C}}^\#, \mathcal{E}_{(\mu+\mu(\psi), \mu^\vee - \mu(\psi) - \mu(\alpha))^\#}).$$

Note that $\mathcal{E}_{(\mu+\mu(\psi), \mu^\vee - \mu(\psi) - \mu(\alpha))^\#}$ is isomorphic to the Serre dual $\mathcal{E}'_{\Lambda^\#[\kappa]}$ of $\mathcal{E}_{\Lambda^\#[\kappa]}$.

For $\sigma \in \text{Aut}(\mathbb{C})$, we can conjugate s_ψ , $s_{\psi^{-1}}$, f_0 and f'_0 (and hence f and f') to obtain automorphic forms ${}^\sigma f \in {}^\sigma \pi \otimes {}^\sigma \psi$ and ${}^\sigma f' \in {}^\sigma \pi^\vee \otimes {}^\sigma \alpha^{-1} \otimes {}^\sigma \psi^{-1}$. These are also associated with ${}^\sigma G(\mathbb{A}_f)$ -equivariant maps ${}^\sigma \gamma$ and ${}^\sigma \gamma'$, and the same procedure as above gives rise to an element

$${}^\sigma(\gamma, \gamma')^\# = ({}^\sigma \gamma, {}^\sigma \gamma')^\#$$

that contributes to

$$H_1^{2d}({}^\sigma S_{\mathbb{C}}^\#, \mathcal{E}'_{\sigma \Lambda^\#[\kappa]}).$$

We define

$$Q^{\text{Pet}}(f_0) = \int_{Z(\mathbb{A})G(\mathbb{Q})G(\mathbb{A})} f_0(g)\bar{f}_0(g)\|\nu(g)\|^{2a_0}dg,$$

and we define $Q^{\text{Pet}}(\sigma f_0)$ for $\sigma \in \text{Aut}(\mathbb{C})$ in a similar way. We let $E(\pi) = E(\mu)E_0(\pi)$, and $E(\pi, \psi) = E(\pi)E(\psi)$. By (3.1.1), we can define $Q^{\text{Pet}}(\pi) = Q^{\text{Pet}}(f_0)$ uniquely up to multiples by $E(\pi)$. If σ fixes $E(\pi)$ (in particular, if σ fixes $E(\pi, \psi)$), then $Q^{\text{Pet}}(\sigma f_0) = Q^{\text{Pet}}(f_0)$, and hence we can define

$$Q^{\text{Pet},*}(\pi) = (Q^{\text{Pet}}(\rho\pi))_{\rho \in J_{E(\pi, \psi)}} \in E(\pi, \psi) \otimes \mathbb{C}.$$

We also let

$$(f, f') = \int_{Z(\mathbb{A})G(\mathbb{Q})\backslash G(\mathbb{A})} f(g)f'(g)\alpha(\det(g))dg,$$

and get in a similar fashion an element

$$(f, f')^* = (({}^\rho f, {}^\rho f')_{\rho G})_{\rho \in J_{E(\pi, \psi, \alpha)}} \in E(\pi, \psi, \alpha) \otimes \mathbb{C},$$

where $E(\pi, \psi, \alpha) = E(\pi, \psi)\mathbb{Q}(\alpha)$.

Lemma 3.3.1. *Keep the notation and assumptions as above. Then*

$$(f, f')^* \sim_{E(\pi, \psi, \alpha) \otimes F^{\text{Gal}}} (2\pi i)^{2a_0} Q^{\text{Pet},*}(\pi) P^*(\psi; \alpha)^{-1}.$$

Proof. This is completely similar to the computations in Section 2.9 of [Har97]. \square

Remark 3.3.1. The L -function $L^{*, \text{mot}, S}(s, \pi \otimes \psi, \text{St}, \alpha)$ can be seen as valued in $E(\pi, \psi, \alpha) \otimes \mathbb{C}$.

3.4. Eisenstein series and zeta integrals. Let α be an algebraic Hecke character of F as above. For $s \in \mathbb{C}$, let $I(s, \alpha)$ be the induced representation

$$I(s, \alpha) = \{f : G^{(2)}(\mathbb{A}) \rightarrow \mathbb{C} : f(pg) = \delta_{GP, \mathbb{A}}(p, \alpha, s)f(g), g \in G^{(2)}(\mathbb{A}), p \in GP(\mathbb{A})\},$$

where $\delta_{GP, \mathbb{A}}(p, \alpha, s) = \alpha(\det(A(p))) \|N_{L/K} \det A(p)\|_{\mathbb{A}_K}^{\frac{n}{2}+s} \|\nu(p)\|_{\mathbb{A}_K}^{-\frac{n}{2}-ns}$. The local inductions $I(s, \alpha)_v$ and finite and archimedean inductions $I(s, \alpha)_f$ and $I(s, \alpha)_\infty$ are defined similarly. A section of $I(s, \alpha)$ is a function $\phi(\cdot, \cdot)$, that to each $s \in \mathbb{C}$ assigns an element $\phi(\cdot, s) \in I(s, \alpha)$, with a certain continuity property. Local sections are defined similarly. For $\text{Re}(s) \gg 0$, we can define the Eisenstein series

$$E_{\phi, s}(g) = \sum_{\sigma \in GP(\mathbb{Q})\backslash G^{(2)}(\mathbb{Q})} \phi(\sigma g, s),$$

which converges absolutely to an automorphic form on $G^{(2)}(\mathbb{A})$. This extends meromorphically to a function of $s \in \mathbb{C}$.

From now on, fix $m > n - \frac{k}{2}$ an integer satisfying (3.2.1). Let $f \in \pi \otimes \psi$ and $f' \in \pi^\vee \otimes \alpha^{-1} \otimes \psi^{-1}$ as above. For any section ϕ of $I(s, \alpha)$, we define the modified Piatetski-Shapiro-Rallis zeta integral to be

$$Z(s, f, f', \phi) = \int_{Z^\sharp(\mathbb{A})G^\sharp(\mathbb{Q})\backslash G^\sharp(\mathbb{A})} E_{\phi, s}(i(g, g'))f(g)f'(g')dgdg',$$

where Z^\sharp is the center of G^\sharp . Suppose moreover that f, f' and ϕ are factorizable as $\otimes'_v f_v, \otimes'_v f'_v \otimes \alpha_v^{-1}$ and $\prod'_v \phi_v$. Note that we are taking $f'_v \in \pi_v^\vee \otimes \psi_v^{-1}$,

with $f'_v \otimes \alpha_v^{-1}$ the function sending g to $f'_v(g)\alpha^{-1}(\det(g))$. At almost all places v , $\pi_v \otimes \psi_v$ is unramified and f_v and f'_v are normalized spherical vectors of $\pi_v \otimes \psi_v$ and $\pi_v^\vee \otimes \psi_v^{-1}$ respectively, with the local pairing $(f_v, f'_v) = 1$. Define the local zeta integrals as

$$Z_v(s, f, f', \phi) = \int_{U_v} \phi_v(i(h_v, 1), s) c_{f, f', v}(h_v) dh_v,$$

where U_v is the local unitary group at the place v for V , and

$$c_{f, f', v}(h_v) = (f_v, f'_v)^{-1}(\pi_v(h_v)f_v, f'_v)$$

is a normalized matrix coefficient for π_v . We let S be a big enough set of primes of K containing the archimedean primes (in practice we take S to be the set consisting of the archimedean places S_∞ , the places at which G is not quasi-split and the places v where π_v is ramified or f_v or f'_v is not a standard spherical vector). Write $S = S_f \cup S_\infty$, and let

$$Z_f(s, f, f', \phi) = \prod_{v \in S_f} Z_v(s, f, f', \phi)$$

and

$$Z_\infty(s, f, f', \phi) = \prod_{v \in S_\infty} Z_v(s, f, f', \phi).$$

We can conjugate sections ϕ_f by an element $\sigma \in \text{Aut}(\mathbb{C})$ as in the discussion before Lemma 6.2.7 of [Har93].

Lemma 3.4.1. *There exists a finite section $\phi_f(\cdot, s) \in I(s, \alpha)_f$ with $\phi_f(\cdot, m - \frac{n}{2})$ taking values in $\mathbb{Q}(\alpha)$ such that*

$$Z_f\left(m - \frac{n}{2}, f, f', \phi_f\right) \neq 0.$$

Moreover, for any $\sigma \in \text{Aut}(\mathbb{C})$, we have

$$\sigma\left(Z_f\left(m - \frac{n}{2}, f, f', \phi_f\right)\right) = Z_f\left(m - \frac{n}{2}, \sigma f, \sigma f', \sigma \phi_f\right).$$

In particular,

$$Z_f\left(m - \frac{n}{2}, f, f', \phi_f\right) \in E(\pi, \psi, \alpha).$$

Proof. The existence of ϕ_f with the first property follows as in Lemma 4.5.2 of [Gue16] or Lemma 3.5.7 of [Har97] (see also the proof of Theorem 4.3 of [Har08]). The description of the action of σ follows from Lemma 6.2.7 of [Har93]. \square

From now on, fix ϕ_f as in Lemma 3.4.1. We consider the element $G(\alpha) \in E(\pi, \psi, \alpha) \otimes \mathbb{C}$ defined in Subsection 2.5, and denote its ρ -component by $G(\alpha)_\rho$, for $\rho : E(\pi, \psi, \alpha) \hookrightarrow \mathbb{C}$. If $\sigma \in \text{Aut}(\mathbb{C})$, we let $G(\alpha)_\sigma = G(\alpha)_\rho$, where ρ is the restriction of σ to $E(\pi, \psi, \alpha)$. Define a section $\varphi_{m, \kappa, \sigma}$ of $I\left(s + m - \frac{n}{2}, \sigma \alpha\right)$ by

$$\varphi_{m, \kappa, \sigma}(g, s) = \mathbb{J}_{m, \kappa}\left(g, s + m - \frac{n}{2}\right) \otimes (2\pi i)^{[F^+ : \mathbb{Q}](m + \kappa)n} G(\alpha)_\sigma^n(\sigma \phi_f)(g, s + m - \frac{n}{2}).$$

The element $\mathbb{J}_{m, \kappa}$ is defined in [Har07], (1.2.7) (with a misprint correction, see [Gue16], 4.3). The Eisenstein series $E_{m, \kappa} = E_{m, \kappa, 1} = E_{\varphi_{m, \kappa, 1}}$ has no pole at $s = 0$ (see for example (1.2.5) of [Har08], where $\chi = \alpha \|N_{F/F^+}\|^{-\kappa/2}$), and

thus this defines an automorphic form, also denoted by $E_{m,\kappa}$, on $G^{(2)}(\mathbb{A})$, which can be seen as an element of $H^0(S_{\mathbb{C}}^{(2)}, \mathcal{E}_{m,\kappa}^{\text{can}})$ (see [Gue16], 4.3). Using the differential operators $\Delta_{m,\kappa}$, as explained in *op. cit.*, we can define sections

$$\tilde{\varphi}_{m,\kappa,\sigma} = \Delta_{m,\kappa} \varphi_{m,\kappa,\sigma}$$

for $\sigma \in \text{Aut}(\mathbb{C})$, and a corresponding Eisenstein series

$$\tilde{E}_{m,\kappa} = E_{\tilde{\varphi}_{m,\kappa,1}}.$$

Then $\tilde{E}_{m,\kappa}$ equals $\Delta_{m,\kappa} E_{m,\kappa}$ when restricted to $G^{\sharp}(\mathbb{A})$.

Proposition 3.4.1. *The Eisenstein series $E_{m,\kappa}$ and $\tilde{E}_{m,\kappa}$ are rational over $\mathbb{Q}(\alpha)$ with respect to the canonical models of $S_{\mathbb{C}}^{(2)}$ and $\mathcal{E}_{m,\kappa}$. Moreover, for any $\sigma \in \text{Aut}(\mathbb{C})$,*

$${}^{\sigma} E_{m,\kappa} = E_{\varphi_{m,\kappa,\sigma}},$$

and a similar equation holds for $\tilde{E}_{m,\kappa}$.

Proof. This follows by combining the ideas of Lemma 3.3.5.3 of [Har97] and Proposition 4.3.1 of [Gue16]. Namely, in the latter, we just need to note that the character $\tilde{\lambda}$ is now given by

$$\tilde{\lambda}(p) = (N_{F/\mathbb{Q}} \det(A(p)))^{-m} \nu(p)^{-[F^+:\mathbb{Q}]nm} t_{\alpha}(\det(A(p)))^{-1},$$

where t_{α} is the algebraic character of $\text{Res}_{F/\mathbb{Q}} \mathbb{G}_{m,F}$, defined over $\mathbb{Q}(\alpha)$, inverse of the infinity type of α , so that the restriction to $\text{Sh}(\mathbb{G}_{m,\mathbb{Q}}, N)$ is the Tate automorphic vector bundle $\mathbb{Q}(-[F^+:\mathbb{Q}]n(m+\kappa))$. \square

We define

$$Z^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) = \left(Z \left(m - \frac{n}{2}, {}^{\sigma} f, {}^{\sigma} f', \tilde{\varphi}_{m,\kappa,\sigma} \right) \right)_{\sigma \in \text{Aut}(\mathbb{C})}.$$

The elements of this family only depend on the restrictions of elements $\sigma \in \text{Aut}(\mathbb{C})$ to $E(\pi, \psi, \alpha)$, and hence we can consider

$$Z^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) = \left(Z \left(m - \frac{n}{2}, {}^{\rho} f, {}^{\rho} f', \tilde{\varphi}_{m,\kappa,\rho} \right) \right)_{\rho \in J_{E(\pi,\psi,\alpha)}}$$

as an element of $E(\pi, \psi, \alpha) \otimes \mathbb{C}$. We can also define

$$Z_{\infty}^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) = \left(Z_{\infty} \left(m - \frac{n}{2}, {}^{\rho} f, {}^{\rho} f', \tilde{\varphi}_{m,\kappa,\rho} \right) \right)_{\rho \in J_{E(\pi,\psi,\alpha)}},$$

which is an element of $E(\pi, \psi, \alpha) \otimes \mathbb{C}$. Note that the archimedean part of $\tilde{\varphi}_{m,\kappa,\rho}$ is independent of ρ , and hence so are the archimedean zeta integrals. Finally, we can define

$$Z_f^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) = \left(Z_f \left(m - \frac{n}{2}, {}^{\rho} f, {}^{\rho} f', \tilde{\varphi}_{m,\kappa,\rho} \right) \right)_{\rho \in J_{E(\pi,\psi,\alpha)}}$$

and $Z_f^* \left(m - \frac{n}{2}, f, f', \phi_f \right)$.

Lemma 3.4.2. *Let the notation and assumptions be as above. Then*

$$Z^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) \in E(\pi, \psi, \alpha).$$

Proof. Let

$$(3.4.1) \quad \mathcal{L}_{m,\kappa} : H^{2d}(S_{\mathbb{C}}^{\sharp}, \mathcal{E}'_{\Lambda^{\sharp}[\kappa]}) \rightarrow \mathbb{C}$$

be the map defined by pairing with $\Delta_{m,\kappa} E_{m,\kappa}$ via Serre duality. Then, as in Lemma 4.5.3 of [Gue16], we have

$$Z\left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa,1}\right) = \mathcal{L}_{m,\kappa}\left((\gamma, \gamma')^{\sharp}\right).$$

Let $\sigma \in \text{Aut}(\mathbb{C})$. The conjugate of (3.4.1) by σ is now

$$\mathcal{L}_{m,\kappa} : H^{2d}(\sigma S_{\mathbb{C}}^{\sharp}, \mathcal{E}'_{\sigma\Lambda^{\sharp}[\kappa]}) \rightarrow \mathbb{C},$$

which is given by cup product with ${}^{\sigma}\Delta_{m,\kappa} E_{m,\kappa}$ via Serre duality. It then follows that

$$\sigma\left(Z\left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa,1}\right)\right) = \mathcal{L}_{m,\kappa}\left(\sigma(\gamma, \gamma')^{\sharp}\right),$$

which equals

$$Z\left(m - \frac{n}{2}, {}^{\sigma}f, {}^{\sigma}f', {}^{\sigma}\tilde{\varphi}_{m,\kappa,\sigma}\right)$$

by Proposition 3.4.1 and the same reasoning as above. This finishes the proof of the lemma. \square

The main formula for the doubling method, proved by Li in [Li92], says that

$$(3.4.2) \quad d^S\left(s - \frac{n}{2}, \alpha\right) Z\left(s - \frac{n}{2}, f, f', \phi\right) = (f, f') \prod_{v \in S} Z_v\left(s - \frac{n}{2}, f, f', \phi\right) L^{\text{mot}, S}(s, \pi \otimes \psi, \text{St}, \alpha)$$

for any section ϕ . Here

$$d^S(s, \alpha) = \prod_{j=0}^{n-1} L^S(2s + n - j, \alpha|_{\mathbb{A}_{F^+}^{\times}} \varepsilon_F^j),$$

where ε_F is the quadratic character associated with the quadratic extension F/F^+ . We can write

$$\alpha|_{\mathbb{A}_{F^+}^{\times}} = \alpha_0 \|\cdot\|_{\mathbb{A}_{F^+}^{\times}}^{\kappa}$$

with α_0 of finite order, so that

$$d^S(s, \alpha) = \prod_{j=0}^{n-1} L^S(2s + n - j + \kappa, \alpha_0 \varepsilon_F^j).$$

We let

$$d^*(s, \alpha) = \prod_{j=0}^{n-1} L^*(2s + n - j + \kappa, \alpha_0 \varepsilon_F^j) \in \mathbb{Q}(\alpha_0) \otimes \mathbb{C},$$

and we define similarly $d^{*,S}(s, \alpha)$ by removing the local factors at primes of S . We can deduce from (3.4.2) that

$$(3.4.3) \quad d^{*,S}\left(m - \frac{n}{2}, \alpha\right) Z^*\left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa}\right) = (f, f')^* Z_f^*\left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa}\right) Z_{\infty}^*\left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa}\right) L^{*,\text{mot}, S}(m, \pi \otimes \psi, \text{St}, \alpha)$$

Lemma 3.4.3. *We have*

$$d^{*,S} \left(m - \frac{n}{2}, \alpha \right) \sim_{\mathbb{Q}(\alpha_0)} (2\pi i)^{[F^+:\mathbb{Q}] \left((2m+\kappa)n - \frac{n(n-1)}{2} \right)} D_{F^+}^{\lfloor \frac{n+1}{2} \rfloor / 2} \delta[\varepsilon_F]^{\lfloor \frac{n}{2} \rfloor} G(\alpha)^n.$$

Proof. First, suppose that $0 \leq j \leq n-1$ is even. Note that $2m-j+\kappa$ is even and positive, and hence is a critical integer of $L^*(s, \alpha_0)$, because the motive $\text{Res}_{F^+/\mathbb{Q}}[\alpha_0]$ is purely of type $(0,0)$ and the Frobenius involution acts as $(-1)^\kappa$. Since Deligne's conjecture is known for $\text{Res}_{F^+/\mathbb{Q}}[\alpha_0]$, we get

$$L^*(2m-j+\kappa, \alpha_0) \sim_{\mathbb{Q}(\alpha_0)} (2\pi i)^{[F^+:\mathbb{Q}](2m-j+\kappa)} c^\pm[\alpha_0],$$

where $\pm = (-1)^{2m-j+\kappa} = (-1)^\kappa$. Here we are writing $c^\pm[\alpha_0] = c^\pm(\text{Res}_{F^+/\mathbb{Q}}[\alpha_0])$. Similarly, if $0 \leq j \leq n-1$ is odd, then $2m-j+\kappa$ is a critical integer for the motive $\text{Res}_{F^+/\mathbb{Q}}[\alpha_0 \varepsilon_F]$ and

$$L^*(2m-j+\kappa, \alpha_0 \varepsilon_F) \sim_{\mathbb{Q}(\alpha_0)} (2\pi i)^{[F^+:\mathbb{Q}](2m-j+\kappa)} c^\mp[\alpha_0 \varepsilon_F].$$

We know use Remark 2.2.1 of [Gue16], together with Proposition 2.2 of [Yos94], to get

$$c^\pm[\alpha_0] \sim_{\mathbb{Q}(\alpha_0)} \delta[\alpha_0]$$

and

$$c^\mp[\alpha_0 \varepsilon_F] \sim_{\mathbb{Q}(\alpha_0)} \delta[\alpha_0] \delta[\varepsilon_F] D_{F^+}^{-1/2}.$$

The lemma follows from these computations, combined with (2.5.1) and the fact that

$$d^{*,S} \left(m - \frac{n}{2}, \alpha \right) \sim_{\mathbb{Q}(\alpha)} d^* \left(m - \frac{n}{2}, \alpha \right).$$

□

3.5. Modified periods. A theorem of Garrett ([Gar08]) says that the archimedean zeta integral $Z_\infty \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_m \right)$ is non-zero (and, moreover, belongs to F^{Gal}), and we define the modified (Petersson) period

$$\mathfrak{Q}^{\text{Pet}}(m; \pi, \psi, \alpha) = \frac{(f, f')^{-1}}{Z_\infty \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right)}.$$

It follows from Lemma 3.3.1 that

$$\mathfrak{Q}^{\text{Pet}}(m; \pi, \psi, \alpha) \sim_{E(\pi, \psi, \alpha) \otimes F^{\text{Gal}}} (2\pi i)^{-2a_0} Q^{\text{Pet}}(\pi)^{-1} P(\psi; \alpha).$$

More generally, we can define $\mathfrak{Q}^{\text{Pet},*}(m; \pi, \psi, \alpha) \in (E(\pi, \psi, \alpha) \otimes \mathbb{C})^\times$ as

$$\mathfrak{Q}^{\text{Pet},*}(m; \pi, \psi, \alpha) = \left(\mathfrak{Q}^{\text{Pet}}(m; \pi^\rho, \psi^\rho, \alpha^\rho) \right)_{\rho \in J_{E(\pi, \psi, \alpha)}} = \frac{(f, f')^{*, -1}}{Z_\infty^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right)}.$$

The doubling zeta integral against the Eisenstein series $\tilde{E}_{m,\kappa}$ defines a bilinear form

$$B^\alpha : H_!^d(S_{\mathbb{C}}, \mathcal{E})[\pi \otimes \psi] \times H_!^d(\bar{S}_{\mathbb{C}}, \mathcal{E}^*)[\pi^\vee \otimes \psi^{-1} \otimes \alpha^{-1}] \rightarrow \mathbb{C},$$

which is moreover rational over $E(\pi, \psi, \alpha)$. Here \mathcal{E} and \mathcal{E}^* are the automorphic vector bundles determined by π and $\pi^\vee \otimes \alpha^{-1}$. In particular, with our choice of f and f' , we have that $B^\alpha(f, f') \in E(\pi, \psi, \alpha)$. Moreover, for any $\sigma \in \text{Aut}(\mathbb{C})$,

$$(3.5.1) \quad \sigma(B^\alpha(f, f')) = B^{\sigma\alpha}(\sigma f, \sigma f').$$

By our multiplicity assumptions, any other bilinear form, such as the Petersson integral, must be a scalar multiple of B^α . In particular, there exists an element $Q(\pi, \psi, \alpha) \in \mathbb{C}^\times$ such that

$$B^\alpha(f, f') = Q(\pi, \psi, \alpha)(f, f').$$

We can define $Q^*(\pi, \psi, \alpha) \in (E(\pi, \psi, \alpha) \otimes \mathbb{C})^\times$ by taking

$$Q^*(\pi, \psi, \alpha) = (Q(\rho\pi, \rho\psi, \rho\alpha))_{\rho \in J_{E(\pi, \psi, \alpha)}}.$$

By (3.5.1), we have that

$$(3.5.2) \quad (f, f')^{*, -1} \sim_{E(\pi, \psi, \alpha)} Q^*(\pi, \psi, \alpha).$$

Remark 3.5.1. In the above computations, $Q(\pi, \psi, \alpha)$ depends a priori on the Eisenstein series $\tilde{E}_{m, \kappa}$, and hence on the integer m . However, (3.5.2) shows that, up to multiplication by an element in $E(\pi, \psi, \alpha) \subset E(\pi, \psi, \alpha) \otimes \mathbb{C}$, $Q^*(\pi, \psi, \alpha)$ does not depend on m or the Eisenstein series.

We define

$$\mathfrak{Q}^*(m; \pi, \psi, \alpha) = \frac{Q^*(\pi, \psi, \alpha)}{Z_\infty^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m, \kappa}\right)}.$$

We have that

$$\mathfrak{Q}^{\text{Pet}, *}(m; \pi, \psi, \alpha) \sim_{E(\pi, \psi, \alpha)} \mathfrak{Q}^*(m; \pi, \psi, \alpha).$$

3.6. The main theorem. Before stating our main theorem, we recall all the hypothesis and assumptions that we have made so far. Thus, F/F^+ is a CM extension, Φ is a CM type, and π is an automorphic representation of $G(\mathbb{A})$, satisfying hypotheses 2.3.1 for a parameter $\mu = ((a_{\tau, 1}, \dots, a_{\tau, n})_{\tau \in \Phi}; a_0)$ (recall as well the assumption that π can be conjugated to $\sigma\pi$ with the desired properties). We also assume that $\pi^\vee \cong \pi \otimes \|\nu\|^{2a_0}$, π contributes to antiholomorphic cohomology and satisfies the multiplicity assumption (3.1.1).

We also have algebraic Hecke characters ψ and α of F . The infinity type of ψ is $(m_\tau)_{\tau \in J_F}$, and that of α is given by an integer κ at places of Φ , and by 0 at places outside Φ . We define the number field $E(\pi, \psi, \alpha)$ as in Subsection 3.3.

Theorem 3.6.1. *Keep the notation and assumptions as above, and let $m > n - \frac{n}{2}$ be an integer satisfying (3.2.1). Then*

$$L^{*, \text{mot}, S}(m, \pi \otimes \psi, \text{St}, \alpha) \sim_{E(\pi, \psi, \alpha)} (2\pi i)^{[F^+:\mathbb{Q}](mn - n(n-1)/2)} D_{F^+}^{\lfloor \frac{n+1}{2} \rfloor / 2} \delta_{[E_F]}^{\lfloor \frac{n}{2} \rfloor} \mathfrak{Q}^*(m; \pi, \psi, \alpha).$$

Proof. We use formula (3.4.3). By Lemma 3.4.1, we have that

$$Z_f^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m, \kappa}\right) \sim_{E(\pi, \psi, \alpha)} (2\pi i)^{[F^+:\mathbb{Q}](m+\kappa)n} G(\alpha)^n.$$

Also, Lemma 3.4.2 says that

$$Z^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m, \kappa}\right) \sim_{E(\pi, \psi, \alpha)} 1.$$

The formula in the theorem follows immediately from these, Lemma 3.4.3 and the definition of $\mathfrak{Q}(m; \pi, \psi, \alpha)$. \square

3.7. A refinement. Lemma 3.3.1 gives us a factorization of (f, f') in terms of periods associated to π , ψ and α respectively. This will lead to a finer result on the special values as Theorem 1 in [Gue16].

Unfortunately, the relation in Lemma 3.3.1 is only shown under $\text{Gal}(\overline{\mathbb{Q}}/F^{\text{Gal}})$ -action. We hope to prove a $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ version in the near future.

Before we state the main formula, let us define two more factors which will appear.

Definition 3.7.1. (1) Let $j \in F$ be a purely imaginary element, i.e., $\bar{j} = -j$ where \bar{j} refers to the complex conjugation of j in the CM field F . We define

$$J_F = \prod_{\tau \in \Phi} \tau(j).$$

Its image in $\mathbb{C}^\times/\mathbb{Q}^\times$ does not depend on the choice of the purely imaginary element j or the CM type Φ .

(2) Let E be a number field containing F^{Gal} **a verifier, if not, we should add n in the definition of sign and a difficult lemma to show that this is well-defined.** We fix ρ_0 an element in J_E . For any $\rho \in J_E$, we define a sign $e_\Phi(\rho)$ by $(-1)^{\#\langle \Phi \setminus g\Phi \rangle}$ by taking any $g \in \text{Aut}(\mathbb{C})$ such that $g\rho_0 = \rho$. We can see easily that it does not depend on the choice of g .

We define $e_\Phi = (e_\Phi(\rho))_{\rho \in J_E}$ as an element of $E \otimes \mathbb{C}$.

Corollary 3.7.1. *With the same assumption as in Theorem 3.6.1, we have that*

$$L^{*,\text{mot},S}(m, \pi \otimes \psi, \text{St}, \alpha) \sim_{E(\pi, \psi, \alpha); F^{\text{Gal}}} (2\pi i)^{[F^+:\mathbb{Q}](mn-n(n-1)/2)-2a_0} \mathfrak{J}_F^{[n/2]} (D_{F^+}^{1/2})^n e_\Phi^{mn} Q^*(\pi)^{-1} P^*(\psi, \alpha).$$

Remark 3.7.1. (1) We identify $(2\pi i)^{mnd(F^+)} \mathfrak{J}_F^{[n/2]} (D_{F^+}^{1/2})^n$ with

$$1 \otimes (2\pi i)^{mnd(F^+)} \mathfrak{J}_F^{[n/2]} (D_{F^+}^{1/2})^n$$

as an element in $E(\Pi, \eta) \otimes \mathbb{C}$.

(2) It is not difficult to see that if $g \in \text{Aut}(\mathbb{C})$ fixes F^{Gal} then it fixes

$$\mathfrak{J}_F^{[n/2]} (D_{F^+}^{1/2})^n e_\Phi^{mn}.$$

Consequently, it can be ignored here since we consider relations under $\text{Gal}(\overline{\mathbb{Q}}/F^{\text{Gal}})$ -action. We keep them here because they are predicted by Deligne's conjecture if we want a finer result under $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ -action. We will discuss this more in Section 5.

Proof. By the proof of Lemma 2.4.2 of [Gue16] and Proposition 2.2 of [Yos94] we know that

$$\delta[\varepsilon_F] \sim_{\mathbb{Q}} J_F D_{F^+}^{1/2}.$$

Moreover, Lemma 3.3.1 implies that

$$\mathfrak{J}^*(m; \pi, \psi, \alpha) \sim_{E(\pi, \psi, \alpha) \otimes F^{\text{Gal}}} (2\pi i)^{-2a_0} Q^*(\pi)^{-1} P^*(\psi, \alpha).$$

It remains to show that $e_\Phi \in E \otimes F^{\text{Gal}}$. In fact, let $\rho \in J_E$ and $g \in \text{Aut}(\mathbb{C}/F^{\text{Gal}})$, we are going to show $e_\Phi \in E \otimes F^{\text{Gal}}$. We take $h \in \text{Aut}(\mathbb{C})$ such that $\rho = h\rho_0$. By definition $e_\Phi(\rho) = (-1)^{\#\langle \Phi \setminus h\Phi \rangle}$ and $e_\Phi(g\rho) =$

$(-1)^{\#(\Phi \setminus gh\Phi)}$. Since g fixes F^{Gal} , we have $gh\Phi = h\Phi$ and hence $e_{\Phi}(g\rho) = e_{\Phi}(\rho)$. We conclude that $e_{\Phi} \in E \otimes F^{\text{Gal}}$ by Definition-Lemma 1.1 of [LIN15a]. The corollary then follows from Theorem 3.6.1. \square

Remark 3.7.2. We expect that Lemma 3.3.1 is true up to factors in $E(\pi, \psi, \alpha)$. **or change the periods to avoid this point** Moreover, we hope to show that

$$Z_{\infty}^* \left(m - \frac{n}{2}, f, f', \tilde{\varphi}_{m,\kappa} \right) \sim_{E(\pi, \psi, \alpha)} e_{\Phi}^{mn}.$$

By the method explained in section 9.4 of [LIN15c] and Blasius's proof of Deligne's conjecture for algebraic Hecke characters ([Bla86]), we can reduce to show that certain archimedean zeta integral belongs to \mathbb{Q} . Garrett proved this for particular cases (see [Gar08]). We hope to show the two points in the future and then the above corollary is true up to $E(\pi, \psi, \alpha)$.

4. APPLICATIONS TO GENERAL LINEAR GROUPS

4.1. Transfer from similitude unitary groups to unitary groups.

Let π be an automorphic representation of $G(V)(\mathbb{A})$. We want to consider the restriction of π to $U(V)(\mathbb{A})$. We sketch the construction of [LS86] in our case.

Definition 4.1.1. (1) Let π_1 and π_2 be two admissible irreducible representation of $G(V)(\mathbb{A})$. We say they are \mathcal{E} -equivalent if there exists a character χ of $U(V)(\mathbb{A}) \backslash G(V)(\mathbb{A})$ such that $\pi_1 \cong \pi_2 \otimes \chi$.
 (2) Let π_0 be an admissible irreducible representation of $U(V)(\mathbb{A})$ and g be an element in $G(V)(\mathbb{A})$. We define π^g , a new representation on $U(V)(\mathbb{A})$, by $\pi^g(x) = \pi(gxg^{-1})$.
 (3) Let $\pi_{0,1}$ and $\pi_{0,2}$ be two admissible irreducible representation of $U(V)(\mathbb{A})$. We say they are \mathcal{L} -equivalent if there exists $g \in G(V)(\mathbb{A})$ such that $\pi_{0,1} \cong \pi_{0,2}^g$.

Lemma 4.1.1. *Let π be an admissible irreducible representation of $G(V)(\mathbb{A})$. The restriction of π to $U(V)(\mathbb{A})$ is a direct sum of admissible irreducible representations in the same \mathcal{L} -equivalence class. This gives a bijection of the \mathcal{E} -equivalence classes of admissible irreducible representations of $G(V)(\mathbb{A})$ and the \mathcal{L} -equivalence classes of admissible irreducible representations of $G(V)(\mathbb{A})$.*

Moreover, if we restrict to the cuspidal spectrum then we get a bijection on equivalence classes of cuspidal representations of both sides.

Proof. The proof is the similar as in Lemma 3.3 and Proposition 3.5 of [LS86] for the special linear group. More details for unitary groups can be found in section 5 of [Clo91]. We sketch the idea there for the last statement.

We write S for the maximal split central torus of G . It is isomorphic to \mathbb{G}_m . Its intersection with U is then isomorphic to $\mu_2 \subset \mathbb{G}_m$. As in [Clo91], we denote this intersection by M .

Let ω be a Hecke character of $S(\mathbb{Q}) \backslash S(\mathbb{A})$. We write ω_0 for its restriction to $M(\mathbb{Q}) \backslash M(\mathbb{A})$. The space of cuspidal forms $L_0^2(U(\mathbb{Q}) \backslash U(\mathbb{A}), \omega_0)$ is endowed with an action of

$$G^1 := G(\mathbb{Q})S(\mathbb{A})U(\mathbb{A})$$

where $G(\mathbb{Q})$ acts by conjugation, $S(\mathbb{A})$ acts via ω and $U(\mathbb{A})$ acts by right translation. We know G^1 is a closed subgroup of $G(\mathbb{A})$ and the quotient $G(\mathbb{A})/G^1$ is compact. The representation of $G(\mathbb{A})$ given by right translation on the cuspidal spectrum $L_0^2(G(\mathbb{Q})\backslash G(\mathbb{A}), \omega)$ is nothing but

$$\text{Ind}_{G^1}^{G(\mathbb{A})} L_0^2(U(\mathbb{Q})\backslash U(\mathbb{A}), \omega_0).$$

□

Remark 4.1.1. Let π be a cuspidal representation of $GU(\mathbb{A})$. Each constituent in the restriction of π to $U(\mathbb{A})$ has the same unramified components. In particular, they all have the same partial L -function.

Lemma 4.1.2. *Let π_0 be an algebraic cuspidal automorphic representation of $U(\mathbb{A})$. We can always extend it to an algebraic cuspidal automorphic representation of $G(\mathbb{A})$.*

Moreover, if π_0 is tempered at some place, discrete series at some place, or cohomological, then its extension has the same property.

Proof. For the extension, we only need to extend the central character of π_0 to an algebraic Hecke character of $S(\mathbb{Q})\backslash S(\mathbb{A})$ by the above lemma.

In fact, since $M(\mathbb{Q})\backslash M(\mathbb{A})$ is compact, the central character of π_0 is always unitary. Hence it lives in the Pontryagin dual of $S(\mathbb{Q})\backslash S(\mathbb{A})$. We know the Pontryagin dual is an exact functor. Therefore, we can extend it to a unitary Hecke character of $M(\mathbb{Q})\backslash M(\mathbb{A})$. This unitary Hecke character is not necessarily algebraic. Twisting by a real power of the absolute value, we can get an algebraic Hecke character of $S(\mathbb{Q})\backslash S(\mathbb{A})$, which is still an extension of the central character of π_0 .

To show the extension is locally tempered or discrete series if π_0 is, it is enough to notice that for any place v of \mathbb{Q} , $M(\mathbb{Q}_v)\backslash U(\mathbb{Q}_v)$ is a finite index subgroup of $S(\mathbb{Q}_v)\backslash G(\mathbb{Q}_v)$.

For the cohomological property, we refer to (5.18) of [Clo91]. □

4.2. Base change for unitary groups. Recall U is the restriction to \mathbb{Q} of the unitary group over F^+ associated to V . We denote the latter by U_0 . Let π_0 be a cuspidal automorphic representation of $U(\mathbb{A}) = U_0(\mathbb{A}_{F^+})$. Since $U_0(V)(\mathbb{A}_F) \cong GL_n(\mathbb{A}_F)$. By Langlands functoriality, we expect to associate π_0 with a $GL_n(\mathbb{A}_F)$ -representation with expected local components.

More precisely, we can describe the unramified representations at local non-archimedean places by the Satake parameters. We refer to [Min11] for more details. The local base change can be then defined explicitly in terms of the Satake parameters. Let l/k be an extension of local non-archimedean fields and H be a connected reductive group over k . The unramified local base change is a map from the set of isomorphism classes of unramified representations of $H(k)$ to that of $H(l)$. In the global settings, let L/K be an extension of global field and H be a connected reductive group over K . We say that an automorphic representation of $H(\mathbb{A}_L)$ is a weak base change of an automorphic representation of $H(\mathbb{A}_K)$ if it is the local unramified base change at almost every finite unramified places. We refer to section 26 of [Art03] for more details on Langlands functoriality.

The base change for unitary groups is almost completely clear thanks to Kaletha-Minguez-Shin-White ([KMSW14]) and their subsequent articles.

But we don't find a precise statement in their paper for our purpose. We will use the results and arguments in [Lab11]. The following proposition is a slight variation of Théorème 5.4 of [Lab11].

Proposition 4.2.1. *Let Π be a cohomological, conjugate self-dual cuspidal representation of $GL_n(\mathbb{A}_F)$. Then Π is a weak base change of π_0 , a cohomological discrete series representation of $U_0(\mathbb{A}_{F^+})$ such that the infinitesimal character of Π_∞ is compatible with the infinitesimal character of π_∞ by base change. *and also at unramified local places?**

We know π_0 is also cuspidal. Moreover, if Π has regular highest weight, then so is π_0 . In this case, $\pi_{0,\infty}$ is a discrete series representation. If the above blue text is OK, then we can have multiplicity one if (1) the two assumptions in Labesse are satisfied (2) Π is regular.

Proof. The existence of π_0 is proved in [Lab11]. There are two additional assumptions in the beginning of section 5.2 of *loc.it* but they are only used for showing the multiplicity one in the *loc.it*.

The compatibility of infinitesimal characters is also proved in *loc.it* by the calculation on transfer of Lefschetz function.

We now show that π_0 is cuspidal. Let v be a split place of F^+ and w be a place of F above v such that Π_w is the local unramified base change of $\pi_{0,v}$. In particular, we have $U(F_v^+) \cong U(F_w) \cong GL_n(F_w)$. We know Π_w is tempered by the Ramanujan conjecture proved in this case by Clozel ([Clo12]) and also by Cariani ([Car12]). Hence $\pi_{0,v}$ is tempered since it is isomorphic to Π_w if we identify $U(F_v^+)$ with $GL_n(F_w)$. The cuspidality then follows from a theorem of Wallach (c.f. [Wal84]) generalized by Clozel (c.f. [Clo93]).

Finally, it is clear that if the highest weight of Π is regular then so is π_0 . We know a cohomological representation of regular weight is discrete series at infinity by Prop. 4.2 and 5.2 of [LS04]. \square

This is the going down part of the base change for unitary groups. We also state the going up part which has been used before **refer to previous calculations on infinity type of Aut twist, check if already assumed the very regular condition, otherwise need some change**

Proposition 4.2.2. *Let π_0 be a cuspidal, cohomological representation of $U_0(\mathbb{A}_{F^+})$. We assume that the highest weight associated to π_0 is very regular. Then there exists Π , a cohomological representation of $GL_n(\mathbb{A}_F)$ which is a weak base change of π_0 . Moreover, Π is the unramified local base change of π_0 at unramified places and the infinitesimal character of Π_∞ is compatible with that of $\pi_{0,\infty}$ by base change.*

Proof. This is exactly the Corollaire 5.3 of [Lab11]. We recall that the condition (*) in the *loc.cit* is satisfied since the highest weight associated to π_0 is very regular. \square

4.3. Special values of representations of general linear group. Let Π be a cohomological conjugate self-dual cuspidal representation of $GL_n(\mathbb{A}_F)$.

For each $\tau \in \Phi$, let s_τ be an integer in $\{0, 1, \dots, n\}$. We write $I := (s_\tau)_{\tau \in \Phi}$ be an element in $\{0, 1, \dots, n\}^\Phi$. Let V_I be a Hermitian space with respect

to F/F^+ of signature $(n - s_\tau, s_\tau)$. We write $U_{0,I}$ for the associated unitary group over F^+ and GU_I for the associated rational similitude unitary group.

We assume that Π^\vee , the contragredient of Π , descends by base change to a packet of representations of $U_{0,I}(\mathbb{A}_{F^+})$, which contains a representation $\pi_{0,I}$ satisfying Hypothesis 2.3.1.

By Lemma 4.1.2, we can extend $\pi_{0,I}$ to π_I , a cuspidal representation of $GU_I(\mathbb{A})$, which still satisfies Hypothesis 2.3.1.

Remark 4.3.1. By Proposition 4.2.1, we know if Π is cohomological with respect to a regular highest weight then it descends by base change to a cuspidal representation of $U_{0,I}(\mathbb{A}_{F^+})$ which is cohomological with respect to a regular highest weight. In particular, this representation satisfies Hypothesis 2.3.1.

Definition 4.3.1. Let Π be as before. Let $I = (s_\sigma)_{\sigma \in \Sigma} \in \{0, 1, \dots, n\}^\Sigma$. We keep the above notation and define the **automorphic arithmetic period** $P^{(I)}(\pi)$ by $(2\pi i)^{-2a_0} Q_{V_I}(\pi)^{-1}$.

Theorem 4.3.1. *Let Π be as before. We denote the infinity type of Π at $\tau \in \Phi$ by $(z^{a_i(\tau)} \bar{z}^{-a_i(\tau)})_{1 \leq i \leq n}$.*

Let η be an algebraic Hecke character of F with infinity type $z^{a(\tau)} \bar{z}^{b(\tau)}$ at $\tau \in \Phi$. We know that $a(\tau) + b(\tau)$ is a constant independent of τ , denoted by $-\omega(\eta)$.

We suppose that $a(\tau) - b(\tau) + 2a_i(\tau) \neq 0$ for all $1 \leq i \leq n$ and $\tau \in \Phi$. We define $I := I(\Pi, \eta)$ to be the map on Φ which sends $\tau \in \Phi$ to $I(\tau) := \#\{i : a(\tau) - b(\tau) + 2a_i(\tau) < 0\}$. As before, we write $P^{,(I(\Pi, \eta))}(\Pi)$ for $(P^{(I(\rho_{\Pi, \eta})})^{(\rho_{\Pi, \eta})})_{\rho \in J_{E(\Pi, \eta)}} \in E(\Pi, \eta) \otimes \mathbb{C}$.*

Let $m \in \mathbb{Z} + \frac{n-1}{2}$. If $m \geq \frac{n + \omega(\eta)}{2}$ satisfies equation (3.2.1), then we have:

$$(4.3.1) \quad L^*(m, \Pi \otimes \eta) \sim_{E(\Pi, \eta); F^{gal}} (2\pi i)^{m d(F^+)} \mathfrak{J}_F^{[n/2]} (D_{F^+}^{1/2})^n e_{\mathbb{F}}^{mn} P^{*,(I(\Pi, \eta))}(\Pi) \prod_{\tau \in \Phi} p^*(\check{\eta}, \tau)^{I(\tau)} p^*(\check{\eta}, \bar{\tau})^{n-I(\tau)}$$

where $d(F^+)$ is the degree of F^+ over \mathbb{Q} .

Remark 4.3.2. (1) The infinity type stated in the theorem is different from the infinity type in subsection 2.5. Previously when we say ψ be an algebraic Hecke character of F , of infinity type $(m_\tau)_{\tau \in J_F}$ we mean ψ is of infinity type $z^{-m_\tau} \bar{z}^{-m_\tau}$ at $\tau \in J_F$ here.
(2) This theorem is first stated as Theorem 5.2.1 in [LIN15c]. It was proved by assuming a conjecture (c.f. Conjecture 5.1.1 of *loc.cit*) which is nothing but a variation of our Theorem 3.7.1.

Proof. **to be fixed after generalization of α** Let ψ be an algebraic Hecke character of F with infinity type

Moreover, we know

$$L^S \left(m - \frac{n-1}{2}, \pi \otimes \psi, \text{St} \right) = L^S \left(m - \frac{n-1}{2}, BC(\pi|_U) \otimes \check{\psi} \right)$$

where $\tilde{\psi} := \psi/\psi^c$ is a Hecke character on F . Therefore the above results can be applied to describe special values of certain automorphic representations of $GL_n(\mathbb{A}_F)$. \square

5. MOTIVIC INTERPRETATION

5.1. The Deligne conjecture. We firstly recall the statement of the general Deligne conjecture. For details, we refer the reader to Deligne's original paper [Del79]. We adapt the notation in [HL16].

Let \mathcal{M} be a motive over \mathbb{Q} with coefficients in a number field E , pure of weight w . For simplicity, we assume that if w is even then $(w/2, w/2)$ is not a Hodge type of \mathcal{M} . In this case, the motive is critical in the sense of [Del79]. Deligne has defined two elements $c^+(\mathcal{M})$ and $c^-(\mathcal{M}) \in (E \otimes \mathbb{C})^\times$ as determinants of certain period matrices.

For each $\rho \in J_E$, we may define the L -function $L(s, \mathcal{M}, \rho)$. We write $L(s, \mathcal{M}) = L(s, \mathcal{M}, \rho)_{\rho \in J_E}$. If $L(s, \mathcal{M}, \rho)$ is holomorphic at $s = s_0$ for all $\rho \in J_E$, we may consider $L(s_0, \mathcal{M})$ as an element in $E \otimes \mathbb{C}$ as before.

Definition 5.1.1. We say an integer m is **critical** for \mathcal{M} if neither $L_\infty(\mathcal{M}, s)$ nor $L_\infty(\check{\mathcal{M}}, 1 - s)$ has a pole at $s = m$ where $\check{\mathcal{M}}$ is the dual of \mathcal{M} . We call m a **critical value** of \mathcal{M} .

Deligne has formulated a conjecture on special values of motivic L -function as follows.

Conjecture 5.1.1. (the Deligne conjecture) *Let m be a critical point for \mathcal{M} . We write ϵ for the sign of $(-1)^m$. We then have:*

$$(5.1.1) \quad L(m, \mathcal{M}) \sim_E (2\pi i)^{mn^\epsilon} c^\epsilon(\mathcal{M})$$

where $n^\epsilon := \dim_E \mathcal{M}^\epsilon$.

The following lemma can be deduced easily from (1.3.1) of [Del79] (for the proof, see Lemma 3.1 of [LIN15b]).

Lemma 5.1.1. *Let \mathcal{M} be a pure motive of weight w as before. We assume that if w is even then $(w/2, w/2)$ is not a Hodge type of \mathcal{M} . Let $T(\mathcal{M}) := \{p \mid (p, w - p) \text{ is a Hodge type of } \mathcal{M}\}$. An integer m is critical for \mathcal{M} if and only if:*

$$\max\{p \in T(\mathcal{M}) \mid p < w/2\} < m \leq \min\{p \in T(\mathcal{M}) \mid p > w/2\}.$$

In particular, critical values always exists in this case.

Remark 5.1.1. We have assumed that if w is even then $(w/2, w/2)$ is not a Hodge type of \mathcal{M} . In this case, $\dim_E \mathcal{M}$ is even and $n^+ = n^- = \dim_E \mathcal{M}/2$.

It is not easy to relate Deligne's periods to geometric objects directly. In [Har13a] and its generalization in [LIN15c], more motivic periods are defined for motives over a CM field. These motivic periods can be related more easily to geometric objects. The Deligne periods are calculated in terms of these new periods in the above two papers and in [HL16].

5.2. Deligne conjecture for tensor product of motives. We give a special example of the results in [HL16] which fits in our main results.

Let M (resp. M') be a regular motive over F with coefficients in a number field E of rank n (resp. rank 1) and pure of weight w .

We first fix $\rho \in J_E$ an embedding of the coefficient field. For each $\tau \in J_F$, we write the Hodge type of M at τ (and ρ) as $(p_i(\tau), q_i(\tau))_{1 \leq i \leq n}$ with $p_1(\tau) > p_2(\tau) > \dots > p_n(\tau)$. We know that $q_i(\tau) = w - p_i(\tau)$.

We write the Hodge type of M' at τ (and ρ) as $(p(\tau), q(\tau))$. We assume that for any i and τ , $2p_i(\tau) + p(\tau) - q(\tau) \neq 0$.

Let $I(M, M')$ be the map on Φ which sends $\tau \in \Phi$ to $\#\{i : 2p_i(\tau) + p(\tau) - q(\tau) - w > 0\}$.

The motivic periods $Q^{(i)}(M, \tau)$, $0 \leq i \leq n$ and $Q^{(j)}(M, \tau)$, $0 \leq j \leq 1$ are defined in Definition 3.1 of [HL16] as elements in $(E \otimes \mathbb{C})^\times$.

As usual, we identify $E \otimes \mathbb{C}$ with \mathbb{C}^{J_E} . We write the ρ -component of $Q^{(i)}(M, \tau)$ by $Q^{(i)}(M, \tau)_\rho$. We define

$$Q^{*, I(M, M')}(M) := \left(\prod_{\tau \in \Phi} Q^{(I(M, M')(\tau))}(M, \tau)_\rho \right)_{\rho \in J_E}.$$

We remark that the index $I(M, M')$ depends implicitly on the embedding $\rho \in J_E$.

Similarly, we write $Q^{*, (0)}(M, \tau)^{n - I(M', M)(\tau)}$ for $(Q^{(0)}(M, \tau)_\rho)^{n - I(M', M)(\tau)}$ for $\rho \in J_E$.

Proposition 5.2.1. *The Deligne's periods for the motive $\text{Res}_{F/\mathbb{Q}}(M \otimes M')$ satisfy:*

$$c^+ \text{Res}_{F/\mathbb{Q}}(M \otimes M') \sim_E (2\pi i)^{-\frac{|\Phi|n(n-1)}{2}} \mathcal{J}_F^{[n/2]} (D_{F^+}^{1/2})^n \times \prod_{\tau \in \Sigma_\Phi} Q^{*, I(M, M')}(M) \prod_{\tau \in \Phi} Q^{*, (0)}(M, \tau)^{n - I(M', M)(\tau)} Q^{*, (1)}(M, \tau)^{I(M', M)(\tau)}.$$

Moreover, we have

$$c^- \text{Res}_{F/\mathbb{Q}}(M \otimes M') \sim_E e_\Phi^n c^- \text{Res}_{F/\mathbb{Q}}(M \otimes M').$$

Proof. The proposition follows from Propositions 2.11 and 3.13 of [HL16]. We refer to Definition 3.2 of *loc.cit* for the definition of the split index. It is enough to show that $sp(i, M; M', \tau) = 0$ if $i \neq I(M, M')(\tau)$, $sp(i, M; M', \tau) = 1$ if $i = I(M, M')(\tau)$, $sp(0, M'; M, \tau) = n - I(M, M')(\tau)$ and $sp(1, M'; M, \tau) = I(M, M')(\tau)$.

We fix $\tau \in J_F$. We denote $I(M, M')(\tau)$ by t . We have:

$$p_1(\tau) - \frac{p(\tau) + q(\tau) + w}{2} > \dots > p_t(\tau) - \frac{p(\tau) + q(\tau) + w}{2} > -p(\tau) > \\ p_{t+1} - \frac{p(\tau) + q(\tau) + w}{2} > \dots > p_n - \frac{p(\tau) + q(\tau) + w}{2}.$$

Therefore $sp(i, M; M', \tau) = 0$ for $i \neq t$ and $sp(t, M; M', \tau) = 0$ by the definition of split index. The proof for $sp(0, M'; M, \tau) = n - I(M, M')(\tau)$ and $sp(1, M'; M, \tau) = I(M, M')(\tau)$ is similar.

We now prove the second part. We use the notation $n_\tau(\rho)$ and $e_\tau(\rho) = (-1)^{n_\tau(\rho)}$ as in Remark 2.2 of [HL16]. It is easy to see that $n_{\bar{\tau}}(\rho) = n - n_\tau(\rho)$.

Let $e = \prod_{\tau \in \Phi} e_\tau$ be an element in $(E \otimes \mathbb{C})^\times$. Let $g \in \text{Aut}(\mathbb{C})$. Recall that $e_{g\rho}(\tau) = e_\rho(g^{-1}\tau)$ by Remark A.2 of [HL16]. Then

$$e(g\rho) = \prod_{\tau \in \Phi} e_\tau(g\rho) = \prod_{\tau \in \Phi} e_{g^{-1}\tau}(\rho) = (-1)^{n\#(g^{-1}\Phi) \setminus \Phi} = (-1)^{n*\#(\Phi \setminus g\Phi)} e(\rho).$$

Hence $e = \pm e_\Phi^n$ by the definition of e_Φ .

We know $c^-(\text{Res}_{F/\mathbb{Q}}(M \otimes M')) = ec^+(\text{Res}_{F/\mathbb{Q}}(M \otimes M'))$ by Remark 2.2 of [HL16]. □

5.3. Compatibility of the main results with the Deligne conjecture.

Let Π be as in section 4.3. It is conjectured that the representation Π is attached to a motive $M = M(\Pi)$ over F with coefficients in $E(\Pi)$ (c.f. Conjecture 4.5 and paragraph 4.3.3 of [Clo90]).

We fix $\rho \in J_E$. We write the infinity type of Π at $\tau \in \Phi$ as $z^{a_i(\tau)} \bar{z}^{-a_i(\tau)}$. Then the Hodge type of $M(\Pi)$ at τ should be $(-a_i(\tau) + \frac{n-1}{2}, a_i(\tau) + \frac{n-1}{2})_{1 \leq i \leq n}$.

Similarly, we write $M' = M(\eta)$ the conjectural motif associated to η .

We have:

$$(5.3.1) \quad L(s, M \otimes M') = L(s + \frac{1-n}{2}, \Pi \times \eta).$$

We want to compare Theorem 4.3.1 with the Deligne conjecture. The main difficulty is to compare the automorphic periods with the motivic periods. Recall that the automorphic periods $P^{(I)}(\Pi)$ are constructed from different geometric objects. It is hard to relate them with the same motive $M(\Pi)$. However, if we admit the Tate conjecture, we will get

$$(5.3.2) \quad P^{(I)}(\Pi) \sim_{E(\Pi)} Q^{(I)}(\Pi)$$

as in section 4.4 of [HL16]. Roughly speaking, the Tate conjecture says that a motive is determined by its l -adic realizations.

Corollary 5.3.1. *We keep the notations as in Theorem 4.3.1. If we admit the Tate conjecture, then the Deligne conjecture is true up to $\sim_{E(\Pi, \eta); F^{gal}}$ for critical values $m > n + \omega(\eta)/2$ of the conjectural motive $M(\Pi) \otimes M(\eta)$.*

Proof. We compare Proposition 5.2.1, Theorem 4.3.1 equation (5.3.1), equation (5.3.2) and the fact that:

$$Q^{(0)}(M(\eta), \tau) \sim_{E(\tau)} p(\check{\eta}^c, \tau), \text{ and } Q^{(1)}(M(\eta), \tau) \sim_{E(\tau)} p(\check{\eta}, \tau)$$

by Lemma 3.17 of [HL16]. It is easy to verify that $I(\Pi, \eta) = I(M(\Pi), M(\eta))$. It remains to show that if $m > n + \omega(\eta)/2$ critical for $M(\Pi) \otimes M(\eta)$ then

$$m - \frac{n-1}{2} \geq \frac{n + \omega(\eta)}{2} \text{ satisfies equation (3.2.1).}$$

to be fixed after generalization of α □

Extra remarks: 1. In the end of section 1, change α to other notation
2. add a remark on $m > n$ to $m > n/2$ and compare with Deligne conjecture

REFERENCES

- [Art03] J. Arthur, *An introduction to the trace formula*, Harmonic analysis, the trace formula and Shimura varieties (J. Arthur, D. Ellwood, and R. Kottwitz, eds.), Clay Mathematics Proceedings, vol. 4, American Mathematical Society Clay Mathematics Institute, 2003, pp. 1–264. [22](#)
- [BHR94] Don Blasius, Michael Harris, and Dinakar Ramakrishnan, *Coherent cohomology, limits of discrete series, and Galois conjugation*, Duke Math. J. **73** (1994), no. 3, 647–685. MR 1262930 [8](#)
- [Bla86] Don Blasius, *On the critical values of Hecke L -series*, Ann. of Math. (2) **124** (1986), no. 1, 23–63. [21](#)
- [Car12] A. Caraiani, *Local–global compatibility and the action of monodromy on nearby cycles*, Duke Math. J (2012). [23](#)
- [Clo90] L. Clozel, *Motifs et formes automorphes: applications du principe de functorialité*, Automorphic forms, Shimura varieties, and L -functions, vol. 1, Perspectives in Mathematics, no. 10, Boston, MA: Academic Press, 1990, pp. 77–159. [27](#)
- [Clo91] ———, *Représentations galoisiennes associées aux représentations automorphes autoduales de $Gl(n)$* , Publications mathématiques de l’I.H.E.S **73** (1991), 97–145. [21](#), [22](#)
- [Clo93] ———, *On the cohomology of kottwitz’s arithmetic varieties*, Duke Math. J. (1993), 757–795. [8](#), [23](#)
- [Clo12] ———, *Purity reigns supreme*, International Math. Research Notices (2012). [23](#)
- [Del79] P. Deligne, *Valeurs de fonctions L et périodes d’intégrales*, Automorphic forms, representations and L -functions (A. Borel and W. Casselman, eds.), Proceedings of the symposium in pure mathematics of the American mathematical society, vol. 33, American Mathematical Society, 1979. [25](#)
- [Gar08] Paul Garrett, *Values of Archimedean zeta integrals for unitary groups*, Eisenstein series and applications, Progr. Math., vol. 258, Birkhäuser Boston, Boston, MA, 2008, pp. 125–148. [2](#), [18](#), [21](#)
- [GH16] Harald Grobner and Michael Harris, *Whittaker periods, motivic periods, and special values of tensor product L -functions*, Journal of the Institute of Mathematics of Jussieu **FirstView** (2016), 1–59. [8](#)
- [GR14] H. Grobner and A. Raghuram, *On some arithmetic properties of automorphic forms of $gl(m)$ over a division algebra*, Int. J. Number Theory (2014), no. 10, 963–1013. [7](#)
- [Gue16] Lucio Guerberoff, *Period relations for automorphic forms on unitary groups and critical values of L -functions*, Doc. Math. **21** (2016), 1397–1458. [2](#), [5](#), [10](#), [12](#), [13](#), [15](#), [16](#), [17](#), [18](#), [20](#)
- [Har93] Michael Harris, *L -functions of 2×2 unitary groups and factorization of periods of Hilbert modular forms*, J. Amer. Math. Soc. **6** (1993), no. 3, 637–719. [12](#), [15](#)
- [Har97] ———, *L -functions and periods of polarized regular motives*, J. Reine Angew. Math. **483** (1997), 75–161. [2](#), [4](#), [9](#), [14](#), [15](#), [16](#)
- [Har07] ———, *Cohomological automorphic forms on unitary groups. II. Period relations and values of L -functions*, Harmonic analysis, group representations, automorphic forms and invariant theory, Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., vol. 12, World Sci. Publ., Hackensack, NJ, 2007, pp. 89–149. [15](#)
- [Har08] ———, *A simple proof of rationality of Siegel–Weil Eisenstein series*, Eisenstein series and applications, Progr. Math., vol. 258, Birkhäuser Boston, Boston, MA, 2008, pp. 149–185. [15](#)
- [Har13a] M. Harris, *L -functions and periods of adjoint motives*, Algebra and Number Theory (2013), no. 7, 117–155. [25](#)

- [Har13b] Michael Harris, *Beilinson-Bernstein localization over \mathbb{Q} and periods of automorphic forms*, Int. Math. Res. Not. IMRN (2013), no. 9, 2000–2053. MR 3053412 [8](#)
- [HK91] Michael Harris and Stephen S. Kudla, *The central critical value of a triple product L -function*, Ann. of Math. (2) **133** (1991), no. 3, 605–672. [12](#)
- [HL16] M. Harris and J. LIN, *Periods relations and special values of Rankin-Selberg L -functions*, To appear in the proceeding for the conference "Representation theory, number theory, and invariant theory" in honor of Roger Howe., 2016. [25](#), [26](#), [27](#)
- [KMSW14] Tasho Kaletha, Alberto Mínguez, Sug Woo Shin, and Paul-James White, *Endoscopic classification of representations: inner forms of unitary groups*, preprint. [10](#), [22](#)
- [Lab11] Jean-Pierre Labesse, *Changement de base CM et séries discrètes*, On the stabilization of the trace formula, Stab. Trace Formula Shimura Var. Arith. Appl., vol. 1, Int. Press, Somerville, MA, 2011, pp. 429–470. [8](#), [23](#)
- [Li92] Jian-Shu Li, *Nonvanishing theorems for the cohomology of certain arithmetic quotients*, J. Reine Angew. Math. **428** (1992), 177–217. [17](#)
- [LIN15a] J. LIN, *Period relations for automorphic induction and applications, I*, Comptes Rendus Mathématiques **353** (2015). [21](#)
- [LIN15b] ———, *Period relations for automorphic induction and applications, I*, arXiv:1511.03517, 2015. [25](#)
- [LIN15c] ———, *Special values of automorphic L -functions for $gl_n \times gl_{n'}$ over CM fields, factorization and functoriality of arithmetic automorphic periods*, Ph.D. thesis, Université Paris 7, 2015. [21](#), [24](#), [25](#)
- [LS86] J. P. Labesse and J. Schwermer, *On liftings and cusp cohomology of arithmetic groups*, Inventiones mathematicae (1986), no. 83, 383–401. [21](#)
- [LS04] Jian-Shu Li and Joachim Schwermer, *On the Eisenstein cohomology of arithmetic groups*, Duke Math. J. **123** (2004), no. 1, 141–169. [8](#), [23](#)
- [Mil90] James S. Milne, *Canonical models of (mixed) Shimura varieties and automorphic vector bundles*, Automorphic forms, Shimura varieties, and L -functions, Vol. I, Perspect. Math., vol. 10, Academic Press, Boston, MA, 1990, pp. 283–414. [5](#), [11](#)
- [Min11] A. Mínguez, *Unramified representations of unitary groups*, On the stabilization of the trace formula (L. Clozel, M. Harris, J. P. Labesse, and B. C. Ngô, eds.), vol. 1, International Press, 2011. [5](#), [7](#), [22](#)
- [MS10] James S. Milne and Junecue Suh, *Nonhomeomorphic conjugates of connected Shimura varieties*, Amer. J. Math. **132** (2010), no. 3, 731–750. MR 2666906 [5](#)
- [Sch94] Joachim Schwermer, *Eisenstein series and cohomology of arithmetic groups: the generic case*, Invent. Math. **116** (1994), no. 1-3, 481–511. [8](#)
- [Wal84] N. Wallach, *On the constant term of a square integrable automorphic form*, Operator algebras and group representations, volume II, Monographs Stud. Math., vol. 18, Pitman, Boston, MA, London, 1984, pp. 227–237. [8](#), [23](#)
- [Yos94] Hiroyuki Yoshida, *On the zeta functions of Shimura varieties and periods of Hilbert modular forms*, Duke Math. J. **75** (1994), no. 1, 121–191. [9](#), [18](#), [20](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, 25 GORDON STREET,
LONDON WC1H 0AY, UK

E-mail address: l.guerberoff@ucl.ac.uk

INSTITUTE DES HAUTES ÉTUDES SCIENTIFIQUES, 91440 BURES-SUR-YVETTE, FRANCE

E-mail address: linjie@ihes.fr