

**Estimates from below for the spectral function and for
the remainder in local Weyl's law**

Dmitry JAKOBSON and Iosif POLTEROVICH



Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Mai 2005

IHES/M/05/16

ESTIMATES FROM BELOW FOR THE SPECTRAL FUNCTION AND FOR THE REMAINDER IN LOCAL WEYL'S LAW

DMITRY JAKOBSON AND IOSIF POLTEROVICH

ABSTRACT. We obtain asymptotic lower bounds for the spectral function of the Laplacian and for the remainder in local Weyl's law on manifolds. In the negatively curved case, thermodynamic formalism is applied to improve the estimates. Key ingredients of the proof include the wave equation parametrix, a pretrace formula and the Dirichlet box principle.

1. INTRODUCTION AND MAIN RESULTS

1.1. Spectral function and Weyl's law. Let X be a compact Riemannian manifold of dimension $n \geq 2$ with the metric $\{g_{ij}\}$ and the volume V . Let Δ be the Laplacian on X with the eigenvalues $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$ and the corresponding orthonormal basis $\{\phi_i\}$ of eigenfunctions: $\Delta\phi_i = \lambda_i\phi_i$.

Given $x, y \in X$, let

$$N_{x,y}(\lambda) = \sum_{\sqrt{\lambda_i} \leq \lambda} \phi_i(x)\phi_i(y)$$

be the spectral function of the Laplacian. On the diagonal $x = y$ we denote it simply $N_x(\lambda)$. If $N(\lambda) = \#\{\sqrt{\lambda_i} \leq \lambda\}$ is the eigenvalue counting function, then $N(\lambda) = \int_X N_x(\lambda) dV$. Let

$$(1.1.1) \quad \sigma_n = \frac{2\pi^{n/2}}{n\Gamma(n/2)}$$

be the volume of the unit ball in \mathbf{R}^n . The asymptotic behavior of the spectral and the counting functions is given by ([Hor], see also [Shu]):

$$(1.1.2) \quad \begin{aligned} N_{x,y}(\lambda) &= O(\lambda^{n-1}), & x \neq y; \\ N_x(\lambda) &= \frac{\sigma_n}{(2\pi)^n} \lambda^n + R_x(\lambda), & R_x(\lambda) = O(\lambda^{n-1}); \\ N(\lambda) &= \frac{V\sigma_n}{(2\pi)^n} \lambda^n + R(\lambda), & R(\lambda) = O(\lambda^{n-1}). \end{aligned}$$

We refer to the asymptotics of $N_x(\lambda)$ as *local Weyl's law*, the asymptotics for $N(\lambda)$ being the usual Weyl's law for the distribution of eigenvalues.

In the present paper we focus on asymptotic *lower* bounds for the spectral function and for the remainder in local Weyl's law. We recall that $f_1(\lambda) = \Omega(f_2(\lambda))$ for a function f_1 and a positive function f_2 means $\limsup_{\lambda \rightarrow \infty} |f_1(\lambda)|/f_2(\lambda) > 0$.

1991 *Mathematics Subject Classification.* 35P20, 37C30, 58J50, 81Q50.

Key words and phrases. Weyl's law, spectral function, wave kernel, negative curvature, Anosov flow, thermodynamic formalism.

The first author is supported by NSERC, FQRNT, Alfred P. Sloan Foundation fellowship and Dawson fellowship. The second author is supported by NSERC and FQRNT.

Theorem 1.1.3. *Let X be a compact n -dimensional Riemannian manifold, and let $x, y \in X$ be two points that are not conjugate along any shortest geodesic joining them. Then*

$$(1.1.4) \quad N_{x,y}(\lambda) = \Omega\left(\lambda^{\frac{n-1}{2}}\right).$$

In order to formulate the on-diagonal counterpart of Theorem 1.1.3 we introduce the following notation. Let $\kappa_x = \min\{j \geq 1 \mid u_j(x, x) \neq 0\}$, where $u_j(x, x)$ is the j -th local heat invariant. If $u_j(x, x) = 0$ for all $j \geq 1$ we set $\kappa_x = \infty$. We recall that $u_1(x, x) = \frac{\tau(x)}{6}$, where $\tau(x)$ is the scalar curvature of X at the point x .

Theorem 1.1.5. *Let X be an n -dimensional Riemannian manifold and $x \in X$ be an arbitrary point. If $n - 2\kappa_x - 1 > 0$ then*

$$(1.1.6) \quad R_x(\lambda) = \Omega(\lambda^{n-2\kappa_x-1}).$$

If $n - 4\kappa_x - 1 < 0$ and, in addition, X is a manifold without conjugate points, then

$$(1.1.7) \quad R_x(\lambda) = \Omega(\lambda^{\frac{n-1}{2}})$$

Theorems 1.1.3 and 1.1.5 are proved by analyzing the asymptotics of the parametrix for the wave equation. Estimate (1.1.7) should be compared with the celebrated Hardy-Landau lower bound for the remainder in the Gauss circle problem or, equivalently, for the remainder in Weyl's law on a 2-dimensional flat square torus (see [Sou] for the most recent improvement of such a bound). Interestingly, we get the same exponent as in the Hardy-Landau bound for any surface without conjugate points.

1.2. Thermodynamic formalism. Asymptotic lower bounds (1.1.4) (1.1.6) and (1.1.7) can be improved for manifolds of negative curvature, see section 1.3. We assume that for any pair of directions ξ, η the sectional curvature $K(\xi, \eta)$ satisfies

$$(1.2.1) \quad -K_1^2 \leq K(\xi, \eta) \leq -K_2^2.$$

Apart from the usual wave equation techniques (cf. [D-G], [Ber], [K]) our method uses the thermodynamic formalism (see, for example, [Bow], [P-P]). Let G^t be the geodesic flow on the unit tangent bundle SX and let E_ξ^u be the unstable subspace for G^t , $\xi \in SX$. The *Sinai-Ruelle-Bowen potential* is a Hölder continuous function $\mathcal{H} : SX \rightarrow \mathbf{R}$ which for any $\xi \in SX$ is defined by the formula (see [B-R], [Sin2])

$$(1.2.2) \quad \mathcal{H}(\xi) = \frac{d}{dt} \Big|_{t=0} \ln \det dG^t|_{E_\xi^u},$$

For any continuous function $f : SX \rightarrow \mathbf{R}$ one can define the *topological pressure*

$$(1.2.3) \quad P(f) = \sup_{\mu} \left(h_{\mu} + \int f d\mu \right),$$

where the supremum is taken over all G^t -invariant measures μ and h_{μ} denotes the *measure-theoretical entropy* of the geodesic flow (see [Bow]). In particular $P(0) = h$, where h is the *topological entropy* of the flow. It is well-known that for the Sinai-Ruelle-Bowen potential $P(-\mathcal{H}) = 0$ and the corresponding equilibrium measure (i.e. the measure on which the supremum is attained) is the Liouville measure μ_L on the unit tangent bundle:

$$(1.2.4) \quad h_{\mu_L} = \int_{SX} \mathcal{H} d\mu_L$$

1.3. Estimates for negatively curved manifolds. Slightly modifying the usual Ω notation, we write $f_1(\lambda) = \Omega_\infty(f_2(\lambda))$ for any $f_1(\lambda)$ and $f_2(\lambda) > 0$ if

$$\limsup_{\lambda \rightarrow \infty} \frac{|f_1(\lambda)|}{f_2(\lambda)} = \infty.$$

Theorem 1.3.1. *The remainder in the local Weyl's law on an n -dimensional compact negatively curved manifold satisfies:*

$$(1.3.2) \quad R_x(\lambda) = \begin{cases} \Omega_\infty \left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta} \right) & \forall \delta > 0, \quad n \leq 5; \\ \Omega(\lambda^{n-3}), & n \geq 6. \end{cases}$$

For $n = 2$, a similar statement is proved in [K], see a discussion below.

Theorem 1.3.3. *On a compact n -dimensional negatively curved manifold the spectral function $N_{x,y}(\lambda)$ satisfies for any $\delta > 0$ and $x \neq y$:*

$$(1.3.4) \quad N_{x,y}(\lambda) = \Omega_\infty \left(\lambda^{\frac{n-1}{2}} (\log \lambda)^{\frac{P(-\mathcal{H}/2)}{h} - \delta} \right).$$

Theorems 1.3.1 and 1.3.3 are proved in section 5.

Let us estimate the power $\frac{P(-\mathcal{H}/2)}{h}$ in terms of curvatures. It was shown in [Sin1] that on an n -dimensional negatively curved manifold satisfying (1.2.1),

$$(n-1)K_2 \leq h_{\mu_L} \leq h \leq (n-1)K_1$$

we can obtain (using the definition of pressure and (1.2.4)):

$$(1.3.5) \quad P \left(-\frac{\mathcal{H}}{2} \right) \geq h_{\mu_L} - \int \mathcal{H} \mu_L + \frac{1}{2} \int \mathcal{H} d\mu_L = \frac{h_{\mu_L}}{2} \geq \frac{(n-1)K_2}{2}$$

and therefore

$$\frac{P(-\mathcal{H}/2)}{h} \geq \frac{K_2}{2K_1} > 0.$$

1.4. Discussion. Remainder estimates in (1.1.2) for $N(\lambda)$ and $N_x(\lambda)$ are attained for round spheres and hence are sharp. One can replace $O(\lambda^{n-1})$ by $o(\lambda^{n-1})$ in the bounds for $R(\lambda)$ (respectively, for $R_x(\lambda)$) if the initial directions of closed geodesics (respectively, of geodesic loops at x) form a set of measure zero (for $R(\lambda)$ see [D-G], for $R_x(\lambda)$ see [Saf], [S-V, section 1.8], and also [S-Z]).

Both local and integrated remainder estimates for Weyl's law were actively studied under various geometric conditions (see [CdV], [Iv], [Vol], [P-T], etc.). Of particular interest are manifolds of negative curvature. In this case, $R_x(\lambda) = O(\lambda^{n-1}/\log \lambda)$ ([Ber]). It is conjectured in [Ran] that on a surface of constant negative curvature $R(\lambda) = O(\lambda^{\frac{1}{2} + \epsilon})$ for any $\epsilon > 0$. In the same article it is proved that an average over λ of $N_{x,y}(\lambda)$ for $x \neq y$ satisfies such an upper bound. On surfaces corresponding to quaternionic lattices $R(\lambda) = \Omega \left(\frac{\sqrt{\lambda}}{\log \lambda} \right)$ (see [Hej]). Hence, if there are no additional assumptions, the bound conjectured in [Ran] is almost the best possible. However, it is believed that on a *generic* negatively curved surface $R(\lambda) = O(\lambda^\epsilon)$ for any $\epsilon > 0$.

The present paper originated as an attempt to improve the estimate $R_x(\lambda) = \Omega(\sqrt{\lambda})$ obtained in an unpublished Princeton Ph.D. thesis [K] for surfaces of arbitrary negative curvature. The main novelty of our approach compared to [K] is the use of thermodynamic formalism. It is also noticed that the restriction to the negatively curved case considered in [K] can be replaced by much more general

assumptions, see Theorems 1.1.3 and 1.1.5. We estimate $N_{x,y}(\lambda)$ not just on the diagonal, but also for $x \neq y$. Our results apply to arbitrary dimensions, while [K] deals only with surfaces. Working in higher dimensions makes the analysis of the parametrix more difficult. In particular, asymptotics have different nature in high and low dimensions, see Theorem 1.3.1.

On surfaces of constant negative curvature (1.3.2) coincides with the bound in [K], but the techniques of thermodynamic formalism for hyperbolic flows allow us to improve the results obtained in [K] for manifolds of variable negative curvature. We do not need the hypothesis $K_1/K_2 < 2$ of [K] to get a logarithmic improvement in the estimates and, moreover, we get higher powers of the logarithm. Note that the results of [Ran] cited above make it unlikely that in two dimensions the power of λ in a lower bound for $N_x(\lambda)$ can be made greater than $1/2$.

We would like to emphasize that the powerful techniques of hyperbolic dynamics can be quite useful for proving spectral estimates on negatively curved manifolds (see also [Vol] where results of Bowen were applied to prove upper bounds for the remainder in Weyl's law).

2. WAVE KERNEL, SPECTRAL FUNCTION AND THE PRETRACE FORMULA

2.1. Smoothed Fourier transform of the wave kernel. The even part of the wave kernel on X satisfies

$$(2.1.1) \quad e(t, x, y) = \sum_{i=0}^{\infty} \cos(\sqrt{\lambda_i}t) \phi_i(x) \phi_i(y)$$

It is the fundamental solution of the wave equation

$$(2.1.2) \quad \begin{aligned} (\partial^2/\partial t^2 - \Delta)e(t, x, y) &= 0, \\ e(0, x, y) &= \delta(x - y), \\ (\partial/\partial t)e(0, x, y) &= 0. \end{aligned}$$

Take a smooth function $\psi \in C_0^\infty(\mathbf{R})$ such that $\text{supp } \psi \subseteq [-1, 1]$, it is even and monotone decreasing on $[0, 1]$, $\psi \geq 0$, $\psi(0) = 1$. Fix two large positive parameters λ, T and choose a compactly supported smooth test function

$$(2.1.3) \quad (1/T)\psi(t/T) \cos(\lambda t).$$

Given $x, y \in M$, denote the integral in the right-hand side by

$$(2.1.4) \quad k_{\lambda, T}(x, y) = \int_{-\infty}^{\infty} \frac{\psi(t/T)}{T} \cos(\lambda t) e(t, x, y) dt$$

Substituting (2.1.1) into (2.1.4) we obtain

$$(2.1.5) \quad k_{\lambda, T}(x, y) = \sum_{i=0}^{\infty} \phi_i(x) \phi_i(y) H_{\lambda, T}(\sqrt{\lambda_i}),$$

where

$$(2.1.6) \quad H_{\lambda, T}(r) = \int_{-\infty}^{\infty} \frac{\psi(t/T)}{T} \cos(\lambda t) \cos(rt) dt = \frac{\widehat{\psi}(T(r + \lambda)) + \widehat{\psi}(T(r - \lambda))}{2},$$

where

$$\widehat{\psi}(s) = \int_{-\infty}^{\infty} e^{is\zeta} \psi(\zeta) d\zeta$$

is the Fourier transform of ψ . Replacing the sum in (2.1.5) by an integral, we get the following representation of $k_{\lambda,T}(x, y)$:

$$(2.1.7) \quad \int_0^\infty H_{\lambda,T}(r) dN_{x,y}(r) = \int_0^\infty \frac{\widehat{\psi}(T(r+\lambda)) + \widehat{\psi}(T(r-\lambda))}{2} dN_{x,y}(r) = k_{\lambda,T}(x, y)$$

Formula (2.1.7) plays a key role in our analysis.

On the diagonal, we shall use the following notation:

$$(2.1.8) \quad \int_0^\infty H_{\lambda,T}(r) dR_x(r) = \widetilde{k}_{\lambda,T}(x)$$

Note that the contribution of the main term in Weyl's law has been subtracted.

2.2. Relation between $N_{x,y}(\lambda)$ and $k_{\lambda,T}(x, y)$, $R_x(\lambda)$ and $\widetilde{k}_{\lambda,T}(x)$. Assume that T is either fixed or $T(\lambda) \rightarrow \infty$ as $\lambda \rightarrow \infty$. The following lemma will be used to prove Theorems 1.1.3 and 1.1.5.

Lemma 2.2.1. (i) Let $N_{x,y}(\lambda) = o(\lambda^a)$, $a > 0$. Then $k_{\lambda,T}(x, y) = o(\lambda^a)$. (ii) Let $R_x(\lambda) = o(\lambda^a)$. Then $\widetilde{k}_{\lambda,T}(x) = o(\lambda^a)$.

Proof. We shall prove (i), the proof of part (ii) is analogous. Denote $\widehat{\psi}'$ by Ψ . Since ψ has compact support, Ψ is Schwartz class. By the assumption of the lemma, for any $\epsilon > 0$, $N_{x,y}(\lambda) < \epsilon\lambda^a$ for large enough λ . Consider the left hand side of (2.1.7):

$$(2.2.2) \quad \int_0^\infty H_{\lambda,T}(r) dN_{x,y}(r).$$

Taking into account (2.1.7) and integrating (2.2.2) by parts we obtain

$$(2.2.3) \quad k_{\lambda,T}(x, y) \leq \frac{\epsilon T}{2} \int_0^\infty |\Psi(T(r-\lambda))| r^a dr + \frac{\epsilon T}{2} \int_0^\infty |\Psi(T(r+\lambda))| r^a dr.$$

Since Ψ is Schwartz class the second term of (2.2.3) is $O(1)$. Changing variables in the first term of (2.2.3), we obtain

$$\frac{\epsilon T}{2} \int_{-\lambda}^\infty |\Psi(Ts)|(s+\lambda)^a ds = \frac{\epsilon \lambda^a}{2} \int_{-\lambda/T}^\infty |\Psi(u)| \left(1 + \frac{u}{\lambda T}\right)^a du \leq C\epsilon \lambda^a$$

for some constant $C > 0$, where the last inequality again follows from the fact that Ψ is Schwartz class. Since ϵ can be chosen arbitrarily small, we get $k_{\lambda,T}(x, y) = o(\lambda^a)$, and this completes the proof of the lemma. \square

A modification of this lemma is used to prove Theorems 1.3.1 and 1.3.3.

Lemma 2.2.4. (i) Let $N_{x,y}(\lambda) = O(\lambda^a (\log \lambda)^b)$, $a, b > 0$. Then $k_{\lambda,T}(x, y) = O(\lambda^a (\log \lambda)^b)$. (ii) Let $R_x(\lambda) = O(\lambda^a (\log \lambda)^b)$, $a, b > 0$. Then $\widetilde{k}_{\lambda,T}(x) = O(\lambda^a (\log \lambda)^b)$.

Proof. This lemma is proved similarly to the previous one. Again, we shall prove (i), the proof of (ii) is analogous.

By the assumption of the lemma there exists a constant $C > 0$ such that $N_{x,y}(\lambda) < C\lambda^a (\log(1+\lambda))^b$. Writing a representation of $k_{\lambda,T}(x, y)$ analogous to

(2.2.3) we get that the second term is $O(1)$ as before, and the first term can be rewritten as

$$\begin{aligned} & \frac{CT}{2} \int_{-\lambda}^{\infty} |\Psi(Ts)|(s+\lambda)^a (\log(1+\lambda+s))^b ds = \\ & \frac{C\lambda^a (\log \lambda)^b}{2} \int_{-\lambda T}^{\infty} |\Psi(u)| \left(1 + \frac{u}{\lambda T}\right)^a \left(1 + \frac{\log(1 + \frac{1+u/T}{\lambda})}{\log \lambda}\right)^b du = O(\lambda^a (\log \lambda)^b). \end{aligned}$$

Remark 2.2.5. Statements similar to Lemmas 2.2.1 and 2.2.4 are proved in [K].

2.3. The pretrace formula. If X is of negative curvature (or, more, generally, has no conjugate points) it is useful to consider the fundamental solution $E(t, x, y)$ of the wave equation (2.1.2) on the universal cover M of X . Then given $x, y \in X$, we have

$$(2.3.1) \quad e(t, x, y) = \sum_{\omega \in \Gamma} E(t, x, \omega y),$$

where the sum is taken over $\Gamma = \pi_1(X)$. Let $K_{\lambda, T}(x, y)$ be the analogue of $k_{\lambda, T}(x, y)$ corresponding to the wave kernel $E(t, x, y)$ on M :

$$(2.3.2) \quad K_{\lambda, T}(x, y) = \int_{-\infty}^{\infty} \frac{\psi(t/T)}{T} \cos(\lambda t) E(t, x, y) dt$$

Then (2.1.7) becomes

$$(2.3.3) \quad k_{\lambda, T}(x, y) = \int_0^{\infty} H_{\lambda, T}(r) dN_{x, y}(r) = \sum_{\omega \in \Gamma} K_{\lambda, T}(x, \omega y).$$

3. ASYMPTOTICS OF THE SMOOTHED FOURIER TRANSFORM

3.1. Parametrix. In this section we review the Hadamard parametrix construction for the wave equation (cf. [Ber, Prop. 27], see also [Zel]). We present it in the setting relevant for the proofs of Theorems 1.3.1 and 1.3.3: X is a negatively curved manifold, hence the parametrix is globally defined on M . In the proofs of Theorems 1.1.3 and 1.1.5 we make the necessary adjustments in order to use this construction.

We shall work in Riemannian normal coordinates centered at $x \in M$. Given $x, y \in M$ let $r = d(x, y)$ and let $E(t, x, y)$ be the fundamental solution of (2.1.2) M appearing in (2.3.1). The parametrix for $E(t, x, y)$ is given by:

$$(3.1.1) \quad E(t, x, y) = \frac{1}{\pi^{\frac{n-1}{2}}} |t| \sum_{j=0}^{\infty} u_j(x, y) \frac{(r^2 - t^2)_+^{j - \frac{n-3}{2} - 2}}{4^j \Gamma(j - \frac{n-3}{2} - 1)} \pmod{C^\infty},$$

The expression (3.1.1) is understood in the sense of generalized functions [G-S]. The coefficients $u_j(x, y)$ are the solutions of the transport equations ([Ber]) along the geodesic joining x and y (since X is negatively curved, such a geodesic on M is unique). In particular $u_0(x, y) = g^{-1/2}(y)$, where $g = \sqrt{\det g_{ij}}$. The on-diagonal values $u_j(x, x)$ are known as the *local heat invariants*, since they are the coefficients in the short time asymptotic expansion for the heat kernel.

We substitute (3.1.1) into (2.1.4) to get

$$(3.1.2) \quad K_{\lambda,T}(x, y) = \int_{-\infty}^{\infty} \frac{\psi(t/T) \cos(\lambda t) |t|}{T \pi^{\frac{n-1}{2}}} \sum_{j=0}^J u_j(x, y) \frac{(r^2 - t^2)_-^{j - \frac{n-3}{2} - 2}}{4^j \Gamma(j - \frac{n-3}{2} - 1)} dt \\ + E_{\lambda,T}(x, y)$$

where $J \geq 2([n/2] + 1) + 1$ and $E_{\lambda,T}(x, y) = O(1) \exp(O(T))$. ([Ber, p.266]).

3.2. Leading term cancellation in the pretrace formula. The results of sections 3.2–3.5 are essentially not new and could be deduced from [Ber]. We refer to [D-G, p. 46] and [Ber, p. 259] for the proper regularization of $E(t, x, x)$ at $t = 0$.

Lemma 3.2.1. *The following relation holds:*

$$(3.2.2) \quad \int_0^{\infty} H_{\lambda,T}(r) dR_x(r) = \int_{-\infty}^{\infty} \frac{\psi(t/T) \cos(\lambda t)}{T \pi^{\frac{n-1}{2}}} \sum_{j=1}^J \frac{u_j(x, x) |t|^{2j-n}}{4^j \Gamma(j - \frac{n-3}{2} - 1)} dt + \\ \sum_{g \in \Gamma \setminus \{\text{Id}\}} \int_{-\infty}^{\infty} \frac{\psi(t/T) \cos(\lambda t) |t|}{T \pi^{\frac{n-1}{2}}} \sum_{j=0}^J \frac{u_j(x, \gamma x) (r^2 - t^2)_-^{j - \frac{n-3}{2} - 2}}{4^j \Gamma(j - \frac{n-3}{2} - 1)} dt + O(1) \exp(O(T))$$

In other words, the contribution of the 0-th term of the parametrix (3.1.1) on the diagonal into the formula (3.1.2) cancels out the contribution of the leading term in Weyl's law on the left-hand side of (2.3.3).

Proof. Substituting (1.1.1) into (1.1.2) we get:

$$dN_x(r) = \frac{r^{n-1}}{2^{n-1} \pi^{n/2} \Gamma(n/2)} + dR_x(r),$$

and hence the contribution of the leading term in Weyl's law into the left-hand side of (2.3.3) is:

$$(3.2.3) \quad \int_{-\infty}^{\infty} \int_0^{\infty} \frac{\psi(t/T) r^{n-1} \cos(rt) dr dt}{2^{n-1} \pi^{n/2} \Gamma(n/2) T}$$

The contribution of the 0-th term of the parametrix on the diagonal to (3.1.2) is

$$(3.2.4) \quad \int_{-\infty}^{\infty} \frac{\psi(t/T) \cos(\lambda t) |t|^{-n}}{\pi^{\frac{n-1}{2}} \Gamma(\frac{1-n}{2}) T},$$

where $\frac{|t|^{-n}}{\Gamma(\frac{1-n}{2})}$ is understood as a generalized function in the sense of ([G-S]). If $n = 2m$ is even, $|t|^{-2m} = t^{-2m}$ in the sense of generalized functions and (see [Zel], [G-S]):

$$(3.2.5) \quad t^{-2m} = \text{Re}(t + i0)^{-2m} = \frac{(-1)^m}{(2m-1)!} \int_0^{\infty} r^{2m-1} \cos(rt) dr.$$

Substituting this into (3.2.4) and taking into account that

$$\Gamma\left(\frac{1}{2} - m\right) = \frac{(-1)^m m! 4^m \sqrt{\pi}}{(2m)!}$$

one obtains equality between (3.2.4) and (3.2.3). If $n = 2m + 1$ is odd, we have ([G-S]):

$$(3.2.6) \quad \frac{|t|^{-n}}{\Gamma(\frac{1-n}{2})} = \frac{(-1)^m \delta^{(2m)}(t) m!}{(2m)!} = \frac{m!}{\pi (2m)!} \int_0^{\infty} r^{2m} \cos(rt) dr,$$

where the second equality follows by taking the inverse Fourier transform of $\delta^{(2m)}(t)$. Substituting (3.2.6) into (3.2.4) we again get equality between (3.2.4) and (3.2.3). Formula (3.2.2) is then a consequence of (3.1.2) and (2.3.3). This completes the proof of the lemma. \square

Lemma 3.2.1 implies that

$$(3.2.7) \quad \tilde{k}_{\lambda,T}(x) = \sum_{\omega \in \Gamma \setminus \text{Id}} K_{\lambda,T}(x, \gamma x) + \tilde{K}_{\lambda,T}(x),$$

where

$$(3.2.8) \quad \tilde{K}_{\lambda,T}(x) = K_{\lambda,T}(x, x) - \int_{-\infty}^{\infty} \frac{\psi(t/T) \cos(\lambda t) |t|^{-n}}{\pi^{\frac{n-1}{2}} \Gamma(\frac{1-n}{2}) T} dt.$$

Formula (3.2.7) indicates that in order to estimate the remainder $R_x(\lambda)$ one has to study the leading terms in the right hand side of (3.2.7) as $\lambda \rightarrow \infty$. The possible candidates are the first on-diagonal term of the parametrix and the 0-th off-diagonal terms of the parametrix. As we shall prove below, in dimensions $n \leq 5$ the contribution of 0-th off-diagonal terms provides the principal contribution, while if $n \geq 6$ this is done by the first on-diagonal term.

In the sequel, we shall consider the cases of n even and n odd separately.

3.3. Leading term in even dimensions. Let the dimension $n = 2m$ be even. The formula (3.1.1) becomes

$$(3.3.1) \quad E(t, x, y) = \frac{1}{\pi^{m-1/2}} |t| \sum_{j=0}^{\infty} (-1)^j u_j(x, y) \frac{(r^2 - t^2)_-^{j-m-1/2}}{4^j \Gamma(j - m + 1/2)} \pmod{C^\infty}.$$

Lemma 3.3.2. *The contribution of the leading ($j = 1$) on-diagonal term of the parametrix as $\lambda \rightarrow \infty$ in (3.2.7) is $o(1/T)$ for $m = 1$, and*

$$\int_{-\infty}^{\infty} \frac{\psi(t/T) t^{2-2m} u_1(x, x) \cos(\lambda t) dt}{4\pi^{m-\frac{1}{2}} \Gamma(3/2 - m) T} = Q_1 \frac{\lambda^{2m-3}}{T} + O(\lambda^{2m-5})$$

for $m \geq 2$, where Q_1 is some non-zero constant.

Proof. This lemma can be extracted from [Ber, p. 266, formula (56)]. For $m = 1$ the expression under the integral is smooth and the statement follows from the Riemann-Lebesgue lemma. For $m \geq 2$ one should expand $\psi(t/T)$ into Taylor series near 0 up to the order $2m - 2$. Note that all terms containing odd powers of t vanish after the integration. Using the fact that the Fourier transform of the generalized function $t^{-2\alpha}$ is of order $\lambda^{2\alpha-1}$ ([G-S]), we obtain the principal term and the error estimate. Note also that $u_1(x, x) = \tau(x)/6$, where $\tau(x) < 0$ is the scalar curvature of M , and hence $u_1(x, x) \neq 0$. \square

We do not compute the constant Q_1 (as well as the constants Q_2, \dots, Q_6 defined below) since their explicit value is not important. It only matters that all these constants are non-zero which one can easily check.

Lemma 3.3.3. *The contribution of the leading (0-th) off-diagonal term of the parametrix to (3.1.2) as $\lambda \rightarrow \infty$ is*

$$\int_r^\infty \frac{\psi(t/T) u_0(x, y) t \cos(\lambda t) dt}{\pi^{m-\frac{1}{2}} T \Gamma(\frac{1}{2} - m) (t^2 - r^2)^{m+\frac{1}{2}}} = Q_2 \frac{\psi(r/T) \lambda^{m-\frac{1}{2}} \sin(\lambda r + \phi_m)}{\sqrt{g(x, y) r^{2m-1}} T} + O(\lambda^{m-\frac{3}{2}}),$$

where Q_2 is some nonzero constant, $r = d(x, y)$, and $\phi_m = \pi/4$ if m is odd, $\phi_m = 3\pi/4$ if m is even.

Proof. The orders of the leading term and the remainder follow from [Ber, pp. 267-268]. To compute the leading term explicitly, we rewrite the left-hand side as

$$\int_r^\infty \frac{\psi(t/T) u_0(x, y) t \cos(\lambda t) dt}{\pi^{m-\frac{1}{2}} T \Gamma(\frac{1}{2} - m) (t^2 - r^2)^{m+\frac{1}{2}}} = Q'_2 \int_r^\infty \frac{\psi(t/T) t \cos(\lambda t)}{T \sqrt{g(x, y)} (t+r)^{m+\frac{1}{2}}} d\left(\frac{1}{(t-r)^{m-\frac{1}{2}}}\right),$$

where Q'_2 is some non-zero constant. Let us integrate by parts m times. In order to get the highest power of λ , all the differentiations should be applied to $\cos \lambda t$. Otherwise, the power of λ decreases at least by 1 which implies the error estimate (the same argument is used in Lemma 3.4.4). Differentiating a cosine m times one gets either a cosine or a sine, depending on the parity of m . This explains different phases ϕ_m for m even and odd. The last step is to make a change of variables $s = t - r$ and to obtain an Erdelyi-type integral whose asymptotics is well-known (cf. [Don, Proposition A.1]). \square

Remark 3.3.4. In [K, Lemma III.2] an equivalent statement is obtained for $m = 1$.

3.4. Leading term in odd dimensions. Let $n = 2m + 1$ be odd. The formula (3.1.1) becomes

$$(3.4.1) \quad E(t, x, y) = \frac{|t|}{\pi^m} \sum_{j=0}^{\infty} (-1)^j u_j(x, y) \frac{(r^2 - t^2)_-^{j-m-1}}{4^j \Gamma(j-m)} \pmod{C^\infty}.$$

Lemma 3.4.2. *The contribution of the the first ($j = 1$) on-diagonal term of (3.4.1) to (3.2.7) is*

$$(3.4.3) \quad \frac{(-1)^m u_1(x, x) (m-1)!}{4\pi^m (2m-2)! T} \int_{-\infty}^{\infty} \psi(t/T) \cos(\lambda t) \delta^{(2m-2)}(t) dt = Q_3 \frac{\lambda^{2m-2}}{T} + O(\lambda^{2m-4}),$$

where Q_3 is some nonzero constant.

Proof. This lemma can be extracted from [Ber, p. 268]. To get the left hand side of (3.4.3) from (3.4.1) we use the first equality in (3.2.6). The asymptotics then follows from the fact that in order to get the highest power of λ , all $2m - 2$ derivations coming from $\delta^{(2m-2)}(t)$ should be applied to $\cos(\lambda t)$. Note that only an even number of derivations applied to $\cos(\lambda t)$ produces a non-zero contribution, hence the orders of the principal and the error terms differ by 2. \square

In order study the contribution of the leading off-diagonal term of (3.4.1) in (3.1.2) we transform it as follows:

$$\begin{aligned} \frac{(-1)^m u_0(x, y) |t| (r^2 - t^2)_-^{m-1}}{\pi^m \Gamma(-m)} &= \frac{u_0(x, y) |t|}{\pi^m} (-1)^m \delta^{(m)}(r^2 - t^2) \\ &= \frac{u_0(x, y) |t| (-1)^m}{\pi^m (r+t)^{m+1}} \delta^{(m)}(r-t) = \frac{u_0(x, y) |t|}{\pi^m (r+t)^{m+1}} \delta^{(m)}(t-r) \end{aligned}$$

Lemma 3.4.4. *The contribution of the leading (0-th) off-diagonal term of (3.4.1) in (3.1.2) is equal to*

$$\frac{-u_0(x, y)}{T\pi^m} \int_{-\infty}^{\infty} \frac{\psi(t/T)|t|\cos(\lambda t)}{(r+t)^{m+1}} \delta^{(m)}(t-r) dt = Q_4 \frac{\lambda^m \psi(r/T) \sin(\lambda r + \phi_m)}{\sqrt{g(x, y)} r^m T} + O(\lambda^{m-1}),$$

where Q_4 is some nonzero constant, $r = d(x, y)$, and $\phi_m = 0$ if m is odd, $\phi_m = \pi/2$ if m is even.

Proof. This lemma follows from [Ber, p. 269]. In order to compute the principal term explicitly one has to integrate m times by parts applying all derivations to $\cos(\lambda t)$. Different phases ϕ_m for m even and odd appear for the same reason as in the proof of Lemma 3.3.3. The principal term is given by

$$Q'_4 \lambda^m \frac{u_0(x, y)}{T\pi^m} \int_{-\infty}^{\infty} \frac{\psi(t/T)|t|\sin(\lambda t + \phi_m)}{(r+t)^{m+1}} \delta(t-r) dt = Q_4 \frac{\lambda^m \psi(r/T) \sin(\lambda r + \phi_m)}{\sqrt{g(x, y)} r^m T},$$

where Q_4, Q'_4 are some non-zero constants. \square

3.5. Leading term: summary.

Proposition 3.5.1. *The integral $K_{\lambda, T}(x, y)$ defined by (2.3.2) satisfies for any $x \neq y \in M$ as $\lambda \rightarrow \infty$:*

$$(3.5.2) \quad K_{\lambda, T}(x, y) = \frac{Q_5 \lambda^{\frac{n-1}{2}} \psi(r/T)}{T \sqrt{g(x, y)} r^{n-1}} \sin(\lambda r + \phi_n) + O(\lambda^{\frac{n-3}{2}}).$$

Here $r = d(x, y)$, $\phi_n = \frac{\pi}{4}(3 - (n \bmod 4))$, Q_5 is a non-zero constant.

Proposition 3.5.3. *Assume that $n > 3$. Then the integral $\tilde{K}_{\lambda, T}(x)$ defined by (3.2.8) satisfies as $\lambda \rightarrow \infty$:*

$$(3.5.4) \quad \tilde{K}_{\lambda, T}(x) = \frac{Q_6 \lambda^{n-3}}{T} + O(\lambda^{n-5}),$$

where Q_6 is a non-zero constant.

In dimensions $n = 2, 3$ the term $\tilde{K}_{\lambda, T}(x)$ is negligible.

Proof of Propositions 3.5.1 and 3.5.3. The propositions follow from (3.1.2) and Lemmas 3.3.2, 3.3.3, 3.4.2, 3.4.4. The terms in (3.1.2) for $j > 0$ (if $x \neq y$) and for $j > 1$ (if $x = y$) contribute only to the remainders in (3.5.2), (3.5.4). \square

3.6. Proof of Theorem 1.1.3. Let $x, y \in X$ be two arbitrary points, r be the distance between them. Assume that x and y are not conjugate along any shortest geodesics joining them (and since the geodesics are shortest, they contain no conjugate points between x and y as well). In particular, this implies that there is only a finite number of shortest geodesics joining x and y . Moreover, there exists $\epsilon > 0$ such that any other geodesic joining x and y has length greater than $r + \epsilon$. Take $T = r + \epsilon$ and consider the formula (2.1.4). The function ψ is supported on $[-1, 1]$, hence one can approximate the wave kernel $e(t, x, y)$ in (2.1.4) by summing up the parametrices (3.1.1) (cf. [Kan, Theorem 4.3]) along geodesics of length r only. Since x and y are not conjugate along such geodesics, the parametrices are well defined and the difference between $e(t, x, y)$ and the sum of the parametrices is a smooth function in t for $t \in [0, T]$. Therefore, we can apply the results of the

previous section working directly on X (and not on M). In particular Lemma 3.5.1 gives us

$$(3.6.1) \quad k_{\lambda,T}(x, y) = Q_7 \lambda^{\frac{n-1}{2}} \sin(\lambda r + \phi_n) + O(\lambda^{\frac{n-3}{2}}),$$

where Q_3 is some nonzero constant. Assume for contradiction $N_{x,y}(\lambda) = o(\lambda^{\frac{n-1}{2}})$. Set $a = \frac{n-1}{2}$ in Lemma 2.2.1. Compare Lemma 2.2.1 with (3.6.1) and fix a small number $\nu > 0$. We obtain a contradiction choosing a sequence $\lambda_k \rightarrow \infty$, such that $|\sin(\lambda_k r + \phi_n)| > \nu$. \square

3.7. Proof of the Theorem 1.1.5. First, let us prove (1.1.6). In this case, we also work directly on X . Take $T = \epsilon > 0$, where ϵ is small enough so that any geodesic loop returning to x has length bigger than ϵ . Throughout sections 3.1-3.6 we assumed that X has negative curvature, which implied $u_1(x, x) \neq 0$ at any point $x \in X$. In order to adjust Proposition 3.5.3 for an arbitrary manifold set $\kappa_x = \min\{j \geq 1 \mid u_j(x, x) \neq 0\}$ as in section 1.1. If $u_j(x, x) = 0$ for all $j \geq 1$ we set $\kappa_x = \infty$. Then if $n - 2\kappa_x - 1 > 0$, (3.5.4) takes the form

$$(3.7.1) \quad k_{\lambda,T}(x, x) = \frac{Q_8 \lambda^{n-2\kappa_x-1}}{T} + O(\lambda^{n-2\kappa_x-3}).$$

Repeating the same arguments as in the previous section we complete the proof of (1.1.6).

To prove the inequality (1.1.7), assume that X is a manifold without conjugate points. Hence the parametrix is globally defined on M (see [Zel]). Consider the orbit Γx of x in the universal cover M . Let

$$r = \inf_{\text{Id} \neq \omega \in \Gamma} d(x, \omega x)$$

(it is attained for finitely many ω -s), and suppose there are no points in $\Gamma x \setminus \{x\}$ whose distance to x lies in $[r, r+\epsilon]$; choose $T = r+\epsilon/2$ in (2.3.2). Only the closest to x lattice points contribute to the right hand side of (2.3.3). This contribution is of order $\lambda^{\frac{n-1}{2}}$. The contribution of $\omega = \text{Id}$ is of order $O(\lambda^{n-2\kappa_x-1})$. If $n - 4\kappa_x - 1 < 0$, $\lambda^{\frac{n-1}{2}}$ gives a better lower bound. This completes the proof of (1.1.7). \square

Clearly, applicability of (1.1.6) and (1.1.7) depends on the geometry of a manifold. For instance, on flat tori (1.1.6) does not give any information since $u_j(x, x) = 0$ for all $j \geq 1$. However, tori have no conjugate points and one can use (1.1.7).

4. SPECTRAL ESTIMATES FROM BELOW ON NEGATIVELY CURVED MANIFOLDS

4.1. Sums over geodesic segments. Let X be a negatively curved manifold with the sectional curvature satisfying (1.2.1), M be the universal cover of X , and let $x, y \in M$. Since X has no conjugate points, the parametrix is well-defined for all times on M .

Our strategy to strengthen Theorems 1.1.3 and 1.1.5 is as follows. Let T grow with λ and assume that we can force all the $\sin(\lambda r + \phi_n)$ in Proposition 3.5.1 be of the same sign and bounded away from zero (see section 5.2). Consider the sum over geodesic segments starting at the point x :

$$(4.1.1) \quad S_{x,y}(T) = \sum_{r_\omega \leq T, \omega \in \Gamma} \frac{1}{\sqrt{g(x, \omega y)} r_\omega^{n-1}},$$

where $x \neq \omega y$ for all $\omega \in \Gamma$, $r_\omega = d(x, \omega y)$ and $g = \sqrt{\det g_{ij}}$. The idea is to estimate (4.1.1) from below by a function going to infinity as $\lambda \rightarrow \infty$.

Theorem 4.1.2. *There exists a constant $C_0 > 0$ such that*

$$(4.1.3) \quad S_{x,y}(T) \geq \frac{C_0}{T} e^{P(-\frac{\mathcal{H}}{2}) \cdot T}$$

as $T \rightarrow \infty$, where P is the topological pressure (1.2.3) and \mathcal{H} is the Sinai-Ruelle-Bowen potential (1.2.2).

It was shown in (1.3.5) that $P(-\frac{\mathcal{H}}{2}) \geq (n-1)K_2/2$, hence $S_{x,y}(T)$ grows exponentially in T .

In the Appendix we shall prove a stronger version of (4.1.3):

$$(4.1.4) \quad S_{x,y}(T) \geq C_0 e^{P(-\frac{\mathcal{H}}{2}) \cdot T}$$

However, a weaker estimate (4.1.3) is sufficient for the proofs of Theorems 1.3.1 and 1.3.3.

The proof of Theorem 4.1.2 is divided into several steps described below.

4.2. Jacobi fields and the geodesic flow. Fix a point $x \in M$ and let $\gamma(t)$ be a minimizing geodesic from x to $y \in M$ such that $\gamma(0) = x$, $\gamma(r) = y$, $\gamma'(0) = v$, where $v \in T_x M$ is a unit vector. Choose an orthonormal basis in $T_x M$ consisting of v and vectors e_1, e_2, \dots, e_{n-1} . To every vector e_j , $j = 1, \dots, n-1$, there corresponds a perpendicular Jacobi field $Y_j(t)$ along the geodesic γ with the initial conditions $Y_j(0) = 0$, $Y_j'(0) = e_j$. Then (see [Ber, p. 27])

$$(4.2.1) \quad g(x, y) = \frac{\sqrt{\det(Y_j(r), Y_k(r))}}{r^{n-1}}$$

Let us recall the construction of the canonical (Sasaki) metric on the tangent bundle TM (see [E]). Let $\Pi : TM \rightarrow M$ be the projection map and let $\xi = (x, v) \in TM$, $\pi(\xi) = x$. Let $\mathcal{K} : T_\xi(TM) \rightarrow T_x M$ be the connection map. There exists a canonical splitting $T_\xi TM = T_\xi(T_x M) \oplus \text{Horiz}(\xi)$ into vertical $\text{Im } \mathcal{K} = T_\xi(T_x M)$ and horizontal $\text{Horiz}(\xi) = \text{Im } d\pi$ subspaces. For any two vectors $z, w \in T_\xi(TM)$ the Sasaki metric on TM is defined by

$$(4.2.2) \quad (z, w)_{T_\xi(TM)} = (d\Pi z, d\Pi w)_{T_x M} + (\mathcal{K}z, \mathcal{K}w)_{T_x M}.$$

The induced metric on the unit tangent bundle SM is also referred to as Sasaki metric.

Consider the geodesic flow $G^t : TM \rightarrow TM$. Then we have ([E]):

$$(4.2.3) \quad Y_j(t) = d\Pi \circ dG^t(\tilde{e}_j), \quad Y_j'(t) = \mathcal{K} \circ dG^t(\tilde{e}_j),$$

where $\tilde{e}_j = \mathcal{K}^{-1}e_j$, $j = 1, \dots, n-1$. From the definition of the Sasaki metric it follows that since e_j are orthonormal vectors, \tilde{e}_j are also orthonormal.

4.3. Jacobian of the geodesic flow. Consider now the unit tangent bundle SM . On negatively-curved manifolds there exists another natural splitting of $T_\xi(SM)$ into a direct sum of DG^t -invariant subspaces. It comes from the Anosov property of the geodesic flow G^t on SM :

$$(4.3.1) \quad T_\xi(SM) = E_\xi^u \oplus E_\xi^o \oplus E_\xi^s.$$

Here E_ξ^u is the *unstable* subspace of dimension $(n-1)$, E_ξ^s is the *stable* subspace of dimension $(n-1)$, and E_ξ^o is a one-dimensional subspace tangent to the flow.

Lemma 4.3.2. *There exists a universal constant $C_1 > 0$ such that*

$$(4.3.3) \quad \sqrt{g(x, y)} r^{\frac{n-1}{2}} < C_1 \det dG^r|_{E_\xi^u},$$

where $\Pi(\xi) = x, r = d(x, y) > 0$ and E_ξ^u is the unstable subspace of $T_\xi(SM)$.

Proof. Due to (4.2.1) and (4.2.3) it remains to show that

$$(4.3.4) \quad \det(d\Pi \circ dG^r(\tilde{e}_j), d\Pi \circ dG^r(\tilde{e}_k)) \leq C_1^2 (\det dG^r|_{E_\xi^u})^2$$

for some constant $C_1 > 0$. First, we argue that

$$(4.3.5) \quad \det(dG^r(\tilde{e}_j), dG^r(\tilde{e}_k))_{T_\xi(SM)} \geq \det(d\Pi \circ dG^r(\tilde{e}_j), d\Pi \circ dG^r(\tilde{e}_k)),$$

where the scalar products on the left are taken with respect to the Sasaki metric (4.2.2). Indeed, the matrix on the left is the sum of the matrix on the right and a symmetric positive definite matrix (coming from the second term in (4.2.2)) and hence it has a greater determinant.

Denote $\text{Vert}(\xi) = T_\xi(S_x M)$ the vertical component of the canonical splitting of SM . Note that $\det(dG^r(\tilde{e}_j), dG^r(\tilde{e}_k))_{T_\xi(SM)} = (\text{Jac}_{\text{Vert}(\xi)} G^r)^2$, where $\text{Jac}_{\text{Vert}(\xi)} G^r$ is the Jacobian of the map G^r restricted to $\text{Vert}(\xi)$ ([H-K, p. 607]). We need to prove

$$(4.3.6) \quad \frac{\text{Jac}_{\text{Vert}(\xi)} G^r}{\text{Jac}_{E_\xi^u} G^r} \leq C_1.$$

It is sufficient to prove this for an integer r . Indeed, a volume comparison argument implies $\frac{\text{Jac}_{\text{Vert}(\xi)} G^r}{\text{Jac}_{\text{Vert}(\xi)} G^{[r]}} \leq c$, where c is a universal constant depending only on the curvature bounds. At the same time, it is known that $\text{Jac}_{E_\xi^u} G^t$ is a monotone increasing function in t and hence $\text{Jac}_{E_\xi^u} G^r \geq \text{Jac}_{E_\xi^u} G^{[r]}$. Taking a logarithm in (4.3.6) and writing $G^r = G^1 \circ \dots \circ G^1$ (r times) we get:

$$|\log \text{Jac}_{\text{Vert}(\xi)} G^r - \log \text{Jac}_{E_\xi^u} G^r| \leq \sum_{i=1}^{r-1} |\log \text{Jac}_{dG^i \text{Vert}(\xi)} G^1 - \log \text{Jac}_{E_{G^i \xi}^u} G^1|.$$

Let us show that the distance between the subspaces $dG^i \text{Vert}(\xi)$ and $E_{G^i \xi}^u$ is converging to zero exponentially fast. Here distance is understood in the following sense (see [ArAv, p. 191]). For any $v \in dG^i \text{Vert}(\xi)$ let $v = v_u + v_s$ where $v_u \in E_{G^i \xi}^u$, $v_s \in E_{G^i \xi}^s$. Set

$$(4.3.7) \quad \text{dist}(dG^i \text{Vert}(\xi), E_{G^i \xi}^u) = \sup_{\|v\|=1, v \in dG^i \text{Vert}(\xi)} \frac{\|v_s\|}{\|v_u\|}$$

It then follows from the Anosov property of the geodesic flow and the results of [E, p. 456] that (4.3.7) is well defined and

$$\text{dist}(dG^i \text{Vert}(\xi), E_{G^i \xi}^u) \leq C_1' e^{-\alpha i}$$

for some positive constants C_1' and α . We now remark that $\log \text{Jac}_V G^1$ depends smoothly on a subspace $V \subset T_\eta(SM)$, and therefore

$$|\log \text{Jac}_{dG^i \text{Vert}(\xi)} G^1 - \log \text{Jac}_{E_{G^i \xi}^u} G^1| \leq C_1'' e^{-\alpha i},$$

where C_1'' is some positive constant that can be chosen independently of i by a compactness argument (G^1 on SM is a lift of G^1 from a compact manifold SX). Hence

$$|\log \text{Jac}_{\text{Vert}(\xi)} G^r - \log \text{Jac}_{E_\xi^u} G^r| \leq C_1'' \sum_{i=0}^{r-1} e^{-\alpha i} \leq \frac{C_1''}{1 - e^{-\alpha}}.$$

This completes the proof of the lemma. \square

As an immediate corollary of Lemma 4.3.3 we get

$$(4.3.8) \quad S_{x,y}(T) \geq \frac{1}{C_1} \sum_{r_\omega \leq T, \omega \in \Gamma} \frac{1}{\sqrt{\det dG^{r_\omega}|_{E_{\xi_\omega}^u}}},$$

where $\xi_\omega = (x, v_\omega) \in SM$, $v_\omega = \gamma'_\omega(0)$, γ_ω is the shortest geodesic joining x and ωy .

Denote

$$(4.3.9) \quad Z(r_\omega, \xi_\omega) = \int_0^{r_\omega} \frac{d}{d\tau} \Big|_{\tau=0} \ln \det \left(dG^{s+\tau} \Big|_{E_{\xi_\omega}^u} \right) ds = \ln \det dG^{r_\omega} \Big|_{E_{\xi_\omega}^u} = \int_0^{r_\omega} \mathcal{H}(G^s \xi_\omega) ds,$$

where \mathcal{H} is the Sinai-Ruelle-Bowen potential (1.2.2). Here we used the invariance of the unstable foliation with respect to the geodesic flow. Hence,

$$(4.3.10) \quad S_{x,y}(T) \geq \frac{1}{C_1} \sum_{r_\omega \leq T, \omega \in \Gamma} \exp \left(-\frac{1}{2} \int_0^{r_\omega} \mathcal{H}(G^s \xi_\omega) ds \right).$$

4.4. From geodesic segments to closed geodesics. Our next aim is to estimate from below the sum (4.3.8) taken over geodesic segments starting at x by a sum over closed geodesics, and then apply techniques of the thermodynamic formalism. It is well-known that each conjugacy class $[\omega] \subset \Gamma$ corresponds to a unique closed geodesic $\gamma_{[\omega]}$ on X of length $l_{[\omega]}$. Choose a representative $\omega_c \in [\omega]$ corresponding to this closed geodesic and assume that it joins $q \in M$ and $\omega_c q \in M$, such that $d(x, q) \leq D$ and $d(\omega y, \omega_c q) \leq D$, where D is the diameter of X . Set $\xi_{[\omega]} = (q, \gamma'_{[\omega]}(0)) \in SM$. To distinguish between distances on M and on SM we shall write d_M and d_{SM} respectively. Here d_{SM} is induced by the Sasaki metric.

Lemma 4.4.1. *There exists $C_2 > 0$ such that for any $x, y \in M$ and $\omega \in \Gamma$, $d(x, \omega y) \geq D$ we have*

$$(4.4.2) \quad |Z(r_\omega, \xi_\omega) - Z(l_{[\omega]}, [\xi_{[\omega]}])| \leq C_2.$$

Let $\gamma_1 = [x_1, y_1]$ and $\gamma_2 = [x_2, y_2]$ be two geodesic segments in M such that $d_M(x_1, x_2) < D$ and $d_M(y_1, y_2) < D$. We shall prove the following general statement implying Lemma 4.4.1:

$$(4.4.3) \quad |Z(r_1, \xi_1) - Z(r_2, \xi_2)| \leq C_2,$$

where $\xi_j = (x_j, \gamma'_j(0))$, $j = 1, 2$ and $r_j = d_M(x_j, y_j)$.

Remark 4.4.4. It suffices to prove (4.4.3) for two geodesics starting at the same point on M : apply it first to (y_1, z_1) and (y_1, z_2) , then to (y_1, z_2) and (y_2, z_2) .

Accordingly, let $v_1, v_2 \in SM$, such that $\Pi(v_1) = x_1 = x_2 = \Pi(v_2)$. Let $\gamma_{v_1}(t) = \gamma_1(t)$ and $\gamma_{v_2}(t) = \gamma_2(t)$ be the corresponding geodesics in M .

Note: to simplify notations, instead of writing $(\gamma(s), \gamma'(s))$ for an element of SM we shall write $\gamma'(s)$.

A key ingredient in the proof of Lemma 4.4.1 is the following

Proposition 4.4.5. *Suppose that $d_M(\gamma_1(t), \gamma_2(t)) \leq D$. Then there exist two constants $A_1 > 0, B_1 > 0$ such that for $0 \leq s \leq t$,*

$$(4.4.6) \quad d_{SM}(\gamma_1'(s), \gamma_2'(s)) \leq A_1 e^{B_1(s-t)}$$

4.5. Proof of Proposition 4.4.5. This proof was communicated to the authors by D. Dolgopyat. We first remark that it follows from [Schr, Lemma 2.1] that there exists $\alpha_1 > 0$ such that

$$(4.5.1) \quad \text{dist}(\gamma_1(s), \gamma_2(s)) \leq \alpha_1 e^{K_2(s-t)},$$

thus establishing the desired estimate on M .

It follows easily from the definition of d_{SM} that

$$(4.5.2) \quad d_{SM}(v_1, v_2) \leq \sqrt{d_M(x_1, x_2)^2 + \pi^2}$$

By increasing α_1 if necessary, we may assume without loss of generality that (4.5.1) holds for $s \leq t + 1$.

We remark that it follows easily from (4.5.1) and (4.5.2) that (4.4.6) would hold if $d_M(\gamma_1(s+1), \gamma_2(s+1)) \geq 1/3$. Accordingly, it suffices to establish (4.4.6) for such $0 < s < t$ that $d_M(\gamma_1(s+1), \gamma_2(s+1)) < 1/3$.

Consider now the following four points on M :

$$(4.5.3) \quad P_1 = \gamma_1(s), P_2 = \gamma_2(s), Q_1 = \gamma_1(s+1), Q_2 = \gamma_2(s+1).$$

We know that

$$\begin{aligned} d_1 &:= d_M(P_1, P_2) \leq \alpha_1 e^{K_2(s-t)}, d_1 \leq 1/3; \\ d_2 &:= d_M(Q_1, Q_2) \leq \alpha_1 e^{K_2(s+1-t)}, d_2 \leq 1/3; \\ d_M(P_1, Q_1) &= d_M(P_2, Q_2) = 1. \end{aligned}$$

It follows from the triangle inequality that

$$(4.5.4) \quad 1 - d_1 \leq d_3 := d_M(P_1, Q_2) \leq 1 + d_1.$$

Denote by $\gamma_3(s), 0 \leq s \leq d_3$ the geodesic segment connecting P_1 and Q_2 on M . Consider now the following vectors in SM :

$$(4.5.5) \quad w_1 = \gamma_1'(s), w_2 = \gamma_2'(s), w_3 = \gamma_3'(0), w_4 = -\gamma_2'(s+1), w_5 = -\gamma_3'(d_3).$$

The idea is to estimate $d_{SM}(w_1, w_2)$ using the triangle inequality

$$(4.5.6) \quad d_{SM}(w_1, w_2) \leq d_{SM}(w_1, w_3) + d_{SM}(w_2, w_3).$$

We first remark that $d_{SM}(w_1, w_3)$ is equal to the angle at the vertex P_1 in the geodesic triangle $P_1Q_1Q_2$ with sides d_1, d_2, d_3 . We compare the triangle $P_1Q_1Q_2$ with the planar triangle having the same sides. It follows from $d_2 < 1/3$, (4.5.4), elementary planar trigonometry and comparison theorem that the angle $\theta = \angle Q_1P_1Q_2$ is less than $\pi/6$. Hence, $\sin \theta > 3\theta/\pi$. Therefore we have

$$(4.5.7) \quad d_{SM}(w_1, w_3) = \theta < \frac{\pi \sin \theta}{3} < \frac{\pi d_2}{3d_3} < \frac{\pi d_2}{2} \leq \frac{\pi \alpha_1 e^{K_2}}{2} e^{K_2(s-t)}$$

Here we use that $\sin \theta < d_2/d_3$ which follows from comparison with a planar triangle, and that $d_3 > 2/3$ which follows from the triangle inequality. It remains to estimate $d_{SM}(w_2, w_3) = d_{SM}(-w_2, -w_3)$. Consider the vector $w_6 = -\gamma_3'(d_3 - 1)$. We see that

$$d_{SM}(-w_2, -w_3) \leq d_{SM}(-w_2, w_6) + d_{SM}(w_6, -w_3) \leq d_{SM}(-w_2, w_6) + d_1,$$

so it suffices to estimate $d_{SM}(-w_2, w_6)$.

Now,

$$-w_2 = G^1(w_4), \quad w_6 = G^1(w_5),$$

where G^1 is the time one map of the geodesic flow G^t . The distance between w_4 and w_5 is equal to the angle at Q_2 of the geodesic triangle $P_1Q_2P_2$ with sides $d_3, 1, d_1$. By comparing that triangle to the isometric planar triangle, one can show as in (4.5.7) that

$$(4.5.8) \quad d_{SM}(w_4, w_5) \leq \frac{\pi d_1}{2} \leq \frac{\pi \alpha_1}{2} e^{K_2(s-t)}.$$

Now, consider the smooth diffeomorphism G^1 restricted to the unit tangent space of the quadrilateral $P_1Q_1P_2Q_2$. By compactness it increases distances by at most a factor α_2 . It follows from (4.5.8) that

$$(4.5.9) \quad d_{SM}(-w_2, w_6) \leq \alpha_2 d_{SM}(w_4, w_5) \leq \frac{3\pi \alpha_1 \alpha_2}{4} e^{K_2(s-t)}.$$

The inequality (4.4.6) now follows from (4.5.6), (4.5.7) and (4.5.9). This finishes the proof of the proposition. \square

4.6. Proof of Lemma 4.4.1. By Remark 4.4.4, in order to prove (4.4.3) it suffices to consider two vectors $\xi_1, \xi_2 \in SM$ with $\pi(\xi_1) = \pi(\xi_2)$. Let γ_1 and γ_2 be geodesics as in Lemma 4.4.5: say, $\gamma_1 = [x, \omega y]$ and $\gamma_2 = [x, \omega_c q]$. Let l_1, l_2 be the lengths of γ_1, γ_2 respectively. It follows from the triangle inequality that $|l_1 - l_2| \leq D$. It thus suffices to estimate the difference

$$|Z(r_1, \xi_1) - Z(r_2, \xi_2)| = \left| \int_0^{r_1} \mathcal{H}(G^s \xi_1) ds - \int_0^{r_2} \mathcal{H}(G^s \xi_2) ds \right|$$

Since \mathcal{H} is uniformly bounded on SX by compactness, and thus also on SM , the inequality (4.4.3) will remain true (with a different constant) if we increase or decrease the length of γ_j by a uniformly bounded amount. Since $|l_1 - l_2| \leq D$, we may thus assume without loss of generality that γ_j -s have the same length t .

Accordingly, to prove (4.4.2), it suffices to show that there exist $B_2 > 0$ such that for $t \geq D$,

$$(4.6.1) \quad \left| \int_0^t [\mathcal{H}(G^s \xi_1) - \mathcal{H}(G^s \xi_2)] ds \right| \leq B_2.$$

It follows from the Hölder continuity of H ([B-R],[Sin2]) that there exist two constants $A_2 > 0, \beta > 0$ such that for any $v_1, v_2 \in SM$,

$$|\mathcal{H}(v_1) - \mathcal{H}(v_2)| \leq A_2 \cdot d_{SM}(v_1, v_2)^\beta.$$

It follows from Proposition 4.4.5 exist two constants $A_1, B_1 > 0$ such that for $0 \leq s \leq t$,

$$d_{SM}(G^s \xi_1, G^s \xi_2) \leq A_1 e^{B_1(s-t)}$$

Call the integral in (4.6.1) $I(t)$. Now,

$$\begin{aligned} |I(t)| &\leq \int_0^t |\mathcal{H}(G^s \xi_1) - \mathcal{H}(G^s \xi_2)| ds \leq A_2 \int_0^t d_{SM}(G^s \xi_1, G^s \xi_2)^\beta ds \\ &\leq A_2 A_1^\beta \int_0^t e^{B_1 \beta (s-t)} ds \leq \frac{A_2 A_1^\beta}{B_1 \beta}. \end{aligned}$$

This proves (4.6.1) with $B_2 = A_2 A_1^\beta / (B_1 \beta)$. The proof of Lemma 4.4.1 is complete. \square

4.7. **Proof of Theorem 4.1.2.** By Lemma 4.4.1 and (4.3.10) we have

$$(4.7.1) \quad S_{x,y}(T) \geq C_3 \sum_{\omega \in \Gamma: 3D \leq l_{[\omega]} \leq T-2D} \exp\left(-\frac{Z(l_{[\omega]}, \xi_{[\omega]})}{2}\right)$$

for some constant $C_3 > 0$. We sum over the interval $3D \leq l_{[\omega]} \leq T - 2D$ since by triangle inequality the difference between the length of the segment and the corresponding closed geodesic is at most $2D$, and due to the condition of Lemma 4.4.1 we consider segments of length $\geq D$. It is a relatively rough bound. Indeed, we take into account the contribution of just one element from every conjugacy class. In the Appendix we refine this estimate taking into account the number of elements in each conjugacy class.

We now apply quite a deep fact on the equidistribution of closed geodesics for Anosov flows proved by methods of thermodynamic formalism. Using (4.7.1) we write:

$$(4.7.2) \quad S_{x,y}(T) \geq \frac{C_3}{T} \sum_{\omega \in \Gamma: 3D \leq l_{[\omega]} \leq T-2D} l_{[\omega]} \exp\left(-\frac{Z(l_{[\omega]}, \xi_{[\omega]})}{2}\right)$$

On the other hand, the sum (4.7.2) over closed geodesics satisfies the following asymptotic relation for $T \rightarrow \infty$: (cf. [Par], [P-P, (7.1)] and [M-S, p. 109]),

$$(4.7.3) \quad \sum_{l_{[\omega]} \leq T} l_{[\omega]} \exp\left(-\frac{Z(l_{[\omega]}, \xi_{[\omega]})}{2}\right) = \frac{C_4}{P(-\mathcal{H}/2)} e^{P(-\mathcal{H}/2)T} (1 + o(1)),$$

where

$$C_4 = \int_{SX} d\mu_{-\mathcal{H}/2}.$$

Here $\mu_{-\mathcal{H}/2}$ is the *equilibrium state* for $-\mathcal{H}/2$ and $P(-\mathcal{H}/2)$ is the pressure defined by (1.2.3). As $T \rightarrow \infty$, the contribution of closed geodesics of lengths $l_{[\omega]} \leq 3D$ and $T - 2D \leq l_{[\omega]} \leq T$ to the sum in (4.7.3) is bounded by a constant. Therefore, (4.7.3) implies (4.1.3) and this completes the proof of Theorem 4.1.2. \square

5. PROOF OF THE MAIN RESULTS.

5.1. **Proof of Theorem 1.3.3 for $n \not\equiv 3 \pmod{4}$.** Given $\delta > 0$, let

$$\alpha = \frac{(1 - \delta)P(-\mathcal{H}/2)}{h}.$$

Assume for contradiction that the bound (1.3.4) doesn't hold and thus for some $x, y \in X, x \neq y$, the off-diagonal spectral counting function $N_{x,y}(\lambda)$ satisfies the following upper bound:

$$(5.1.1) \quad N_{x,y}(\lambda) = O\left(\lambda^{(n-1)/2}(\log \lambda)^\alpha\right).$$

By Lemma 2.2.4 we conclude that there exists a constant $C_5 > 0$ such that for any λ we have

$$(5.1.2) \quad |k_{\lambda,T}(x, y)| < C_5 \lambda^{(n-1)/2} (\log \lambda)^\alpha.$$

Using (2.3.3) we write

$$k_{\lambda,T}(x, y) = \sum_{\omega \in \Gamma} K_{\lambda,T}(x, \omega y).$$

We want to estimate the right-hand side from below. It follows from Proposition 3.5.1 that

$$(5.1.3) \quad k_{\lambda,T}(x,y) = \sum_{\omega \in \Gamma, r_\omega \leq T} \frac{Q\lambda^{\frac{n-1}{2}}}{T} \frac{\psi(\frac{r_\omega}{T})}{\sqrt{g(x,\omega y) r_\omega^{n-1}}} \sin(\lambda r_\omega + \phi_n) + O(\lambda^{\frac{n-3}{2}}) \exp(O(T)).$$

The sum in (5.1.3) can be taken over $r_\omega \leq T$ since $\text{Supp } \psi = [-1, 1]$. By results of Margulis ([M-S]), the number of summands in (5.1.3) is less than $C_6 e^{hT}$ for some positive constant C_6 . Since the error for each term is $O(\lambda^{(n-3)/2})$ by Proposition 3.5.1, for the whole sum the error is $O(\lambda^{\frac{n-3}{2}}) \exp(O(T))$.

5.2. Dirichlet box principle. We want all the terms $\sin(\lambda r_\omega + \phi_n)$ in (5.1.3) to have the same sign, and $|\sin(\lambda r_\omega + \phi_n)|$ to be bounded from below by some positive constant. Let $\{r_1, r_2, \dots, r_{N_1}\}$ be all the distinct values of r_ω appearing in (5.1.3). It suffices to choose λ so that for all $1 \leq j \leq N_1$,

$$(5.2.1) \quad |e^{i\lambda r_j} - 1| < 1/10.$$

In that case all the angles $\lambda r_j + \phi_n$ are close to

$$\phi_n = \pi/4(3 - n \bmod 4).$$

Since $n \not\equiv 3 \pmod{4}$, $\sin(\lambda r_j + \phi_n) \approx \sin \phi_n$ all have the same signs and $|\sin(\lambda r_j + \phi_n)| \geq C_7$ for some positive constant C_7 .

To establish (5.2.1) we apply the following Lemma from [P-R] (a similar approach was also used in [R-S] to estimate error terms from below):

Lemma 5.2.2. *Given r_1, r_2, \dots, r_{N_1} and $M_1 > 0, Y > 1$ there exists $\lambda \in [M_1, M_1 Y^{N_1}]$ such that for all $1 \leq j \leq N_1$,*

$$|e^{i\lambda r_j} - 1| < 1/Y.$$

By Lemma 5.2.2, given $M_1 > 0$ we can choose $M_1 \leq \lambda \leq M_1 \cdot 10^{N_1}$ such that (5.2.1) will hold for all $1 \leq j \leq N_1$. We recall that $N_1 < C_6 e^{hT}$.

Moreover, given $\delta > 0$ we can choose ψ so that $\psi(x) \geq 1/2$ for $|x| \leq 1 - \delta/2$. Accordingly, if the radius r_j in (5.1.3) satisfies $r_j \leq T(1 - \delta/2)$, then $\psi(r_j/T) \geq 1/2$.

Therefore, given M_1 , there exists λ such that

$$(5.2.3) \quad \ln M_1 \leq \ln \lambda \leq \ln M_1 + (\ln 10) C_6 e^{hT}.$$

It then follows from (5.1.3) that there exists $A_3 > 0$ such that

$$(5.2.4) \quad |k_{\lambda,T}(x,y)| \geq \frac{A_3 \lambda^{(n-1)/2}}{T} \sum_{\omega: d_M(x,\omega y) \leq (1-\delta/2)T} \frac{1}{\sqrt{g(x,\omega y) r_\omega^{n-1}}} + O(\lambda^{\frac{n-3}{2}}) \exp(O(T))$$

In the sequel, we shall choose $T = O(\ln \ln \lambda)$. It follows that the error term in (5.2.4) is

$$(5.2.5) \quad O\left(\lambda^{\frac{n-3}{2}} (\log \lambda)^{O(1)}\right), \quad \text{provided } T = O(\ln \ln \lambda).$$

Using the estimate (4.1.3) proved in Theorem 4.1.2 we conclude from (5.2.4) and (5.2.5) that

$$(5.2.6) \quad |k_{\lambda,T}(x,y)| \geq \frac{B_3}{T^2} \lambda^{\frac{n-1}{2}} e^{P(-\mathcal{H}/2)(1-\frac{\delta}{2})T},$$

for some constant $B_3 > 0$.

To obtain contradiction with (5.1.2), we should find $\lambda > 0, T > 0$ satisfying $\lambda \geq M_1$ and (5.2.3)

$$C_5 \lambda^{(n-1)/2} (\log \lambda)^\alpha \leq \frac{B_3}{T^2} \lambda^{(n-1)/2} e^{P(-\mathcal{H}/2)(1-\frac{\delta}{2})T}.$$

This translates to

$$(5.2.7) \quad \begin{aligned} \ln \ln \lambda &\leq \frac{1}{\alpha} [T P(-\mathcal{H}/2)(1-\delta/2) + \ln B_3 - \ln C_5 - 2 \ln T] \\ &= \frac{h(1-\delta/2)}{1-\delta} T + \frac{1}{\alpha} [\ln B_3 - \ln C_5 - 2 \ln T] \end{aligned}$$

We can rewrite (5.2.3) as

$$(5.2.8) \quad \ln \ln M_1 \leq \ln \ln \lambda \leq hT + \ln \ln M_1 + \ln C_6 + \ln \ln 10.$$

By Lemma 5.2.2, for any M_1 we can find some λ satisfying (5.2.8) such that the estimate (5.2.4) holds.

Let β be a small constant satisfying

$$0 < \beta < \frac{\delta/2}{1-\delta}.$$

We choose M_1 in (5.2.8) to so that

$$\ln \ln M_1 = h\beta T.$$

Then (5.2.8) becomes

$$(5.2.9) \quad h\beta T \leq \ln \ln \lambda \leq h(1+\beta)T + \ln C_6 + \ln \ln 10$$

It follows that $T \approx \frac{1}{h} \ln \ln \lambda$, and thus (5.2.5) holds, implying (5.2.6).

To prove Theorem 1.3.3, it suffices to show that if we choose T large enough, then (5.2.9) would *imply* (5.2.7). Indeed, in that case *any* choice of λ satisfying (5.2.9) would automatically contradict (5.1.1), and *some* such choice exists by Lemma 5.2.2.

The largest terms in both (5.2.9) and (5.2.7) are linear in T , so it suffices to compare the coefficients of T in those inequalities. The coefficient in (5.2.7) is equal to $\frac{h(1-\delta/2)}{1-\delta}$ while the coefficient in (5.2.9) is equal to $h(1+\beta)$. By the choice of β , the coefficient of T in (5.2.9) is *smaller* than that in (5.2.7). It follows that for large T , the right-hand side of (5.2.7) is *larger* than the right-hand side of (5.2.9), therefore (5.2.9) implies (5.2.7), finishing the proof of Theorem 1.3.3 for $n \not\equiv 3(\text{mod } 4)$. \square

5.3. Proof of Theorem 1.3.3 for $n \equiv 3(\text{mod } 4)$. The only difference between the proofs in cases $n \not\equiv 3(\text{mod } 4)$ and $n \equiv 3(\text{mod } 4)$ is that in the latter case, the angle ϕ_n in Proposition 3.5.1 is equal to zero. Accordingly, the leading term in Proposition 3.5.1 is proportional to $\sin(\lambda r_\omega)$ and thus we have to ensure that λr_ω don't get too close to $\pi \mathbf{Z}$. In addition, we want to choose λ so that $\sin \lambda r_\omega$ all have the same sign. Accordingly, we cannot use Lemma 5.2.2 directly and have to replace it by a different statement.

The idea of the modification is as follows: the biggest contribution to the sum (4.1.3) in Theorem 4.1.2 comes from the terms $\omega \in \Gamma$ where $r_\omega = d_M(x, \omega y)$ is

large. Indeed, for any $t > 0$ every term in the sum $S_{x,y}(t)$ is not greater than 1, therefore the number of terms gives a trivial upper bound: $S_{x,y}(t) \leq C_6 e^{ht}$. Take

$$(5.3.1) \quad A_4 > \frac{h}{P(-\mathcal{H}/2)(1-\delta/2)}.$$

and write $S_{x,y}(T) = S_1(T) + S_2(T)$, where $S_1(T)$ is the sum taken over all ω such that $r_\omega < T/A_4$ and $S_2(T)$ corresponds to

$$(5.3.2) \quad T/A_4 \leq r_\omega \leq T.$$

Comparing the exponents we deduce that $S_1(T)$ grows slower than $S_{x,y}(T)$. Indeed, $P(-\mathcal{H}/2)(1-\delta/2)T > hT/A_4$ by (5.3.1). Therefore, for T large the contribution of $S_1(T)$ to $S_{x,y}(T)$ is negligible in the sense that $S_{x,y}(T) = S_2(T)(1+o(1))$.

Let $\{r_1, r_2, \dots, r_{N_2}\}$ be all the $N_2 = O(e^{hT})$ distinct values of r_ω appearing in $S_2(T)$. Choose Y satisfying

$$(5.3.3) \quad Y > \frac{3A_4}{\pi}.$$

By Lemma 5.2.2, given $M_2 > 0$ we can find $\mu_1 \in [M_2, M_2 Y^{N_2}]$ such that

$$(5.3.4) \quad \text{dist}(\mu_1 r_j, 2\pi\mathbf{Z}) < 1/Y, \quad 1 \leq j \leq N_2.$$

The reason for choosing $Y > 3A_4/\pi$ will be explained below.

It suffices to find λ such that $\sin \lambda r_\omega$ all have the same sign for $T/A_4 \leq r_\omega \leq T$, and to estimate all those $\sin \lambda r_\omega$ from below. Let $2\pi k_j$ be the closest multiple of 2π to $\mu_1 r_j$ in (5.3.4), and let

$$b_j = \mu_1 r_j - 2\pi k_j.$$

We shall prove the following

Lemma 5.3.5. *Let $T/A_4 \leq r_1 < r_2 < \dots < r_{N_2} \leq T$, $T \gg 1$, and let $|b_j| \leq 1/Y$, $j = 1, \dots, N_2$. Then there exists $\mu \in [0, 1]$ such that*

$$(5.3.6) \quad \sin(\mu r_j + b_j) \geq \frac{1}{2T}, \quad 1 \leq j \leq N_2.$$

Proof of Lemma 5.3.6. We want to find $\mu \in [0, 1]$ such that

$$(5.3.7) \quad \frac{1}{T} \leq \mu r_j + b_j \leq \pi - \frac{1}{T}, \quad 1 \leq j \leq m.$$

The equality (5.3.7) is equivalent to

$$\mu \in I_j := \left[\frac{1}{Tr_j} - \frac{b_j}{r_j}, \frac{\pi}{r_j} - \frac{1}{Tr_j} - \frac{b_j}{r_j} \right].$$

Thus, it suffices to show that

$$(5.3.8) \quad \emptyset \neq \bigcap_{j=1}^m I_j \cap [0, 1].$$

The maximum P_1 of the left endpoints of I_j is given by

$$P_1 = \max_j \left\{ \frac{1}{Tr_j} - \frac{b_j}{r_j} \right\} \leq \frac{A_4}{T^2} + \frac{A_4}{YT}.$$

On the other hand, the minimum P_2 of the right endpoints of I_j is given by

$$P_2 = \min_j \left\{ \frac{\pi}{r_j} - \frac{1}{Tr_j} - \frac{b_j}{r_j} \right\} \geq \frac{\pi}{T} - \frac{A_4}{T^2} - \frac{A_4}{YT}.$$

To prove (5.3.8) it suffices to show that $P_1 < P_2$. Accordingly, it suffices to show that

$$\frac{A_4}{T^2} + \frac{A_4}{YT} < \frac{\pi}{T} - \frac{A_4}{T^2} - \frac{A_4}{YT}.$$

Now, it follows easily that the above inequality holds if we choose $Y > 3A_4/\pi$ (as in (5.3.3)) and T large enough. It follows that (5.3.8) holds, finishing the proof of Lemma 5.3.5. \square

The rest of the proof of the Theorem 1.3.3 for $n \equiv 3 \pmod{4}$ is the same as in the case $n \not\equiv 3 \pmod{4}$. The only difference is that in the analogue of (5.2.6) we get an extra T in the denominator due to the dependence on T in (5.3.6). However, this does not affect the final result since $T = O(\log \log \lambda)$ and $1/T^3$ is absorbed by the factor $-\delta$ in the power of $\log \lambda$ in (1.3.4). \square

5.4. Proof of Theorem 1.3.1 for $n \leq 5$. The rate of growth of the error term on the diagonal is determined by the exponents of the leading terms in Propositions 3.5.1 (for $\omega \neq \text{Id}$) and 3.5.3 (for $\omega = \text{Id}$). For $n \leq 4$ the leading terms in Proposition 3.5.1 are of order $\lambda^{(n-1)/2}$ and grow faster than then $\tilde{K}_{\lambda,T}(x)$. For $n = 5$, the powers of λ in the leading terms in Proposition 3.5.1 are of the same order as $\tilde{K}_{\lambda,T}(x)$, since $(n-1)/2 = n-3 = 2$ for $n = 5$. However, if we let $\lambda, T \rightarrow \infty$ then the sum of the contributions for $\omega \neq \text{Id}$ dominates due to the logarithmic improvement guaranteed by Theorem 4.1.2.

Note that the geodesic segments considered in section 4.1 in the case of Theorem 1.3.1 are geodesic loops. Consequently, the sum (4.1.1) should be taken over $\omega \in \Gamma \setminus \text{Id}$. We remark that the geodesic *loops*, and not the closed geodesics (though they are used in the proof), contribute to (4.1.1) and hence to the growth of $R_x(\lambda)$ (cf. section 1.4, [Bogom, p.175]).

The rest of the proof of Theorem 1.3.1 is identical to that of Theorem 1.3.3. In the case $n = 3$ the application of the Dirichlet principle is used appropriately as in section 5.3. \square

5.5. Proof of Theorem 1.3.1 for $n \geq 6$. The reason we get a different statement in Theorem 1.3.1 for $n \geq 6$ is that $\tilde{K}_{\lambda,T}(x)$ grows faster in λ than the sum of the leading terms for $\omega \neq \text{Id}$ given by Proposition 3.5.1. Indeed, $\tilde{K}_{\lambda,T}(x)$ is of order λ^{n-3} , while for $\omega \neq \text{Id}$ the leading terms are of order $\lambda^{(n-1)/2} < \lambda^{n-3}$ if $n \geq 6$.

Therefore, in this case the proof goes exactly as in the first part of Theorem 1.1.5. In particular, $T > 0$ is fixed. The only simplification compared to Theorem 1.1.5 is that on negatively curved manifolds $u_1(x, x) \neq 0$ at every point, since $u_1(x, x)$ is a multiple of the scalar curvature which is strictly negative. Therefore, in the notations of Theorem 1.1.5, $\kappa_x = 1$ for any x , and hence $R_x(\lambda) = \Omega(\lambda^{n-3})$. \square

Acknowledgements. The authors are deeply grateful to D. Dolgopyat for explaining them hyperbolic dynamics leading to the results in section 4. We would also like to thank V. Ivrii, M. Jakobson, V. Jaksic, Y. Kannai, Ya. Pesin, M. Pollicott, L. Polterovich, P. Sarnak, R. Sharp, A. Shnirelman, M. Shubin, J. Toth and D. Wise for useful discussions. This paper was completed while the first author visited IHES, and its hospitality is greatly appreciated.

6. APPENDIX: PROOF OF (4.1.4)

6.1. Refining the estimate. In this section we shall prove the following

Proposition 6.1.1. *Let $S_{x,y}(T)$ be as in (4.1.1). Then*

$$S_{x,y}(T) \geq C_0 e^{P(-\frac{T}{2}) \cdot T}$$

To prove Proposition 6.1.1, we group the terms in (4.7.1) according to conjugacy classes of $\omega \in \Gamma$. It is well-known that each conjugacy class corresponds to a unique closed geodesic on X . This correspondence can be defined as follows. The universal cover M of X can be tiled by identical copies of a fundamental domain Θ for X . Those copies are indexed by elements of $\Gamma = \pi_1(X)$. Indeed, pick a copy of $\Theta \in \mathcal{S}$ and call it Θ_e , where e is the identity element in Γ . Then for any other copy $\tilde{\Theta}$ there exists a unique $\omega \in \Gamma$ such that $\omega\Theta_e = \tilde{\Theta}$; we then let $\tilde{\Theta} = \Theta_\omega$. We thus have

$$\omega h^{-1} \Theta_h = \Theta_\omega, \quad \forall \omega, h \in \Gamma.$$

Given a *simple* closed geodesic γ on X , choose its lift to M passing through Θ_e (which we shall also call γ). Choose a basepoint $z \in \gamma \cap \Theta_e$, and let $\gamma = [z, \omega z], \omega \in \Gamma$; this defines ω . Now, γ intersects other copies of the fundamental domain besides Θ_e ; denote those copies by

$$(6.1.2) \quad \Theta_{h_1}, \Theta_{h_2}, \dots, \Theta_{h_n}$$

Here $h_1 = e, h_n = \omega$ and Θ_{h_i} -s are numbered consecutively along γ . One can show [Mil] that Γ is generated by $\{h : \partial\Theta_h \cap \partial\Theta_e \neq \emptyset\}$. Let $\{a_1, a_1^{-1}, \dots, a_k, a_k^{-1}\}$ denote a set of generators. Then each of the group elements $h_{i+1}h_i^{-1}, 1 \leq i \leq n-1$ is one of the generators (it maps Θ_{h_i} to $\Theta_{h_{i+1}}$). We set

$$a_{j_i} := h_{i+1}h_i^{-1}.$$

Thus to a geodesic γ we have associated an element ω and a word $w(\gamma)$ in a_j -s given by

$$(6.1.3) \quad a_{j_n} a_{j_{n-1}} \dots a_{j_2} a_{j_1} = \omega$$

It is clear that

$$(6.1.4) \quad n \geq l(\gamma) / \text{diam}(\Theta),$$

where $l(\gamma)$ is the length of the closed geodesic γ . In fact, it is shown in [Mil, Lemma 2] that (for a fixed set of generators) the ratio $l(\gamma) / L_\Gamma(w(\gamma))$, the denominator being the word length of $w(\gamma)$, is uniformly bounded above and below. However, we need only one side of this estimate, namely inequality (6.1.4).

If we choose a basepoint z on γ not in Θ_e but in adjacent fundamental domain Θ_{h_1} , then the corresponding $\omega \in \Gamma$ becomes $\tilde{\omega} = h_1 \omega h_1^{-1}$, and the word corresponding to $\tilde{\omega}$ is equal to (cf. (6.1.3))

$$a_{j_1} a_{j_n} a_{j_{n-1}} \dots a_{j_2} = a_{j_1} w(\gamma) a_{j_1}^{-1}$$

is a *cyclic shift* of $w(\gamma)$. Among all such cyclic shifts, choose one (call it $w_1(\gamma)$) which will have the *smallest* word length (such shift is called a *cyclically reduced* form of $w(\gamma)$). We shall prove the following

Proposition 6.1.5. *All cyclic shifts of a primitive cyclically reduced $w_1 = w_1(\gamma)$ are different elements of Γ .*

Proof of Proposition 6.1.5. It was shown by Preissman in [Pre] that any nontrivial commutative subgroup of Γ is infinite cyclic. Let us now assume for contradiction that two cyclic shifts of w_1 coincide. We can assume without loss of generality that one shift is trivial and equals w_1 itself. So suppose that

$$w_1 = u_1 u_2 = u_2 u_1$$

for two nontrivial words $u_1, u_2 \in \Gamma$. Then u_1 and u_2 generate a nontrivial commutative subgroup of Γ . Therefore, by Preissman's theorem it is cyclic and there exists $u_3 \in \Gamma$ such that $u_1 = u_3^k, u_2 = u_3^l$ for some $k, l \in \mathbf{Z}$. We claim that k and l have the same sign, otherwise the word $w_1 = u_1 u_2$ has cancellations and is not cyclically reduced. It follows that $w_1 = u_3^{k+l}$ and is therefore not primitive, which is a contradiction. This finishes the proof of the Proposition 6.1.5. \square

6.2. Proof of Proposition 6.1.1. We would like to get a better estimate for (4.1.1) than the one proved in section 4.7. To do that, consider a conjugacy class in Γ and the corresponding closed geodesic γ . We associate to γ a word $w(\gamma)$ of length $L_\Gamma(w(\gamma))$ as in section 6.1. By Proposition 6.1.5, to each primitive γ , there correspond $l_\Gamma(w(\gamma))$ cyclic shifts of this word, and by (6.1.4)

$$L_\Gamma(w(\gamma)) \geq \frac{l(\gamma)}{\text{diam}(\Theta)}.$$

It is now clear from the argument of section 4.7, that to each primitive closed geodesic γ we can associate at least $L_\Gamma(w(\gamma))$ segments connecting x to a point on the orbit Γy . The conclusion of Lemma 4.4.1 applies to each of these segments. Accordingly, the estimate (4.7.2) can be improved to

$$S_{x,y}(T) \geq \frac{C_3}{\text{diam}(\Theta)} \sum_{\omega \in \Gamma: 3D \leq l_{[\omega]} \leq T-2D} l_{[\omega]} \exp\left(-\frac{Z(l_{[\omega]}, \xi_{[\omega]})}{2}\right),$$

where $l_{[\omega]} = l(\gamma)$ is the length of the closed geodesic γ corresponding to the group element ω .

An application of (4.7.3) would complete the proof of Proposition 6.1.1, provided one can show that the sum over *primitive* closed geodesics is comparable to the "total" sum (4.7.3). Indeed, it easily follows that the contribution to (4.7.3) from *imprimitive* closed geodesics is

$$O\left(T e^{\frac{P(-\mathcal{H}/2)T}{2}}\right)$$

and grows much slower than $e^{P(-\mathcal{H}/2)T}$. Here we have used that every imprimitive geodesic corresponds to a primitive geodesic of at least twice smaller length. Therefore, the asymptotics in (4.7.3) still holds if we restrict summation to the primitive closed geodesics. \square

REFERENCES

- [A] D. Anosov. *Geodesic flows on closed Riemannian manifolds of negative curvature*. Trudy Mat. Inst. Steklov. 90 (1967).
- [ArAv] V. Arnold, A. Avez. *Ergodic problems of classical mechanics*, W.A. Benjamin, Inc., 1968.
- [Ber] P. Berard. *on the wave equation on a compact riemannian manifold without conjugate points*. Math. Z. 155 (1977), 249–276.
- [Bogom] E. Bogomolny. *Smoothed wavefunctions of chaotic quantum systems*. Physica D 31 (1988), 169–189.

- [Bow] R. Bowen *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*. Lecture Notes in Math. 470. Springer, 1975.
- [B-R] R. Bowen and D. Ruelle. *The ergodic theory of Axiom A flows*. Invent. Math. 29 (1975), no. 3, 181–202.
- [Ch] I. Chavel. *Riemannian geometry - a modern introduction*. Cambridge Tracts in Math. 108, Cambridge Univ. Press, 1995.
- [CdV] Y. Colin de Verdière. *Spectre conjoint d'opérateurs pseudo-différentiels qui commutent. II. Le cas intégrable*. Math. Z. 171 (1980), no. 1, 51–73.
- [Don] H. Donnelly. *On the wave equation asymptotics of a compact negatively curved surface*. Invent. Math. 45 (1978), 115–137.
- [D-G] J. Duistermaat and V. Guillemin. *The spectrum of positive elliptic operators and periodic bicharacteristics*. Inventiones Math. 29 (1975), 39–75.
- [E] P. Eberlein. *When is a geodesic flow of Anosov type? I, II*. Jour. Diff. Geometry 8 (1973), 437–463, 565–577.
- [G-S] I.M. Gelfand, G. E. Shilov, *Generalized functions*, vol. 1, Academic Press, New York, 1964.
- [Hej] D. Hejhal. *Selberg trace formula for $PSL(2, \mathbf{R})$, Vol. I*. Lecture Notes in Math. 548, Springer, 1976.
- [Hör] L. Hörmander. *The spectral function of an elliptic operator*. Acta Math. 121 (1968), 193–218.
- [Iv] V. Ivrii. *Precise spectral asymptotics for elliptic operators*. Lecture notes in math. 1100, Springer, 1984.
- [Kan] Y. Kannai. *Off diagonal short time asymptotics for fundamental solutions of diffusion equations*, Commun. Partial Differ. Equations 2 (1977), no. 8, 781–830.
- [H-K] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*. Encyclopedia of Math. and its Apps, 54. Cambridge University Press, Cambridge, 1995.
- [K] A. Karnaukh. *Spectral count on compact negatively curved surfaces*. Ph.D. thesis, Princeton University (1996), 1–48.
- [M-S] G. Margulis. *On Some Aspects of the Theory of Anosov Systems*, Springer, 2004. With a survey by Richard Sharp *Periodic Orbits of Hyperbolic Flows*.
- [Mil] J. Milnor. *A note on the curvature of the fundamental group*. Jour. Diff. Geom. 2 (1967), 1–7.
- [Par] W. Parry. *Equilibrium states and weighted uniform distribution of closed orbits*. Dynamical Systems (College Park, MD 1986–87), Lecture Notes in Math. 1342, 617–625. Springer, 1988.
- [P-P] W. Parry and M. Pollicott. *Zeta functions and closed orbit structure for hyperbolic systems*. Asterisque, 187–188 (1990), 1–256.
- [P-T] Y. Petridis and J. Toth. *The remainder in Weyl's law for Heisenberg manifolds*. J. Diff. Geom. 60 (2002), no. 3, 455–483.
- [P-R] R. Phillips and Z. Rudnick. *The circle problem in the hyperbolic plane*. J. Funct. Anal. 121 (1994), no. 1, 78–116.
- [Pre] A. Preissman. *Quelques propriétés globales des espaces de Riemann*. Comm. Math. Helv. 15 (1943), 175–216.
- [Ran] B. Randol. *A Dirichlet series of eigenvalue type with applications to asymptotic estimates*. Bull. London Math. Soc. 13 (1981), 309–315.
- [R-S] M. Rubinstein and P. Sarnak. *Chebyshev's bias*. Experiment. Math. 3 (1994), no. 3, 173–197.
- [Saf] Y. Safarov. *Asymptotics of a spectral function of a positive elliptic operator without a non-trapping condition*. Funct. Anal. Appl. 22 (1988) no. 3, 213–223.
- [S-V] Y. Safarov and D. Vassiliev. *The asymptotic distribution of eigenvalues of partial differential operators*. Translations of Mathematical Monographs, 155. AMS, 1997.
- [Schr] V. Schroeder. *Bounded geodesics in manifolds of negative curvature*. Math. Z. 235 (2000), no. 4, 817–828.
- [Sin1] Y. Sinai. *The asymptotic behavior of the number of closed geodesics*. AMS Transl. 73 (1968), 227–250.
- [Sin2] Y. Sinai. *Gibbs measures in ergodic theory*. Uspehi Mat. Nauk 27 (1972), 4(166), 21–64. Translation: Russian Math. Surveys 27 (1972), 21–69.
- [Shu] M. Shubin, *Pseudodifferential operators and spectral theory*, Springer-Verlag, 1987.
- [S-Z] C. Sogge and S. Zelditch. *Riemannian manifolds with maximal eigenfunction growth*. Duke Math. J. 114 (2002), no. 3, 387–437.

- [Sou] K. Soundararajan. *Omega bounds for the divisor and circle problems*, IMRN 36 (2003), 1987-1998.
- [Vol] A. Volovoy, *The Hamilton flow conditions associated with Weyl's conjecture*, Ann. Global Anal. Geom. vol 8, No. 2 (1990), 127-136.
- [Zel] S. Zelditch, *Lectures on wave invariants*, in: *Spectral theory and geometry*, edited by B. Davies and Y. Safarov, LMS lecture note series 273, Cambridge University Press, 1999.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE STR.
WEST, MONTRÉAL QC H3A 2K6, CANADA.
E-mail address: `jakobson@math.mcgill.ca`

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL CP 6128
SUCC CENTRE-VILLE, MONTRÉAL QC H3C 3J7, CANADA.
E-mail address: `iossif@dms.umontreal.ca`