

Multifractal Analysis via Scaling Zeta Functions and Recursive Structure of Lattice Strings

Rolando DE SANTIAGO, Michel L. LAPIDUS, Scott A. ROBY
and John A. ROCK



Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Juillet 2012

IHES/M/12/19

Multifractal Analysis via Scaling Zeta Functions and Recursive Structure of Lattice Strings

Rolando de Santiago, Michel L. Lapidus, Scott A. Roby, and John A. Rock

ABSTRACT. The multifractal structure underlying a self-similar measure stems directly from the weighted self-similar system (or weighted iterated function system) which is used to construct the measure. This follows much in the way that the dimension of a self-similar set, be it the Hausdorff, Minkowski, or similarity dimension, is determined by the scaling ratios of the corresponding self-similar system via Moran's theorem. The multifractal structure allows for our definition of scaling regularity and α -scaling zeta functions motivated by the geometric zeta functions of [24] and, in particular, the partition zeta functions of [9, 22]. Some of the results of this paper consolidate and partially extend the results of [9, 22] to a multifractal analysis of certain self-similar measures supported on compact subsets of a Euclidean space. Specifically, the α -scaling zeta functions generalize the partition zeta functions of [9, 22] when the choice of the family of partitions is given by the natural family of partitions determined by the self-similar system in question. Moreover, in certain cases, self-similar measures can be shown to exhibit lattice or nonlattice structure with respect to specified scaling regularity values in a sense which extends that of [24]. Additionally, in the context provided by generalized fractal strings viewed as measures, we define generalized self-similar strings, allowing for the examination of many of the results presented here in a specific overarching context and for a connection to the results regarding the corresponding complex dimensions as roots of Dirichlet polynomials in [24]. Furthermore, generalized lattice strings and recursive strings are defined and shown to be very closely related.

2010 *Mathematics Subject Classification.* Primary: 11M41, 28A12, 28A80. Secondary: 28A75, 28A78, 28C15, 33C05, 37B10, 37F35, 40A05, 40A10, 65Q30.

Key words and phrases. Fractal string, generalized self-similar string, recursive string, geometric zeta function, multifractal (or α -scaling) zeta function, scaling regularity, self-similar system, iterated function system, self-similar set, Besicovitch subset, self-similar measure, similarity dimension, Hausdorff dimension, Minkowski dimension, geometric multifractal spectrum, symbolic multifractal spectrum, scaling multifractal spectrum, complex dimensions, α -scaling complex dimensions, tapestry of complex dimensions, Cantor set, Cantor string, Fibonacci string, Lucas string, lattice vs. nonlattice, linear recurrence relation, hypergeometric series.

The work of the second author (M. L. Lapidus) was partially supported by the US National Science Foundation under the research grant DMS-1107750, as well as by the Institut des Hautes Etudes Scientifiques (IHES) where the second author was a visiting professor in the Spring of 2012 while this paper was written.

1. Introduction and summary

The abscissae of convergence of the α -scaling zeta functions associated with a self-similar measure provide a multifractal spectrum akin to the classic symbolic multifractal spectrum called the *scaling multifractal spectrum*. The technique described herein allows for partial yet extensive generalizations of the main results on the abscissae of convergence of partition zeta functions found in [22] and especially [9]. In a variety of cases, the Hausdorff dimensions of Besicovitch subsets of a self-similar set are recovered. In other cases, the classic symbolic multifractal spectrum of a self-similar measure is recovered. Moran's theorem is fully recovered in the special case where the measure in question (supported on a given self-similar set) is the natural mass distribution associated with a specific probability distribution.

Along with the development of α -scaling zeta functions, a generalized setting for self-similar and lattice generalized fractal strings is developed and *recursive strings* are introduced in this paper. Indeed, *generalized self-similar strings* provide a framework in which one can analyze certain special cases of α -scaling zeta functions. It is also shown that *generalized lattice strings* and *recursive strings* are intimately related.

In terms of applications, multifractal analysis is the study of a variety of mathematical, physical, dynamical, probabilistic, statistical, and biological phenomena from which families of fractals may arise. Such objects and behaviors are often modeled by mass distributions, or measures, with highly irregular and intricate structure. These multifractal measures, or simply multifractals, stem from phenomena such as weather, structure of lightning, turbulence, distribution of galaxies, spatial distribution of oil and minerals, cellular growth, internet trac, and nancial time series. See [5, 6, 9, 12–14, 25–27, 29–32, 34]

The structure of this paper is summarized as follows:

Section 2 provides a summary of results on classical approaches to multifractal analysis of self-similar measures established in the literature which are most pertinent to the results presented in this paper. In particular, the manner in which words are associated to the structure of self-similar measures (i.e., the way in which symbolic dynamics is employed) is discussed. See [3, 5–7, 10, 12–15, 27–32, 34] for classical and known results on self-similar sets and multifractal analysis of self-similar measures and other multifractal objects.

In Section 3, definitions and results regarding (generalized) fractal strings and complex dimensions from [24] are recalled. Further, *generalized self-similar strings* are defined and are shown to have, by design, complex dimensions that are completely determined by the roots of Dirichlet polynomials, as examined (thoroughly) in [24]. That is, generalized self-similar strings provide a context in which the self-similar structures considered throughout the paper can be related to the study of Dirichlet polynomials performed in [24]. In addition to [24], see [9, 17–19, 22, 25, 33] for further notions and uses of fractal strings and complex dimensions in a variety of contexts.

In Section 4, *generalized lattice strings* and *recursive strings* are defined and the intimate connections between them are discussed. In particular, it is shown that every generalized lattice string is a recursive string and exhibits properties which are determined by a naturally corresponding linear recurrence relation. It is also shown that the complex dimensions of a recursive string are the complex

dimensions of a naturally corresponding generalized lattice string. See [8] for a more detailed development of the connections between generalized lattice strings, linear recurrence relations, and recursive strings.

In Section 5, scaling regularity is used to define families of fractal strings associated with a given self-similar measure, giving rise to the definition of α -scaling zeta functions and the notion of multifractal spectrum as the abscissae of convergence of these zeta functions. This technique is motivated by and partially extends the results on partition zeta functions found in [9, 22, 24, 33].

Finally, Section 6 provides preliminary investigations of some further problems which expand upon the results of this paper. In particular, the α -scaling zeta functions of certain self-similar measures are shown to actually be hypergeometric series. This relationship is central to the material studied in [11]. See [2] for more information on hypergeometric series. Also, a family of self-similar measures which do not satisfy any of the conditions of the theorems and corollaries in Section 5 is investigated, motivating further research.

2. Multifractal analysis of self-similar systems

Multifractal analysis of a measure ν concerns the fractal geometry of objects such as the sets E_t of points $x \in E$ for which the measure $\nu(B(x, r))$ of the closed ball $B(x, r)$ with center x and radius r satisfies

$$\lim_{r \rightarrow 0^+} \frac{\log \nu(B(x, r))}{\log r} = t,$$

where $t \geq 0$ is the *local Hölder regularity* and E is the support of ν . Roughly speaking, multifractal analysis is the study of the ways in which a Borel measure behaves locally like r^t .

2.1. Multifractal spectra. The multifractal spectra of Definitions 2.1 and 2.12 along with Proposition 2.13 below are presented as found in [32], as well as the corresponding references therein. See especially the work of Cawley and Mauldin in [6].

DEFINITION 2.1. The *geometric Hausdorff multifractal spectrum* f_g of a Borel measure ν supported on E is given by

$$f_g(t) := \dim_H(E_t),$$

where $t \geq 0$, \dim_H is the Hausdorff dimension, and

$$E_t := \left\{ x \in E : \lim_{r \rightarrow 0^+} \frac{\log \nu(B(x, r))}{\log r} = t \right\}.$$

2.2. Self-similar systems. Self-similar systems lie at the heart of many of the results presented in this paper.

DEFINITION 2.2. Given $N \in \mathbb{N}$, $N \geq 2$, a *self-similar system* $\Phi = \{\Phi_j\}_{j=1}^N$ is a finite family of contracting similarities on a complete metric space (X, d_X) . Thus, for all $x, y \in X$ and each $j = 1, \dots, N$ we have

$$d_X(\Phi_j(x), \Phi_j(y)) = r_j d_X(x, y),$$

where $0 < r_j < 1$ is the *scaling ratio* (or Lipschitz constant) of Φ_j for each $j = 1, \dots, N$.

The *attractor* of Φ is the nonempty compact set $F \subset X$ defined as the unique fixed point of the contraction mapping

$$(1) \quad \Phi(\cdot) := \bigcup_{j=1}^N \Phi_j(\cdot)$$

on the space of compact subsets of X equipped with the Hausdorff metric. That is, $F = \Phi(F)$. The set F is also called the *self-similar set* associated with Φ .

A self-similar system (or set) is *lattice* if there is a unique real number r and positive integers k_j such that $0 < r < 1$ and $r_j = r^{k_j}$ for each $j = 1, \dots, N$. Otherwise, the self-similar system (or set) is *nonlattice*.

REMARK 2.3. For clarity of exposition, only self-similar systems on some Euclidean space \mathbb{R}^d ($d \in \mathbb{N}$), with $X \subset \mathbb{R}^d$, are considered.¹ Furthermore, throughout the paper we consider only self-similar systems which satisfy the *open set condition*. (See [6, 9, 12, 14, 24].) Recall that a self-similar system Φ satisfies the open set condition if there is a nonempty open set $V \subset \mathbb{R}^d$ such that $\Phi(V) \subset V$ and $\Phi_j(V) \cap \Phi_k(V) = \emptyset$ for each $j, k \in \{1, \dots, N\}$ where $j \neq k$. Results presented in [32] and [34], for example, specifically do *not* require the open set condition to be satisfied.

The notion of self-similar ordinary fractal strings which are either lattice or nonlattice as defined below follows from [24, Ch. 2]. These notions are extended and generalized in various ways throughout this paper.

DEFINITION 2.4. Let Φ be a self-similar system on \mathbb{R} such that $\sum_{j=1}^N r_j < 1$ and satisfying the open set condition on a compact interval I . Denote the endpoints of I by a_1 and a_2 , and assume that there are $j_1, j_2 \in \{1, \dots, N\}$ such that $a_1 \in \Phi_{j_1}(I)$ and $a_2 \in \Phi_{j_2}(I)$. The complement of the attractor F , given by $I \setminus F$, is a *self-similar ordinary fractal string*. Let I° denote the interior of I . The lengths of the connected components of $I^\circ \setminus \Phi(I)$, called the *gaps* of Φ , are denoted by $\mathbf{g} = (g_1, \dots, g_K) \in (0, \infty)^K$ where $K \in \mathbb{N}$. If, additionally, there is a unique real number r and positive integers k_j such that $0 < r < 1$ and $r_j = r^{k_j}$ for each $j = 1, \dots, N$, then $I \setminus F$ is *lattice*. Otherwise, $I \setminus F$ is *nonlattice*.

The following notation, which is motivated by the notation of self-similar (ordinary) fractal strings in [24, Ch. 2], allows for a clearer comparison between the main results herein and the classical results found in, for instance, [6, 12, 14].

NOTATION 2.5. For each $k \in \mathbb{N} \cup \{0\}$, let $\mathcal{J}_k = \{1, \dots, N\}^k$ denote the set of all finite sequences of length k in the symbols $\{1, \dots, N\}$ (i.e., words). For $k = 0$, let \mathcal{J}_0 be the set consisting of the empty word. Let $\mathcal{J} = \bigcup_{k=0}^{\infty} \mathcal{J}_k$; hence, \mathcal{J} is the set of all finite sequences (or words) in the symbols $\{1, \dots, N\}$. Let \mathcal{J}_∞ denote the set of all one-sided infinite sequences (or words) in the symbols $\{1, \dots, N\}$. For $J \in \mathcal{J}$, let $|J|$ denote number of components (i.e., the length) of J and define $|J| = \infty$ if $J \in \mathcal{J}_\infty$. For a word J (in either \mathcal{J} or \mathcal{J}_∞) and each $n \in \mathbb{N}$, $n \leq |J|$, let $J|n$ denote the truncation of J at its n th component and let $J|0$ denote the empty word. More specifically, $J|n = j_1 j_2 \dots j_n$ if J begins with the letters j_1, j_2, \dots, j_n . For $J \in \mathcal{J}$, define the contracting similarity Φ_J by

$$\Phi_J := \Phi_{j_{|J|}} \circ \Phi_{j_{|J|-1}} \circ \dots \circ \Phi_{j_1}.$$

¹Throughout this paper, \mathbb{N} denotes the set of positive integers: $\mathbb{N} = \{1, 2, 3, \dots\}$.

The scaling ratio of Φ_J is given by $r_J = \prod_{q=1}^{|J|} r_{\pi_q(J)}$, where $\pi_q(\cdot)$ is the projection of a word onto its q th component. For the empty word $J|0$, let $\Phi_{J|0}$ denote the identity map and let $r_{J|0} = 1$. For a set $E \subset X$ and a word J , let

$$E_J := \Phi_J(E).$$

Finally, define $\tau : \mathcal{J}_\infty \rightarrow \mathbb{R}^d$ by $\{\tau(J)\} := \bigcap_{n=0}^\infty E_{J|n}$.

In Theorem 9.1 of [12], for instance, it is shown via the Contraction Mapping Principle that (1) uniquely defines the attractor F as the fixed point of the map $\Phi(\cdot)$. Moreover, for any compact non-empty set E such that $\Phi_j(E) \subset E$ for each $j = 1, \dots, N$, we have

$$F = \bigcap_{n=0}^\infty \bigcup_{|J|=n} E_J,$$

where the notation “ $|J| = n$ ” indicates that, for each $n \in \mathbb{N} \cup \{0\}$, the corresponding union runs over all $J \in \mathcal{J}$ such that $|J| = n$.

For the support F , its Hausdorff dimension $\dim_H(F)$ (or, equivalently, its Minkowski dimension denoted by $\dim_M(F)$) is given by the solution of the Moran equation (2); see [12, 14, 28].

THEOREM 2.6 (Moran’s Theorem). *Let Φ be a self-similar system with scaling ratios $\{r_j\}_{j=1}^N$ that satisfies the open set condition. Then the Hausdorff (and Minkowski) dimension of the attractor F is given by the unique (and hence, positive) real solution D of the equation*

$$(2) \quad \sum_{j=1}^N r_j^\sigma = 1, \quad \sigma \in \mathbb{R}.$$

REMARK 2.7. The proof of Moran’s Theorem, as presented in [12] and [14] for instance, makes use of the mass distribution principle (see mass distribution principle 4.2 in [12]) and the “natural mass distribution” μ which is the self-similar measure determined, as described in the next section, by the self-similar system Φ and the probability vector $\mathbf{p} = (r_1^D, \dots, r_N^D)$ and which is supported on the attractor (or self-similar set) F .

2.3. Self-similar measures and scaling regularity. The multifractal measures in the context of this paper are constructed as follows (cf. [6, 9, 14]).

Let $\Phi = \{\Phi_j\}_{j=1}^N$ be a self-similar system that satisfies the open set condition with scaling ratios $\mathbf{r} = (r_1, \dots, r_N)$, where $0 < r_j < 1$ for each $j = 1, \dots, N$. Let $\mathbf{p} = (p_1, \dots, p_N)$ be a probability vector such that $0 \leq p_j \leq 1$ (hence $\sum_{j=1}^N p_j = 1$). A self-similar measure μ supported on the attractor F of the self-similar system Φ can be constructed via the probability vector \mathbf{p} and the equation

$$(3) \quad \mu(E) = \sum_{j=1}^N p_j \cdot \mu(\Phi_j^{-1}(E)),$$

which holds for all compact subsets E of \mathbb{R}^d . The self-similar measure μ is uniquely defined as the unique fixed point of the contraction implied by (3) on the space of regular Borel measures with unit total mass equipped with the L -metric (see [14]). In this setting, we refer to the pair (Φ, \mathbf{p}) as a *weighted self-similar system*. When

$\mathbf{p} = (r_1^D, \dots, r_N^D)$, the resulting self-similar measure μ is called the *natural Hausdorff measure* (or *natural mass distribution*) associated with the attractor F .

It is worth noting that the analysis of a given self-similar measure μ developed below depends only on the scaling ratios \mathbf{r} of the corresponding self-similar system Φ and the probability distribution determined by \mathbf{p} .

NOTATION 2.8. For each $J \in \mathcal{J}$, let

$$r_J = \prod_{q=1}^{|J|} r_{\pi_q(J)} \quad \text{and} \quad p_J = \prod_{q=1}^{|J|} p_{\pi_q(J)}.$$

Thus, for $J \in \mathcal{J}_\infty$ and each $k \in \mathbb{N}$ we have $r_{J|k} = r_{j_1} \cdots r_{j_k}$ and $p_{J|k} = p_{j_1} \cdots p_{j_k}$.

REMARK 2.9. There is a reason to distinguish between the elements of \mathcal{J}_∞ and those of \mathcal{J} . Specifically, the classical symbolic Hausdorff multifractal spectrum of Definition 2.12 below is defined in terms of the truncation of the elements of \mathcal{J}_∞ whereas our results, found in Section 5, are stated in terms of elements of \mathcal{J} .

DEFINITION 2.10. Let $J \in \mathcal{J}$. The *scaling regularity* of J is the value $A_{\mathbf{r}, \mathbf{p}}(J)$ given by

$$A_{\mathbf{r}, \mathbf{p}}(J) := \log_{r_J} p_J = \frac{\log p_J}{\log r_J},$$

where r_J and p_J are defined in Notation 2.8. Alternately, $A_{\mathbf{r}, \mathbf{p}}(J)$ is the unique real number α defined by $r_J^\alpha = p_J$.

REMARK 2.11. Note that the scaling regularity $A_{\mathbf{r}, \mathbf{p}}$ depends only on the scaling ratios \mathbf{r} of Φ and the probability vector \mathbf{p} but not on the contracting similarities $\Phi_j \in \Phi$. Indeed, the results presented in this paper are independent of the contracting similarities themselves. However, we do require, as mentioned above, that a self-similar system Φ satisfies the open set condition. Also note that for each $J \in \mathcal{J}$, the scaling regularity of J coincides with the coarse Hölder regularity of E_J ; see [6] and [9], for instance.

2.4. The symbolic Hausdorff multifractal spectrum. Self-similar measures are often called *multifractal measures* since, as will be discussed, whenever a self-similar measure is *not* the natural Hausdorff measure of the support, it decomposes the support into an amalgamation of fractal sets.

DEFINITION 2.12. Let μ be the self-similar measure determined by a weighted self-similar system (Φ, \mathbf{p}) . The *symbolic Hausdorff multifractal spectrum* f_s of μ is given by

$$f_s(t) := \dim_H \left\{ \tau(J) : J \in \mathcal{J}_\infty \text{ and } \lim_{n \rightarrow \infty} A_{\mathbf{r}, \mathbf{p}}(J|n) = t \right\}$$

for $t \geq 0$, where the map τ is defined at the very end of Notation 2.5. Here and henceforth, given $E \subset \mathbb{R}^d$, $\dim_H(E)$ denotes the Hausdorff dimension of E .

The following proposition is a simplified version of a similar proposition in [6].

PROPOSITION 2.13. *Let μ be the unique self-similar measure on \mathbb{R}^d defined by a weighted self-similar system (Φ, \mathbf{p}) which satisfies the open set condition. Then*

$$f_g(t) = f_s(t), \quad t \geq 0.$$

Due to Proposition 2.13 and the fact that scaling regularity plays a central role in Section 5, for a self-similar measure μ , focus is put on the symbolic Hausdorff multifractal spectrum f_s in the remainder of this section. A more complete development of the properties of f_s described here can be found in [6, §1].

REMARK 2.14. The self-similar measure μ uniquely defined by a weighted self-similar system (Φ, \mathbf{p}) attains maximum and minimum scaling regularity values α_{\min} and α_{\max} which, in turn, define the compact interval on which the sets E_t from Definition 2.1 are nonempty. Define $\alpha_j := \log_{r_j} p_j$ for each $j \in \{1, \dots, N\}$. These extreme scaling regularity values are given by

$$\alpha_{\min} = \min \{\alpha_j : j \in \{1, \dots, N\}\}, \quad \alpha_{\max} = \max \{\alpha_j : j \in \{1, \dots, N\}\}.$$

If $\mathbf{p} = (r_1^D, \dots, r_N^D)$, where D is the Hausdorff dimension of the attractor F of the self-similar system Φ (that is, if μ is the natural Hausdorff measure of its support, which is the attractor F), then $\alpha_{\min} = \alpha_{\max} = D$ and the domain of f_s is the singleton $\{D\}$. (In general, the domain of f_s is $[\alpha_{\min}, \alpha_{\max}]$.)

Otherwise, f_s is concave, f'_s near α_{\min} and α_{\max} is unbounded, the unique value t_1 such that $t_1 = f_s(t_1)$ is the *information dimension* of μ , and

$$\max\{f_s(t) : t \in [\alpha_{\min}, \alpha_{\max}]\} = \dim_H(F) = \dim_M(F),$$

where $\dim_M(F)$ denotes the Minkowski (or box) dimension of F ; see [12, Chs. 2 & 3] for the definition of Hausdorff and Minkowski dimension. In the context of an *ordinary fractal string* Ω , we are also concerned with the *inner* Minkowski dimension of its boundary $\partial\Omega$; see [24, §1.1] and Definition 3.7 below.

The following example is also studied in the context of partition zeta functions; see [22], [9, §5.2], and [33].

EXAMPLE 2.15 (Measures on the Cantor set). The Cantor set, denoted C , is the unique nonempty attractor of the lattice self-similar system Φ_C on $[0, 1]$ given by the two contracting similarities $\varphi_1(x) = x/3$ and $\varphi_2(x) = x/3 + 2/3$ with scaling ratios $\mathbf{r} = (1/3, 1/3)$. The Hausdorff dimension, and equivalently the Minkowski dimension, of C is the unique real-valued solution of the corresponding Moran equation (cf. (2)): $2 \cdot 3^{-s} = 1$. Thus, $\dim_H C = \dim_M C = \log_3 2 =: D_C$.

When Φ_C is weighted by $\mathbf{p} = (1/2, 1/2) = (1/3^{D_C}, 1/3^{D_C})$, the corresponding self-similar measure is the natural mass distribution (i.e., the natural Hausdorff measure) μ_C of the Cantor set. Such measures are used to find lower bounds on the Hausdorff dimension of their supports; see [12, Ch. 9] and Remark 5.14 below.

When Φ_C is weighted by $\mathbf{p} = (1/3, 2/3)$, we obtain the self-similar measure β , called the *binomial measure*, which exhibits the following properties: $\alpha_{\min} = 1 - \log_3 2$; $\alpha_{\max} = 1$; and for $t \in [1 - \log_3 2, 1]$, the geometric (and symbolic) Hausdorff multifractal spectrum is given by

$$f_g(t) = f_s(t) = - \left(\frac{1-t}{\log_3 2} \right) \log_3 \left(\frac{1-t}{\log_3 2} \right) - \left(1 - \frac{1-t}{\log_3 2} \right) \log_3 \left(1 - \frac{1-t}{\log_3 2} \right).$$

See [9, §5.2] and [33] for details.

Note that, in the case of μ_C , the only scaling regularity value attained by any corresponding word J is $A_{\mathbf{r}, \mathbf{p}}(J) = \log_3 2$. However, in the case of β , the attained scaling regularity values depend on the vector $\mathbf{k} = (k_1, k_2)$ where $\sum \mathbf{k} := k_1 + k_2 =$

$|J|$, k_1 denotes the number of times 1 appears in J , and k_2 denotes the number of times 2 appears in J . Moreover, for a fixed $n \in \mathbb{N}$ with $\sum \mathbf{k} = n$, we have

$$\#\{J \in \mathcal{J}_n : \#\{q : \pi_q(J) = j\} = k_j, j \in \{1, 2\}\} = \frac{n!}{k_1!k_2!} =: \binom{\sum \mathbf{k}}{\mathbf{k}},$$

where, in general, $\binom{\sum \mathbf{k}}{\mathbf{k}}$ denotes the multinomial coefficient and the projection π_q is defined in Notation 2.5.

This decomposition of the words associated with a given weighted self-similar system via scaling regularity along with the corresponding multinomial coefficients lies at the heart of the approach to multifractal analysis developed in Section 5 below.

The multifractal spectrum of a self-similar measure μ developed in Section 5 of this paper is determined by the abscissae of convergence of the α -scaling zeta functions. The motivation for this approach, and the analogous approach taken in [22] and [9], is a classic theorem of Besicovitch and Taylor and its significant extension which is at the heart of the theory of complex dimensions of fractal strings developed in [24]. (See Theorem 1.10 of [24] along with Theorem 3.4 below.)

2.5. Besicovitch subsets of the attractor of a self-similar system. A probability vector can be used not only to define a self-similar measure supported on a self-similar set, but also to decompose the support of such a measure into a family of disjoint subsets with interesting fractal properties of their own.

DEFINITION 2.16. Let F be the attractor of a self-similar system Φ and let $\mathbf{q} = (q_1, \dots, q_N)$ be a probability vector. The *Besicovitch subset* $F(\mathbf{q}) \subset F$ is defined as follows:

$$F(\mathbf{q}) := \left\{ x \in F : \lim_{k \rightarrow \infty} \frac{\#_j(x|_k)}{k} = q_j, j \in \{1, \dots, N\} \right\},$$

where $x|_k$ is the truncation of x at its k -th term in the expansion implied by Φ via $\tau(J) = x$ and $\#_j(x|_k)$ is the number of times the term j appears in $x|_k$ (really, the number of times j appears in $J|_k$).

REMARK 2.17. A little thought shows that, for a probability vector \mathbf{q} , the Besicovitch subset $F(\mathbf{q})$ is dense in the support F . That is, $\overline{F(\mathbf{q})} = F$, where $\overline{F(\mathbf{q})}$ is the closure of $F(\mathbf{q})$. Further, if the Minkowski dimension $\dim_M(F)$ exists (i.e., if the upper and lower Minkowski (or box) dimensions of F coincide), then Proposition 3.4 of [12] implies that $\dim_M(F(\mathbf{q})) = \dim_M(F)$. (Actually, the upper and lower Minkowski dimensions of a set are always equal, respectively, to those of its closure.) Throughout this paper, either the Minkowski or Hausdorff dimension of a given set will be considered, depending on the context.

Care needs to be taken in the above definition. Some $x \in F$ may have more than one $J \in \mathcal{J}_\infty$ where $x = \tau(J)$: however, this has no effect on the following proposition regarding the Hausdorff dimension of a Besicovitch subset $F(\mathbf{q})$. See [6] for a proof of the following proposition, and see [3] and [10] for related classical results.

PROPOSITION 2.18. *Let $F(\mathbf{q})$ be the Besicovitch subset of the attractor F of a self-similar system Φ with scaling ratios $\mathbf{r} = (r_1, \dots, r_N)$ determined by a probability*

vector \mathbf{q} . Then

$$\dim_H(F(\mathbf{q})) = \frac{\sum_{j=1}^N q_j \log q_j}{\sum_{j=1}^N q_j \log r_j}.$$

3. Fractal strings and complex dimensions

The material found in this section provides a brief summary of pertinent results from the theory of complex dimensions of fractal strings developed by Lapidus and van Frankenhuysen in [24].

3.1. Generalized fractal strings. For a (local) measure η on $(0, \infty)$, denote the total variation of η by $|\eta|$. For a bounded measurable set S we have,²

$$|\eta|(S) = \sup \left\{ \sum_{k=1}^m |\eta(S_k)| \right\},$$

where $m \in \mathbb{N}$ and $\{S_k\}_{k=1}^m$ ranges over all finite partitions of S into disjoint measurable subsets of $(0, \infty)$. Recall that $|\eta| = \eta$ if η is positive and that $|\eta|$ is a positive measure.

DEFINITION 3.1. A *generalized fractal string* is either a local complex or a local positive measure η on $(0, \infty)$ which is supported on a subset of (x_0, ∞) for some $x_0 > 0$. The *dimension* of η , denoted D_η , is the *abscissa of convergence* of the Dirichlet integral $\zeta_{|\eta|}(\sigma) = \int_0^\infty x^{-\sigma} |\eta|(dx)$. That is,

$$D_\eta := \inf \left\{ \sigma \in \mathbb{R} \mid \int_0^\infty x^{-\sigma} |\eta|(dx) < \infty \right\}.$$

The *geometric zeta function* of η is the Mellin transform of η given by

$$\zeta_\eta(s) = \int_0^\infty x^{-s} \eta(dx),$$

for $\operatorname{Re}(s) > D_\eta$. Let $W \subset \mathbb{C}$ be a window³ on an open neighborhood of which ζ_η has a meromorphic extension. By a mild abuse of notation, both the geometric zeta function of η and its meromorphic extension are denoted by ζ_η . The set of (*visible*) *complex dimensions* of η , denoted by \mathcal{D}_η , is given by

$$\mathcal{D}_\eta = \{\omega \in W : \zeta_\eta \text{ has a pole at } \omega\}.$$

²A *local measure* η on $(0, \infty)$ is a \mathbb{C} -valued function on the Borel σ -algebra of $(0, \infty)$ whose restriction to any bounded subinterval is a complex measure. If η is $[0, \infty]$ -valued, then η is simply a locally bounded positive measure on $(0, \infty)$ and it is called a *local positive measure*.

³As in [24], we are interested in the meromorphic extension of the geometric zeta function $\zeta_\mathcal{L}$ to suitable regions. To this end, consider the *screen* S as the contour

$$S : S(t) + it \quad (t \in \mathbb{R}),$$

where $S(t)$ is a Lipschitz continuous function $S : \mathbb{R} \rightarrow [-\infty, D_\mathcal{L}]$. Also, consider the *window* W as the closed set

$$W = \{s \in \mathbb{C} : \operatorname{Re}(s) \geq S(\operatorname{Im}(s))\}$$

and assume that ζ_η has a meromorphic continuation to an open connected neighborhood of W satisfying suitable polynomial growth conditions (as in [24, §5.3]).

In the case where ζ_η has a meromorphic extension to $W = \mathbb{C}$, the set \mathcal{D}_η is referred to as the *complex dimensions* of η .

Generalized fractal strings have two notable predecessors: fractal strings and ordinary fractal strings.

DEFINITION 3.2. A *fractal string* $\mathcal{L} = \{\ell_j\}_{j=1}^\infty$ is a nondecreasing sequence of positive real numbers which tend to zero. Hence, $\lim_{j \rightarrow \infty} \ell_j = 0$.

REMARK 3.3. As in [18], but unlike the classic geometric setting of [24], we do not require $\sum_{j=1}^\infty \ell_j < \infty$. That is, in [24] an *ordinary fractal string* is a bounded open subset Ω of \mathbb{R} and \mathcal{L} denotes the sequence of lengths of the disjoint open intervals whose union is Ω ; see, e.g., [20], [24, Chs. 1 & 2], and [18]. We note, however, that in [21], [24, Chs. 3 & 10], and [18, 19], for example, the underlying sequence of scales is allowed to satisfy $\sum_{j=1}^\infty \ell_j = \infty$.

There is a natural relationship between discrete generalized fractal strings η and fractal strings \mathcal{L} . Recall that the Dirac mass at $x \in (0, \infty)$, denoted by $\delta_{\{x\}}$, is the measure given by

$$\delta_{\{x\}}(S) := \begin{cases} 1, & x \in S, \\ 0, & x \notin S, \end{cases}$$

where $S \subseteq \mathbb{R}$ (for example). So, a fractal string

$$\mathcal{L} = \{\ell_j\}_{j=1}^\infty = \{l_n \mid l_n \text{ has multiplicity } m_n, n \in \mathbb{N}\}$$

defines the generalized fractal string η as follows:

$$\eta = \sum_{j=1}^\infty \delta_{\{\ell_j^{-1}\}} = \sum_{n=1}^\infty m_n \delta_{\{l_n^{-1}\}}.$$

For such η , it immediately follows that $\zeta_{\mathcal{L}} = \zeta_\eta$ and $\mathcal{D}_{\mathcal{L}} = \mathcal{D}_\eta$.

The following theorem, which is Theorem 1.10 of [24], is a restatement of a classical theorem of Besicovitch and Taylor (see [4]) formulated in terms of ordinary fractal strings. (A direct proof can be found in [24], *loc. cit.*) For the definition of (inner) Minkowski dimension as used below, see [24, §1.1] and Definition 3.7.

THEOREM 3.4. *Suppose Ω is an ordinary fractal string with infinitely many lengths denoted by \mathcal{L} . Then the abscissa of convergence of $\zeta_{\mathcal{L}}$ coincides with the (inner) Minkowski dimension of $\partial\Omega$. That is, $D_{\mathcal{L}} = \dim_M(\partial\Omega)$, where $\dim_M(\partial\Omega)$ denotes the (inner) Minkowski dimension of $\partial\Omega$.⁴*

In terms of such meromorphic extensions, a given geometric zeta function encountered throughout this paper falls into one of two categories: (i) the geometric zeta function has a meromorphic extension to all of \mathbb{C} , or (ii) the geometric zeta function is similar to a hypergeometric series and, hence, in general, the appropriate extension is yet to be determined (see Section 6 and [2, 11]). For a self-similar ordinary fractal string (see Definition 2.4), the geometric zeta function has a closed form which allows for a meromorphic extension to all of \mathbb{C} . This closed form is given in the following theorem, which is Theorem 2.3 in [24]. Compare this theorem with Theorem 3.12 below, which is a completely analogous result regarding the closed form of the geometric zeta function of a *generalized self-similar string*.

⁴Unlike in [4], the fact that the inner Minkowski Dimension is used in [24] allows for not requiring any additional assumptions about $\partial\Omega$ or about Ω .

THEOREM 3.5. *Let Ω be a self-similar ordinary fractal string with lengths \mathcal{L} . Then the geometric zeta function $\zeta_{\mathcal{L}}$ has a meromorphic continuation to the whole complex plane, given by*

$$(4) \quad \zeta_{\mathcal{L}}(s) = \frac{L^s \sum_{k=1}^K g_k^s}{1 - \sum_{j=1}^N r_j^s}, \quad s \in \mathbb{C}.$$

Here, $L = \zeta_{\mathcal{L}}(1)$ is the total length of Ω .

EXAMPLE 3.6 (The Fibonacci string). Consider the lattice self-similar system on the interval $[0, 4]$ given by $\Phi_1(x) = x/2$ and $\Phi_2(x) = x/4 + 3$ (i.e., $r_1 = 1/2$ and $r_2 = 1/4 = 1/2^2$). This self-similar system generates an attractor F and a lattice ordinary fractal string $\Omega = [0, 4] \setminus F$ whose lengths are given by the Fibonacci string \mathcal{L}_{Fib} . See [24, §2.3.2]. The Fibonacci string \mathcal{L}_{Fib} is the fractal string with distinct lengths given by $l_n = 2^{-n}$ occurring with multiplicity $m_n = F_n$, where F_n is the n th Fibonacci number and $n \in \mathbb{N} \cup \{0\}$. (Hence, $F_0 = F_1 = 1, F_2 = 2, \dots$) Via Theorem 3.5, the geometric zeta function of \mathcal{L}_{Fib} is given by

$$(5) \quad \zeta_{\mathcal{L}_{\text{Fib}}}(s) := \zeta_{\mathcal{L}_{\text{Fib}}}(s) = \sum_{n=0}^{\infty} F_n 2^{-ns} = \frac{1}{1 - 2^{-s} - 4^{-s}}$$

for $s \in \mathbb{C}$, and the dimension D_{Fib} is the unique real-valued solution of the equation

$$(6) \quad 2^{-2s} + 2^{-s} = 1, \quad s \in \mathbb{C}.$$

Moreover, the complex dimensions of \mathcal{L}_{Fib} are the complex roots of (6).⁵ Thus, we have

$$(7) \quad \mathcal{D}_{\text{Fib}} = \{D_{\text{Fib}} + izp : z \in \mathbb{Z}\} \cup \{-D_{\text{Fib}} + i(z + 1/2)p : z \in \mathbb{Z}\},$$

where $\varphi = (1 + \sqrt{5})/2$ is the Golden Ratio, $D_{\text{Fib}} = \log_2 \varphi$, and the oscillatory period is $p = 2\pi/\log 2$.

Accordingly, the ordinary self-similar fractal string $[0, 4] \setminus F$ is lattice in the sense of Definition 2.4. Furthermore, as they are given by the Fibonacci numbers, the multiplicities $m_n = F_n$ satisfy the linear recurrence relation for $n \geq 2$ given by

$$(8) \quad F_n = F_{n-1} + F_{n-2},$$

with initial conditions $\mathbf{s}_{\text{Fib}} := (F_0, F_1) = (1, 1)$.

Connections between generalized lattice strings and linear recurrence relations are examined in Section 4.

3.2. Minkowski measurability and lattice/nonlattice dichotomy. The following theorem is a partial restatement of Theorem 8.15 of [24] that provides a criterion for the Minkowski measurability of an ordinary fractal string Ω satisfying certain mild restrictions. First, we introduce a few useful tools. (See [24, §1.1] for more information such as detailed definitions of lower and upper Minkowski contents for ordinary fractal strings.)

DEFINITION 3.7. Let $V(\varepsilon)$ be the 1-dimensional Lebesgue measure of the *inner tubular neighborhood* of Ω given by the set $\{x \in \Omega : d(x, \partial\Omega) < \varepsilon\}$. The *inner Minkowski dimension* of $\partial\Omega$, denoted $D = D_{\mathcal{L}}$, is given by

$$D = \inf\{t \geq 0 : V(\varepsilon) = O(\varepsilon^t) \text{ as } \varepsilon \rightarrow 0^+\}.$$

⁵These roots are obtained by solving the quadratic equation $z^2 + z - 1 = 0$ with $z = 2^{-s}, s \in \mathbb{C}$.

The boundary $\partial\Omega$ of an ordinary fractal string Ω is *Minkowski measurable* if the limit $\lim_{\varepsilon \rightarrow 0^+} V(\varepsilon)\varepsilon^{D-1}$ exists in $(0, \infty)$.

In the following theorem, the equivalence of statements (2) and (3) below for an arbitrary ordinary fractal string is established in [20] (without any conditions on \mathcal{L} other than $D_{\mathcal{L}} \neq 0, 1$).

THEOREM 3.8 (Criterion for Minkowski measurability). *Let Ω be an ordinary fractal string whose geometric zeta function $\zeta_{\mathcal{L}}$ has a meromorphic extension which satisfies certain mild growth conditions.⁶ Then the following statements are equivalent:*

- (i) D is the only complex dimension with real part D , and it is simple.
- (ii) $\partial\Omega$ is Minkowski measurable.
- (iii) $\ell_j = Lj^{-1/D}(1 + o(1))$ as $j \rightarrow \infty$, for some $L > 0$.

Theorem 3.8 applies to all lattice self-similar ordinary fractal strings. Specifically, if Ω is a lattice string, then the mild growth conditions are satisfied by $\zeta_{\mathcal{L}}$ and there are no complex dimensions other than D which have real part D , so the boundary of Ω is not Minkowski measurable. On the other hand, Theorem 3.8 does not apply to all nonlattice self-similar strings since there are some for which $\zeta_{\mathcal{L}}$ does not satisfy the growth conditions for a screen of the type described in footnote 6; see [24, Example 5.32]. Nonetheless, we have the following theorem which partially summarizes Theorems 8.23 and 8.36 of [24].

THEOREM 3.9 (Lattice/nonlattice dichotomy). *A self-similar ordinary fractal string Ω is nonlattice if and only if its boundary $\partial\Omega$ is Minkowski measurable.*

REMARK 3.10. An extension of a part of Theorem 3.9 for suitable classes of self-similar tilings (and sets or systems) of higher-dimensional Euclidean spaces is provided in [19], using results of [24, Ch. 8] and [18]. (See also the relevant references therein.) Furthermore, an interesting study of a nonlinear analogue of Theorem 3.9 (and related counter-examples for certain self-conformal sets) in the real line is conducted in [16].

In the next section, we summarize (in an extended self-similar setting) some key results on the generalized fractal strings (i.e., fractal strings viewed as measures) of Lapidus and van Frankenhuysen found in [24, Ch. 4].

3.3. Generalized self-similar strings. In this section, a particular form of generalized fractal string, called *generalized self-similar string*, is defined. The corresponding complex dimensions, by design, are given by the roots of a naturally associated Dirichlet polynomial equation; see [24, Ch. 3]. The geometric zeta function of a generalized self-similar string has a meromorphic continuation established in Theorem 3.12 which allows for the determination of the complex dimensions of the recursive strings defined in Section 4 and the complex dimensions of certain scaling zeta functions in Section 5.

DEFINITION 3.11. A discrete generalized fractal string η is *self-similar*, and may be referred to as a *generalized self-similar string*, if there are $K, N \in \mathbb{N}$ such

⁶Specifically, Theorem 3.8 holds if $\zeta_{\mathcal{L}}$ is *languid* (see [24, Def. 5.2]) for a screen passing between the vertical line $\operatorname{Re}(s) = D_{\mathcal{L}}$ and all the complex dimensions of (the corresponding fractal string) \mathcal{L} with real part strictly less than D , and not passing through 0.

that for some $\mathbf{g} = (g_1, \dots, g_K) \in (0, \infty)^K$, $\mathbf{r} = (r_1, \dots, r_N) \in (0, 1)^N$, and $\mathbf{m} = (m_1, \dots, m_N) \in \mathbb{C}^N$, we have

$$\eta = \sum_{k=1}^K \sum_{J \in \mathcal{J}} m_J \delta_{\{g_k^{-1} r_J^{-1}\}},$$

where $r_J = \prod_{q=1}^{|J|} r_{\pi_q(J)}$ and m_J is defined in an identical fashion. For the empty word $J|0$, we let $m_{J|0} = r_{J|0} = 1$. We refer to components of the vectors \mathbf{g} , \mathbf{r} , and \mathbf{m} (or sometimes the vectors themselves) as the *gaps*, *initial scaling ratios*, and *initial multiplicities* of η , respectively.

The following theorem, which is an immediate consequence of the results of [24, §3.3], but will be useful in the sequel, determines a closed form of the geometric zeta function of a generalized self-similar string (cf. Theorem 3.5 above and Theorem 2.3 and Equation (3.21) in [24]). This closed form allows for the meromorphic continuation of the geometric zeta function to all of the complex plane and, therefore, an extension of the theory of complex dimensions for self-similar strings of [24]. When the multiplicities m_j are all integral and positive, this fact, along with the detailed study (conducted in [24, Ch. 3]) of the periodic or almost periodic distribution of the complex dimensions in the lattice or nonlattice case, respectively, was used in an essential manner in the work of [18] on tube formulas for higher-dimensional self-similar sets and tilings. (See also the earlier papers by the first two authors of [18] quoted therein, along with [19].) Moreover, the proof of the following theorem is included so that one may compare and contrast with the results on α -scaling functions presented in Section 5 below, especially the decomposition of the corresponding multiplicities.

THEOREM 3.12. *Let η be a generalized self-similar string. Then the geometric zeta function of η has a meromorphic continuation to \mathbb{C} given by*

$$\zeta_\eta(s) = \frac{\sum_{k=1}^K g_k^s}{1 - \sum_{j=1}^N m_j r_j^s}, \quad \text{for } s \in \mathbb{C}.$$

PROOF. For $q = 0$, $J|0$ is the empty word and we have $m_{J|0} = r_{J|0} = 1$. For each $q \in \mathbb{N}$ we have

$$\sum_{J \in \mathcal{J}: |J|=q} m_J r_J^s = \sum_{\nu_1=1}^N \cdots \sum_{\nu_q=1}^N m_{\nu_1} r_{\nu_1}^s \cdots m_{\nu_q} r_{\nu_q}^s = \left(\sum_{j=1}^N m_j r_j^s \right)^q.$$

Now, ζ_η is given by

$$\zeta_\eta(s) = \int_0^\infty x^{-s} \eta(dx) = \sum_{k=1}^K \sum_{J \in \mathcal{J}} m_J (g_k r_J)^s = \sum_{k=1}^K g_k^s \left(\sum_{q=0}^\infty \left(\sum_{j=1}^N m_j r_j^s \right)^q \right).$$

If $\text{Re}(s) > D_r$, the series converges since $|\sum_{j=1}^N m_j r_j^s| < 1$. Moreover, we have

$$\zeta_\eta(s) = \frac{\sum_{k=1}^K g_k^s}{1 - \sum_{j=1}^N m_j r_j^s}.$$

Hence, by the Principle of Analytic Continuation, ζ_η has a meromorphic continuation to all of \mathbb{C} given by the last formula. \square

EXAMPLE 3.13 (Self-similar ordinary fractal strings). The geometric zeta function of a self-similar ordinary fractal string is given by the geometric zeta function of an appropriately defined generalized self-similar string. Basically, an ordinary fractal string is self-similar if it is the complement of a self-similar set with respect to a certain type of closed interval. (See [24, Ch. 2].) For instance, the geometric zeta function ζ_{Fib} of the Fibonacci string given in (5) is the geometric zeta function of the generalized self-similar string determined by $K = g_1 = 1$, $\mathbf{r} = (1/2, 1/4)$, and $\mathbf{m} = (1, 1)$. (See Example 3.6.) Another such example is the Cantor string Ω_{CS} which is the complement in $[0, 1]$ of the classic Cantor set. (See [9] and [24] for further information on the Cantor string.)

The self-similar systems which generate the Fibonacci string and the Cantor string, respectively, are lattice. Hence, as described in Section 4, the corresponding generalized self-similar strings are *recursive strings* (see Definition 4.9). We note that in the case of the Fibonacci string, this fact is foreshadowed in the linear recurrence relation (8).

EXAMPLE 3.14 (Generalized Cantor strings). A *generalized Cantor string* μ is a generalized self-similar string of the form

$$\mu = \sum_{n=0}^{\infty} b^n \delta_{\{r^{-n}\}},$$

where $0 < r < 1$ and $b > 0$. (Note that b is not required to be an integer.) That is, as a generalized self-similar string, μ is determined by $N = K = g_1 = 1$, $\mathbf{r} = (r)$, and $\mathbf{m} = (b)$. Generalized Cantor strings are studied in [24, Chs. 8 & 10], where it is shown (among many other things) that the geometric zeta function of a generalized Cantor string has a meromorphic extension to all of \mathbb{C} given by

$$(9) \quad \zeta_{\mu}(s) = \frac{1}{1 - b \cdot r^s}, \quad s \in \mathbb{C}.$$

In the special case where $b = 2^{-1}$ and $r = 3^{-1}$, one obtains a generalized Cantor string which can not be realized geometrically as an ordinary fractal string (since $b = m_1 = 2^{-1}$ is nonintegral). The geometric zeta function of μ , after meromorphic extension to \mathbb{C} , is given by

$$\zeta_{\mu}(s) = \frac{1}{1 - 2^{-1} \cdot 3^{-s}}, \quad s \in \mathbb{C}.$$

Note that the “dimension” $D_{\mu} = -\log_3 2$ is *negative*.⁷ Indeed, by allowing the “multiplicities” $\mathbf{m} \neq \mathbf{0}$ to comprise complex numbers in Definition 3.11, one is able to study self-similar structures with respect to measures which do not necessarily (or rather, do not readily) correspond to geometric objects.

This section concludes with an indication of how the framework of generalized self-similar strings ties to material elsewhere in the literature. Specifically, the *scaling measure* and *scaling zeta function* of a self-similar system, as defined below, are studied in [18] and [24]. In the setting of this paper, a scaling measure is a generalized self-similar string where $m_j = K = g_1 = 1$ for each $j = 1, \dots, N$, and the scaling zeta function is the corresponding geometric zeta function. As mentioned in Section 3.2 above, there are deep connections between the Minkowski measurability

⁷By “dimension” here we mean the abscissa of convergence of ζ_{μ} , as in Definition 3.1.

of the attractor of a self-similar system and the structure of the complex dimensions of corresponding zeta functions such as the scaling zeta function.

DEFINITION 3.15. For a self-similar system Φ , the *scaling measure* is the associated generalized fractal string given by

$$\eta_{\Phi} := \sum_{J \in \mathcal{J}} \delta_{\{r_J^{-1}\}}.$$

The *scaling zeta function* of Φ is given by

$$\zeta_{\Phi}(s) := \frac{1}{1 - \sum_{j=1}^N r_j^s}, \quad s \in \mathbb{C}.$$

According to Definition 3.1, it is just the geometric zeta function (i.e., the Mellin transform) of the generalized fractal string η_{Φ} .

REMARK 3.16. Every scaling measure η_{Φ} defines a generalized self-similar string and if the self-similar system Φ is lattice, then η_{Φ} is a generalized lattice string (see Definition 4.1 below).

4. Generalized lattice strings and linear recurrence relations

In this section we discuss yet another notion of lattice structure. This time, it pertains to generalized self-similar strings. In particular, the results of this section extend accordingly to self-similar sets which are subsets of some Euclidean space (not just the real line) and certain cases of scaling zeta functions of self-similar measures as discussed in Section 5.

Linear recurrence relations are also shown, in this section, to be intimately related to the *generalized lattice strings* defined here.

DEFINITION 4.1. A generalized self-similar string η is *lattice* if there is a unique $0 < r < 1$ called the *multiplicative generator* of η and positive integers k_j such that $r_j = r^{k_j}$ for each $j = 1, \dots, N$. A lattice generalized self-similar string may also be referred to as a *generalized lattice string*.

REMARK 4.2. Every lattice ordinary fractal string (see Definition 2.4 and the Fibonacci and Cantor strings in Examples 3.6 and 3.13) and every generalized Cantor string (see Example 3.14) can be realized as a generalized lattice string.

What follows is a discussion of a connection between linear recurrence relations and the (possibly complex) multiplicities stemming from generalized lattice strings and lattice ordinary fractal strings.

4.1. Linear recurrence relations. A brief summary of some relevant material on linear recurrence relations is provided in this section. For a more detailed introduction to recurrence relations, see [1].

DEFINITION 4.3. A sequence $\{s_n\}_{n=0}^{\infty} \subset \mathbb{C}$ satisfies a *linear recurrence relation* \mathcal{R} if there exist $d \in \mathbb{N}$ and $\mathbf{a} \in \mathbb{C}^d$ with $\pi_d(\mathbf{a}) \neq 0$ such that for all $n \geq d$, we have

$$s_n = a_1 s_{n-1} + \dots + a_d s_{n-d}.$$

The positive integer d is the *degree of linear recursion*. The constant vector $\mathbf{a} := (a_1, \dots, a_d)$ is the *kernel* of \mathcal{R} . The *characteristic equation* of \mathcal{R} is

$$(10) \quad \varphi^d = a_1 \varphi^{d-1} + \dots + a_d, \quad \varphi \in \mathbb{C}.$$

For a given sequence $\{s_n\}_{n=0}^{\infty}$ which satisfies the linear recurrence relation \mathcal{R} , the first d terms of the sequence $\{s_n\}_{n=0}^{\infty}$ are the *initial conditions*, and they are denoted by the vector $\mathbf{s} := (s_0, \dots, s_{d-1})$.

REMARK 4.4. Each linear recurrence relation \mathcal{R} is completely determined by its kernel \mathbf{a} . Also, each sequence $\{s_n\}_{n=0}^{\infty}$ which satisfies a linear recurrence relation \mathcal{R} is completely determined by the corresponding kernel \mathbf{a} and the initial conditions $\mathbf{s} := (s_0, \dots, s_{d-1})$.

Some of the properties of linear recurrence relations can be understood and analyzed in the context of linear algebra. A brief synopsis is provided below; see [1] and [8] for more information.

DEFINITION 4.5. Suppose $\{s_n\}_{n=1}^{\infty}$ satisfies a linear recurrence relation \mathcal{R} . The kernel matrix A and the n th sequence matrix S_n are respectively given by

$$(11) \quad A := \begin{pmatrix} a_1 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{d-1} & 0 & \cdots & 1 \\ a_d & 0 & \cdots & 0 \end{pmatrix}, \quad S_n := \begin{pmatrix} s_{2d-2+n} & \cdots & s_{d-1+n} \\ s_{2d-3+n} & \cdots & s_{d-2+n} \\ \vdots & \ddots & \vdots \\ s_{d-1+n} & \cdots & s_n \end{pmatrix}.$$

THEOREM 4.6. Suppose $\{s_n\}_{n=0}^{\infty}$ satisfies a linear recurrence relation \mathcal{R} . Then $S_0 A^n = S_n$.

PROOF. For each $j \in \mathbb{N}$ we have $S_j A = S_{j+1}$. Thus, for each $n \in \mathbb{N}$ we have

$$(12) \quad S_0 A^n = S_1 A^{n-1} = \cdots = S_n. \quad \square$$

PROPOSITION 4.7. A complex number λ is an eigenvalue of the kernel matrix A of a recurrence relation \mathcal{R} if and only if λ is a solution of the characteristic equation of \mathcal{R} given by (10).

The proof is omitted as it follows immediately from the definitions and some linear algebra.

REMARK 4.8. Recursion relations in the context of measures with another type of self-similarity property are studied in [24, §4.4.1]. There, such measures are allowed to have mass near zero and are assumed to be absolutely continuous with respect to dx/x , the Haar measure on the multiplicative group \mathbb{R}_+^* . These results are compared and contrasted with the results of this paper in [8].

In the next section, linear recurrence relations extend and are related to generalized lattice strings.

4.2. Recursive strings. In this section we present a new type of generalized fractal string, called a *recursive string*. Recursive strings are closely related to generalized lattice strings as described in this section, especially via Theorems 4.11 and 4.13. See [8] for a more detailed analysis of recursive strings, linear recurrence relations, and connections to generalized lattice strings.

DEFINITION 4.9. Given $K \in \mathbb{N}$, $0 < r < 1$, $\mathbf{g} = (g_1, \dots, g_K) \in (0, 1)^K$, and a sequence $\{s_n\}_{n=0}^{\infty} \subset \mathbb{C}$ satisfying a linear recurrence relation \mathcal{R} , the *recursive string*

$\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ is the discrete generalized fractal string given by

$$\eta_{\mathcal{R}}(\mathbf{s}; \cdot) = \sum_{k=1}^K \sum_{n=0}^{\infty} s_n \delta_{\{g_k^{-1} r^{-n}\}}(\cdot).$$

The real number r is called the *multiplicative generator* of $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ and the components of the vector \mathbf{g} (or sometimes \mathbf{g} itself) are(is) called the *gaps* of $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$.

The geometric zeta function, dimension, and complex dimensions of a recursive string $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ are denoted, respectively, by

$$\zeta_{\mathcal{R}}(\mathbf{s}; \cdot) := \zeta_{\eta_{\mathcal{R}}(\mathbf{s}; \cdot)}, \quad D_{\mathcal{R}}(\mathbf{s}) := D_{\eta_{\mathcal{R}}(\mathbf{s}; \cdot)}, \quad \text{and} \quad \mathcal{D}_{\mathcal{R}}(\mathbf{s}) := \mathcal{D}_{\eta_{\mathcal{R}}(\mathbf{s}; \cdot)}.$$

REMARK 4.10. Since every linear recurrence relation \mathcal{R} is completely determined by its kernel \mathbf{a} and every sequence $\{s_n\}_{n=0}^{\infty} \subset \mathbb{C}$ which satisfies \mathcal{R} is further determined by the initial conditions \mathbf{s} , we have that every recursive string $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ is completely determined by the kernel \mathbf{a} , initial conditions \mathbf{s} , gaps \mathbf{g} , and multiplicative generator r .

THEOREM 4.11. *Every generalized lattice string η is a recursive string. That is,*

$$\eta = \sum_{n=0}^{\infty} s_n \delta_{\{r^{-n}\}},$$

where $0 < r < 1$ and the sequence of multiplicities $\{s_n\}_{n=0}^{\infty} \subset \mathbb{C}$ satisfies a linear recurrence relation \mathcal{R} .

PROOF. Since η is lattice, there exist a self-similar system Φ with scaling ratios $\mathbf{r} = (r_1, \dots, r_N)$ and a unique $0 < r < 1$ such that $r_j = r^{k_j}$ for some positive integer k_j and each $j = 1, \dots, N$. Let $k = \max\{k_j : j \in \{1, \dots, N\}\}$. For each $q = 1, \dots, k$, define a_q to be the sum of the (complex) multiplicities of r^q with respect to the self-similar system. That is, $a_q := \sum_{j: r_j = r^q} m_j$. Let $s_0 = 1$ and for $n \in \mathbb{N}$, let s_n be the multiplicity of the length r^n . Since $r^n = r^q r^{n-q}$ for $1 \leq q \leq k$, each instance of r^q will contribute s_{n-q} to the total multiplicity of the length r^n . Specifically, for all $n \geq k$,

$$s_n = \sum_{q=1}^k a_q s_{n-q} = a_1 s_{n-1} + \dots + a_k s_{n-k}.$$

This is the desired linear recurrence relation \mathcal{R} . □

The following corollary provides a well-known fact regarding the Hausdorff dimension and Minkowski dimension of a lattice self-similar system. A simple proof is provided in part to illuminate the deep connections between generalized lattice strings and linear recurrence relations.

COROLLARY 4.12. *Suppose Φ is a lattice self-similar system with attractor F . Then there is a sequence of positive integer multiplicities $\{s_n\}_{n=0}^{\infty}$ which satisfies the linear recurrence relation \mathcal{R} corresponding to Φ such that the scaling measure η_{Φ} satisfies*

$$\eta_{\Phi} = \sum_{n=0}^{\infty} s_n \delta_{\{r^{-n}\}}.$$

Moreover, the Hausdorff dimension and Minkowski dimension of F are given by

$$\dim_H(F) = \dim_M(F) = -\log_r \varphi,$$

where φ is the unique positive root the of characteristic equation of \mathcal{R} .

PROOF. By Theorem 2.6, $\dim_H(F)$ and $\dim_M(F)$ are given by the unique real-valued solution D of (2). Since Φ is lattice, $r_j = r^{k_j}$, for $0 < r < 1$ and $k_j \in \mathbb{N}$. Letting $m = \max\{k_j : j = 1, \dots, N\}$, a_j be the multiplicity of r^j , and $x = r^{-s}$, (2) becomes

$$1 = \sum_{j=0}^m a_j r^{js} = a_1 x^{-1} + \dots + a_m x^{-m}.$$

Multiplying through by x^m yields

$$x^m = a_1 x^{m-1} + \dots + a_m,$$

which is a polynomial in x . Since the coefficients are all positive, Descartes's Rule of Signs implies the existence of a unique positive root φ . Thus, $\varphi = r^{-D}$, which implies $D = -\log_r \varphi$. \square

The following theorem and its corollaries show that a recursive string is nearly a generalized lattice string. In particular, the set of complex dimensions of a recursive string is contained in or equal to the set of complex dimensions of a naturally related generalized lattice string.

THEOREM 4.13. *Let \mathcal{R} be a linear recurrence relation of degree d . Let $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ be the recursive string determined by the kernel \mathbf{a} of \mathcal{R} , initial conditions \mathbf{s} , gaps \mathbf{g} , and multiplicative generator $0 < r < 1$. Then*

$$(13) \quad \zeta_{\mathcal{R}}(\mathbf{s}; s) = g(s) \cdot \frac{\sum_{n=0}^{d-1} s_n r^{ns} - \sum_{l=1}^{d-1} a_l r^{ls} \sum_{n=0}^{d-1-l} s_n r^{ns}}{1 - \sum_{j=1}^d a_j r^{js}}$$

for $s \in \mathbb{C}$ and $g(s) := \sum_{k=1}^K g_k^s$.

PROOF. Assume, for notational simplicity, that $K = g_1 = 1$. Further, for $\operatorname{Re}(s) > D_{\mathcal{R}}(\mathbf{s}; \cdot)$, consider the sum

$$\zeta_{\mathcal{R}}(\mathbf{s}; s) - \sum_{n=0}^{d-1} s_n r^{ns} = \sum_{n=0}^{\infty} s_{n+d} r^{(n+d)s}.$$

We then have successively:

$$\begin{aligned} \zeta_{\mathcal{R}}(\mathbf{s}; s) - \sum_{n=0}^{d-1} s_n r^{ns} &= a_1 \sum_{n=0}^{\infty} s_{n+d-1} r^{(n+d)s} + \dots + a_d \sum_{n=0}^{\infty} s_n r^{(n+d)s} \\ &= a_1 r^s \sum_{n=0}^{\infty} s_{n+d-1} r^{(n+d-1)s} + \dots + a_d r^d \sum_{n=0}^{\infty} s_n r^{ns} \\ &= a_1 r^s \left(\zeta_{\mathcal{R}}(\mathbf{s}; s) - \sum_{n=0}^{d-2} s_n r^{ns} \right) + \dots + a_d r^{ds} \zeta_{\mathcal{R}}(\mathbf{s}; s). \end{aligned}$$

So,

$$\zeta_{\mathcal{R}}(\mathbf{s}; s) \left(1 - \sum_{j=1}^d a_j r^{js} \right) = \sum_{n=0}^{d-1} s_n r^{ns} - \sum_{l=0}^{d-1} a_l r^{ls} \sum_{n=0}^{d-1-l} s_n r^{ns}.$$

Therefore, for $\operatorname{Re}(s) > D_{\mathcal{R}}(\mathbf{s}; \cdot)$, we have

$$(14) \quad \zeta_{\mathcal{R}}(\mathbf{s}; s) = \frac{\sum_{n=0}^{d-1} s_n r^{ns} - \sum_{l=1}^{d-1} a_l r^{ls} \sum_{n=0}^{d-1-l} s_n r^{ns}}{1 - \sum_{j=1}^d a_j r^{js}}.$$

Since the right-hand side of (14) defines a meromorphic function in all of \mathbb{C} , it follows that $\zeta_{\mathcal{R}}(\mathbf{s}; \cdot)$ is meromorphic in \mathbb{C} and is still given by the same expression on \mathbb{C} .

Finally, for the case where $K \neq 1$ or $g_1 \neq 1$, the reasoning is exactly the same as above except for the fact that the factor $g(s) = \sum_{k=1}^K g_k^s$ would need to be included throughout the proof, accordingly. \square

As we have seen, the right-hand side of (13) allows for a meromorphic continuation of $\zeta_{\mathcal{R}}(\mathbf{s}; \cdot)$ to \mathbb{C} . This, in turn, allows for the following two results.

COROLLARY 4.14. *Let \mathcal{R} be a linear recurrence relation and let $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ be a recursive string as in Theorem 4.13. Then there is a generalized lattice string η and an entire function $h_{\mathcal{R}}(\mathbf{s}; s)$ such that, after meromorphic continuation, for all $s \in \mathbb{C}$, we have*

$$(15) \quad \zeta_{\mathcal{R}}(\mathbf{s}; s) = h_{\mathcal{R}}(\mathbf{s}; s) \zeta_{\eta}(s).$$

Moreover, $D_{\mathcal{R}}(\mathbf{s}) = D_{\eta}$ and $\mathcal{D}_{\mathcal{R}}(\mathbf{s}) \subset \mathcal{D}_{\eta}$.

PROOF. For some $d \in \mathbb{N}$, let \mathcal{R} be a linear recurrence relation with kernel $\mathbf{a} = (a_1, \dots, a_d)$. Given a recursive string $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$ determined by the gaps $\mathbf{g} = (g_1, \dots, g_K)$, initial conditions $\mathbf{s} = (s_0, \dots, s_{d-1})$, and multiplicative generator r , consider the generalized lattice string η' determined by the initial multiplicities $\mathbf{m}' = \mathbf{a}$, initial scaling ratios $\mathbf{r}' = (r, r^2, \dots, r^d)$ and a single gap of length 1 (i.e., $\mathbf{g}' = (1)$). We immediately have that $D_{\mathcal{R}}(\mathbf{s}) = D_{\eta'}$.

Now, define $h_{\mathcal{R}}(\mathbf{s}; s)$ by

$$h_{\mathcal{R}}(\mathbf{s}; s) := g(s) \cdot \left(\sum_{n=0}^{d-1} s_n r^{ns} - \sum_{l=1}^{d-1} a_l r^{ls} \sum_{n=0}^{d-1-l} s_n r^{ns} \right),$$

where $g(s) = \sum_{k=1}^K g_k^s$. Then $h_{\mathcal{R}}(\mathbf{s}; \cdot)$ is entire, and by applying Theorems 3.12 and 4.13 to η' and $\eta_{\mathcal{R}}(\mathbf{s}; \cdot)$, respectively, we see that (15) holds with $\eta := \eta'$. Moreover, after meromorphic continuation, (15) holds for all $s \in \mathbb{C}$, and we conclude that $\mathcal{D}_{\mathcal{R}}(\mathbf{s}) \subset \mathcal{D}_{\eta'}$. \square

Note that in Corollary 4.14, we do not conclude that $\mathcal{D}_{\mathcal{R}}(\mathbf{s}) = \mathcal{D}_{\eta'}$ in general. This is due to the fact that some of the zeros of $h_{\mathcal{R}}(\mathbf{s}; \cdot)$ might cancel some of the poles (of the meromorphic extension) of $\zeta_{\eta} = \zeta_{\eta'}$.

The following corollary is an immediate consequence of the combination of Theorems 3.8 and 4.13.

COROLLARY 4.15. *Let Ω be an ordinary fractal string with lengths*

$$\mathcal{L}_{\mathcal{R}} = \{r^n : r^n \text{ has multiplicity } m_n, n \in \mathbb{N} \cup \{0\}\},$$

where $0 < r < 1$ and the multiplicities $\{m_n\}$ satisfy a linear recurrence relation \mathcal{R} . Furthermore, if none of the complex roots of $h_{\mathcal{R}}(\mathbf{s}; s) = 0$ is also a complex root of the Moran equation $1 = \sum_{j=1}^d a_j r^{j s}$, then $\partial\Omega$ is not Minkowski measurable.

The discussion of recursive strings concludes with an example of a recursive string which is not a generalized self-similar string (hence, it is also not a generalized lattice string).

EXAMPLE 4.16 (The Lucas string). Consider the recursive string η_{Luc} , called the *Lucas string*, determined by the kernel $\mathbf{a} = (1, 1)$, initial conditions $\mathbf{s}_{\text{Luc}} := (2, 1)$, a single gap determined by $K = g_1 = 1$, and multiplicative generator $r = 1/2$. The geometric zeta function of η_{Luc} satisfies

$$\zeta_{\text{Luc}}(s) = \sum_{n=0}^{\infty} s_n 2^{-ns},$$

where $\text{Re}(s)$ is large enough and the sequence $\{s_n\}_{n=0}^{\infty}$ satisfies the linear recurrence relation determined by \mathbf{a} with initial conditions \mathbf{s}_{Luc} . That is, $s_0 = 2, s_1 = 1$, and for $n \geq 2$, s_n satisfies the Fibonacci recursion relation (8). The closed form of ζ_{Luc} given by Theorem 4.13 is

$$\zeta_{\text{Luc}}(s) = \frac{2 - 2^{-s}}{1 - 2^{-s} - 4^{-s}}, \quad s \in \mathbb{C}.$$

Note that the sequence $\{s_n\}_{n=1}^{\infty} = \{1, 3, 4, 7, \dots\}$ (where the index n begins at 1) is the sequence of *Lucas numbers*. Moreover, the Fibonacci numbers $\{F_n\}_{n=0}^{\infty}$ and the sequence $\{s_n\}_{n=0}^{\infty}$ both satisfy the linear recurrence relation determined by the kernel $\mathbf{a} = (1, 1)$, but with initial conditions $\mathbf{s}_{\text{Fib}} := (1, 1)$ and $\mathbf{s}_{\text{Luc}} = (2, 1)$, respectively. (See Example 3.6.) Also, note that $2 - 2^{-s} = 0$ if and only if $2^{-s} = 2$, which implies $1 - 2^{-s} - 4^{-s} = -5 \neq 0$. Thus, the set of complex dimensions of the Lucas string, denoted \mathcal{D}_{Luc} , coincides with \mathcal{D}_{Fib} , the set of complex dimensions of the Fibonacci string given by (7) and which are the roots of the Dirichlet polynomial equation $2^{-s} + 4^{-s} = 1$. Also note that $D_{\text{Luc}} = D_{\text{Fib}} = \log_2 \varphi$, where $\varphi = (1 + \sqrt{5})/2$ is the Golden Ratio.

Moreover, $\{s_n\}_{n=0}^{\infty}$ is a sequence of positive integers and $\zeta_{\text{Luc}}(1)$ is positive and finite. Hence, ζ_{Luc} is the geometric zeta function of a suitably defined ordinary fractal string Ω_{Luc} . However, such a fractal string is *not* lattice, in the sense that it cannot be realized as the complement of the attractor of a (lattice or even nonlattice) self-similar system in some closed interval (see Definition 2.4). Indeed, ζ_{Luc} can not be put into the form given in (4). Hence, Theorem 3.9 does not apply. However, Corollary 4.15 applies and, hence, $\partial\Omega_{\text{Luc}}$ is not Minkowski measurable.

EXAMPLE 4.17 (Recursive structure of generalized Cantor strings). Every generalized Cantor string μ is a recursive string; see Example 3.14 and [24, Chs. 8 & 10]. Indeed, we have

$$\mu = \sum_{n=0}^{\infty} s_n \delta_{\{r^{-n}\}},$$

where the sequence $\{s_n\}_{n=0}^{\infty}$ is given by $s_n = b^n$ for every n and satisfies the linear recurrence relation \mathcal{R} determined by the kernel $\mathbf{a} = (b)$ with initial condition $\mathbf{s} = (1)$. By Theorem 4.13, the meromorphic extension of the geometric zeta function ζ_μ given in (9) is recovered as $\zeta_{\mathcal{R}}(\mathbf{s}; \cdot)$. That is,

$$\zeta_\mu(s) = \zeta_{\mathcal{R}}(\mathbf{s}; s) = \frac{1}{1 - b \cdot r^{-s}}, \quad s \in \mathbb{C}.$$

The technique developed in the next section is motivated by the theory of complex dimensions of fractal strings and is designed to perform a multifractal analysis of self-similar measures.

5. Multifractal analysis via scaling regularity and scaling zeta functions

In this section, an approach to multifractal analysis is described in which, for a given weighted self-similar system (Φ, \mathbf{p}) , a family of fractal strings is defined by gathering lengths (or rather, scales) according to scaling regularity values α . Then, a family of α -scaling zeta functions is readily defined and the collection of their abscissae of convergence provides a multifractal spectrum of dimensions, called the *scaling multifractal spectrum*, which is akin to the geometric and symbolic multifractal spectra (see Section 2.1).

The approach given in this section generalizes the results regarding *partition zeta functions* found in [9, §5]. Specifically, the measures studied in [9, §5] are limited to measures supported on subsets of $[0, 1]$, whereas the framework provided below allows for results on self-similar measures which are supported on self-similar sets in Euclidean spaces of any dimension. See Section 2 as well as [14].

We refer to Section 13.3 (of the second edition) of [24] for a survey of aspects of the theory of multifractal zeta functions (and related partition zeta functions), as developed in [17] and [22]. We also refer to [9] and [33] where additional results can be found.

5.1. α -scales and α -scaling zeta functions. Throughout this section, and indeed throughout the paper, only weighted self-similar systems (Φ, \mathbf{p}) with scaling ratios $\mathbf{r} = \{r_j\}_{j=1}^N$ satisfying the open set condition are considered; see Section 2.

DEFINITION 5.1. Let (Φ, \mathbf{p}) be a weighted self-similar system. For a scaling regularity value $\alpha \in \mathbb{R}$, the *sequence of α -scales*, denoted $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$, is the fractal string given by

$$\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha) = \{r_J : J \in \mathcal{J}, A_{\mathbf{r}, \mathbf{p}}(J) = \alpha\},$$

where $A_{\mathbf{r}, \mathbf{p}}$ is defined in Definition 2.10.

Alternately, the *distinct α -scales* are denoted by $l_n(\alpha)$ and the corresponding *multiplicities* are denoted by $m_n(\alpha)$. Thus,

$$m_n(\alpha) := \#\{J \in \mathcal{J} : r_J = l_n(\alpha), A_{\mathbf{r}, \mathbf{p}}(J) = \alpha\},$$

and we can consider the sequence of α -scales to be given by

$$\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha) = \{l_n(\alpha) : l_n(\alpha) \text{ has multiplicity } m_n(\alpha)\}.$$

Given that $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ is a fractal string when there are infinitely many words $J \in \mathcal{J}$ such that $A_{\mathbf{r}, \mathbf{p}}(J) = \alpha$ for some $\alpha \in \mathbb{R}$, one defines a zeta function and a set of complex dimensions for each such α as follows.

DEFINITION 5.2. Consider $\alpha \in \mathbb{R}$ such that $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ is not empty. The α -scaling zeta function $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha; \cdot)$ is the geometric zeta function of the sequence of α -scales $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$. That is,

$$(16) \quad \zeta_{\mathbf{r}, \mathbf{p}}(\alpha; s) := \zeta_{\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)}(s) = \sum_{A_{\mathbf{r}, \mathbf{p}}(J)=\alpha} r_J^s,$$

where $\operatorname{Re}(s) > D_{\mathbf{r}, \mathbf{p}}(\alpha) := D_{\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)}$. If $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ is empty, we set $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha; s) = 0$. The scaling multifractal spectrum $f_{\mathbf{r}, \mathbf{p}}(\alpha)$ is the function given by the abscissa of convergence of $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha; s)$ for each $\alpha \in \mathbb{R}$. That is,

$$f_{\mathbf{r}, \mathbf{p}}(\alpha) := \max\{0, \inf\{\sigma \in \mathbb{R} : \zeta_{\mathbf{r}, \mathbf{p}}(\alpha; \sigma) < \infty\}\},$$

for $\alpha \in \mathbb{R}$.

More precisely, $f_{\mathbf{r}, \mathbf{p}}(\alpha)$ is the maximum of 0 and the abscissa of convergence of the Dirichlet series which defines the α -scaling zeta function $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha; \cdot)$. Thus, $f_{\mathbf{r}, \mathbf{p}}(\alpha) \geq 0$ whenever $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ comprises an infinite number of scales and $f_{\mathbf{r}, \mathbf{p}}(\alpha) = 0$ otherwise. Hence, α is *nontrivial* if $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ comprises an infinite number of scales, otherwise α is *trivial* (see Remark 5.3 and compare Remark 4.5 of [9]). Accordingly, for a nontrivial scaling regularity value α , the set given by

$$\{s \in \mathbb{C} : \operatorname{Re}(s) > f_{\mathbf{r}, \mathbf{p}}(\alpha)\}$$

is the largest open right half-plane on which the Dirichlet series in (16) is absolutely convergent.

REMARK 5.3. Unlike in Remark 4.5 of [9], all trivial scaling regularity values α_0 in the current setting generate an empty sequence of α_0 -scales. Thus, for any $\alpha \in \mathbb{R}$, $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ is either countably infinite or empty. See Remark 5.7 for clarification.

DEFINITION 5.4. Let $W_\alpha \subset \mathbb{C}$ be a window on a connected open neighborhood of which $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha; \cdot)$ has a meromorphic extension. (Again, both the geometric zeta function of $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ and its meromorphic extension are denoted by $\zeta_{\mathbf{r}, \mathbf{p}}(\alpha, \cdot)$.) The set of (*visible*) α -scaling complex dimensions, denoted $\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\alpha)$, is the set of (visible) complex dimensions of the sequence of α -scales $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$ given by

$$\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\alpha) = \{\omega \in W_\alpha : \zeta_{\mathcal{L}} \text{ has a pole at } \omega\}.$$

The *tapestry of complex dimensions* $\mathcal{T}_{\mathbf{r}, \mathbf{p}}$ with respect to the regions W_α is the subset of $\mathbb{R} \times \mathbb{C}$ given by

$$\mathcal{T}_{\mathbf{r}, \mathbf{p}} := \{(\alpha, \omega) : \alpha \text{ is nontrivial, } \omega \in \mathcal{D}_{\mathbf{r}, \mathbf{p}}(\alpha)\}.$$

5.2. Scaling regularity values attained by self-similar measures. The collection of all scaling regularity values $A_{\mathbf{r}, \mathbf{p}}(J)$ attained by the words $J \in \mathcal{J}$ with respect to a given weighted self-similar system (Φ, \mathbf{p}) are found readily. The following notation is used in order to facilitate the statement of the results. Note that, as throughout the paper, only weighted self-similar systems which satisfy the open set condition are considered.

NOTATION 5.5. Let $N \in \mathbb{N}$ with $N \geq 2$. For a pair of N -vectors \mathbf{k} and \mathbf{r} with $\mathbf{r} \in (0, 1)^N$ and $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$, let $\mathbf{r}^{\mathbf{k}} := r_1^{k_1} \cdots r_N^{k_N}$. Furthermore, denote by $\operatorname{gcd}(\mathbf{k})$ the greatest common divisor of the nonzero components of \mathbf{k} . Let $\sum \mathbf{k} := \sum_{j=1}^N k_j$ and

$$(17) \quad \binom{\sum \mathbf{k}}{\mathbf{k}} := \binom{K}{k_1 \dots k_N} = \frac{K!}{k_1! \cdots k_N!},$$

where $K = \sum \mathbf{k}$.

Recall from Notation 2.5 that \mathcal{J} denotes the collections of all finite words on the alphabet $\{1, \dots, N\}$.

LEMMA 5.6. *Let (Φ, \mathbf{p}) be a weighted self-similar system as above and let $J \in \mathcal{J}$ be a nonempty word. Then there exist a unique vector $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$ with $\gcd(\mathbf{k}) = 1$ and a unique positive integer n such that $|J| = n \sum \mathbf{k}$ and*

$$(18) \quad A_{\mathbf{r}, \mathbf{p}}(J) = \alpha(\mathbf{k}) := \log_{\mathbf{r}^{\mathbf{k}}} \mathbf{p}^{\mathbf{k}}.$$

Moreover, for each $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$ we have

$$\#\{J \in \mathcal{J}_{\sum \mathbf{k}} : \#\{q : \pi_q(J) = j\} = k_j, j \in \{1, \dots, N\}\} = \binom{\sum \mathbf{k}}{\mathbf{k}},$$

where $\binom{\sum \mathbf{k}}{\mathbf{k}}$ is given by (17).

PROOF. The results follow immediately from basic combinatorics and the definitions of r_J and p_J given in Notation 2.8. \square

REMARK 5.7. An immediate consequence of Lemma 5.6 is that each $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$ satisfying $\gcd(\mathbf{k}) = 1$ yields a countably infinite collection of words $J \in \mathcal{J}$ which have scaling regularity $A_{\mathbf{r}, \mathbf{p}}(J) = \alpha(\mathbf{k})$. The scales r_J associated with these words essentially constitute the terms of the Dirichlet series which defines the α -scaling zeta function (for $\alpha = \alpha(\mathbf{k})$, as in (18)) in Definition 5.2.

Also, to clarify Remark 5.3, suppose $\alpha_0 \in \mathbb{R}$. If there is a word $J \in \mathcal{J}$ such that $A_{\mathbf{r}, \mathbf{p}}(J) = \alpha_0$, then by Lemma 5.6, α_0 is nontrivial. Indeed, in that case, for each $n \in \mathbb{N}$, there is a word $H_n \in \mathcal{J}$ such that $|H_n| = n|J|$ and $A_{\mathbf{r}, \mathbf{p}}(H_n) = A_{\mathbf{r}, \mathbf{p}}(J) = \alpha_0$.

Furthermore, for each $n \in \mathbb{N}$ we have

$$\#\{J \in \mathcal{J}_{n \sum \mathbf{k}} : \#\{q : \pi_q(J) = j\} = nk_j, j \in \{1, \dots, N\}\} = \binom{n \sum \mathbf{k}}{n \mathbf{k}}.$$

If there are $\mathbf{m}_1, \mathbf{m}_2 \in (\mathbb{N} \cup \{0\})^N$ such that $\gcd(\mathbf{m}_1) = \gcd(\mathbf{m}_2) = 1$, $\mathbf{m}_1 \neq \mathbf{m}_2$, and $\alpha(\mathbf{m}_1) = \alpha(\mathbf{m}_2)$, then the total multiplicity of a given scale r' is given by the sum of all multinomial coefficients $\binom{n \sum \mathbf{k}}{n \mathbf{k}}$ where $r' = \mathbf{r}^{n \mathbf{k}}$. The determination of a closed formula for such total multiplicity is, in general, a difficult problem. However, various special cases are addressed in the remainder of this section and in Section 6. In particular, see Theorems 5.8 and 5.11, along with Corollary 5.12.

5.3. Self-similar and lattice structures within self-similar measures.

For a weighted self-similar system (Φ, \mathbf{p}) as above, let $\alpha_j := \log_{r_j} p_j$ for each $j = 1, \dots, N$ denote the scaling regularity values attained with respect to (Φ, \mathbf{p}) by all words J such that $|J| = 1$. Let N_0 denote the number of distinct values among the α_j . Then $N_0 \leq N$ and we denote these distinct α_j values by β_q , where $q = 1, \dots, N_0$.

THEOREM 5.8. *Let (Φ, \mathbf{p}) be a weighted self-similar system as above. Suppose the collection of distinct scaling ratios $\{\beta_q : q = 1, \dots, N_0\}$ is rationally independent. Then, for each $q \in \{1, \dots, N_0\}$, $\zeta_{\mathbf{r}, \mathbf{p}}(\beta_q; \cdot)$ has a meromorphic continuation*

to all of \mathbb{C} given by

$$(19) \quad \zeta_{\mathbf{r}, \mathbf{p}}(\beta_q; s) = \frac{\sum_{j: \alpha_j = \beta_q} r_j^s}{1 - \sum_{j: \alpha_j = \beta_q} r_j^s}, \quad \text{for } s \in \mathbb{C}.$$

PROOF. Since the β_q values are rationally independent, the only words J with regularity β_q are those with components $\pi_n(J) = j$ where $\alpha_j = \beta_q$. Thus, for each $q \in \{1, \dots, N_0\}$, the β_q -scaling zeta function is equal to the scaling zeta function of a self-similar system Φ_q whose scaling ratios are given by $\{r_j : \alpha_j = \beta_q\}$ less the first term which corresponds to the empty word. So, by Theorem 3.12 we have

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta_q; s) = \zeta_{\Phi_q}(s) - 1 = \frac{\sum_{j: \alpha_j = \beta_q} r_j^s}{1 - \sum_{j: \alpha_j = \beta_q} r_j^s}.$$

□

COROLLARY 5.9. *If the conditions of Theorem 5.8 are satisfied and, additionally, if there exists a unique r such that $0 < r < 1$ and for each j where $\alpha_j = \beta_q$ we have $r_j = r^{u_j}$ for some $u_j \in \mathbb{N}$, then there is a generalized lattice string η such that*

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta_q; s) = h(s)\zeta_\eta(s),$$

where $h(s) := \sum_{j: \alpha_j = \beta_q} r^{u_j s}$ and $\eta := \eta_\Phi$ (as in Definition 3.15).

REMARK 5.10. For each $q \in \{1, \dots, N_0\}$, the function $h(s) := \sum_{j: \alpha_j = \beta_q} r_j^s$ in the numerator of the right-hand side of (19) is entire. Moreover, the complex roots of $h(s) = 0$ are distinct from the poles of ζ_{Φ_q} , so $\mathcal{D}(\beta_q) = \mathcal{D}_{\Phi_q}$. Also, note that the hypotheses of Corollary 5.9 do not require the self-similar system Φ to be lattice. Section 6.2 examines a self-similar measure built upon a nonlattice self-similar system which satisfies the hypotheses of Corollary 5.9 for a particular scaling regularity value.

The α -scaling zeta functions with respect to regularity values α which do not fit any of the conditions required in the results presented in this section are much harder to determine in general; see Section 6.2. However, given particular constraints, the full family of α -scaling zeta functions for certain weighted self-similar systems can be determined, as we shall see in the next section.

5.4. Full families of α -scaling zeta functions. The development that follows determines the α -scaling zeta functions associated with a weighted self-similar system (Φ, \mathbf{p}) where the distinct regularity values β_q are rationally independent and for each q , the corresponding scaling ratios $\{r_j : \alpha_j = \beta_q\}$ are given by a single value t_q such that $0 < t_q < 1$. In this setting, but unlike in Section 5.3, the full family of α -scaling zeta functions as well as all of the corresponding abscissae of convergence are determined.

Let $c_q = \#\{j : \alpha_j = \beta_q\}$ and $\mathbf{c} := (c_1, \dots, c_{N_0})$. Then $\mathbf{c} \in \mathbb{N}^{N_0}$ and $\sum \mathbf{c} = N$. For any pair of N_0 -vectors \mathbf{m} and \mathbf{t} , let $\mathbf{t}^{\mathbf{m}} := t_1^{m_1} \cdots t_{N_0}^{m_{N_0}}$. The multinomial

theorem implies that for each $n, K \in \mathbb{N}$ we have

$$N^n = \sum_{\sum \mathbf{k}=K} \binom{nK}{n\mathbf{k}} = \sum_{\sum \mathbf{m}=K} \binom{nK}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}}.$$

As shown in the following theorem, the products of the form $\binom{nM}{n\mathbf{m}_0} \mathbf{c}^{n\mathbf{m}_0}$, where $n \in \mathbb{N}$, $M = \sum \mathbf{m}_0$, and $\gcd(\mathbf{m}_0) = 1$, are the multiplicities of the α -scales of the scaling zeta functions associated with (Φ, \mathbf{p}) (cf. [9, §5], especially Theorem 5.2, Lemma 5.10, Proposition 5.11, and Theorem 5.12 therein). Also, the following theorem determines the scaling zeta functions $\zeta_{\mathbf{r}, \mathbf{p}}$ and scaling multifractal spectrum $f_{\mathbf{r}, \mathbf{p}}$ associated with a weighted self-similar system (Φ, \mathbf{p}) .

THEOREM 5.11. *Let (Φ, \mathbf{p}) be a weighted self-similar system. For each $q = 1, \dots, N_0$, suppose there exists a unique t_q such that $0 < t_q < 1$ and for each $j = 1, \dots, N$ such that $\alpha_j = \beta_q, r_j = t_q$. Further, suppose the collection $\{\beta_q\}_{q=1}^{N_0}$ is rationally independent. Then there exists a unique vector $\mathbf{v} = (v_1, \dots, v_{N_0})$ such that $\beta_q = \log_{t_q} v_q$, where $p_j = v_q$ for each $j = 1, \dots, N$ such that $\alpha_j = \beta_q, r_j = t_q$ and some $q \in \{1, \dots, N_0\}$.*

Furthermore, the distinct scaling regularity values attained with respect to (Φ, \mathbf{p}) are given by

$$\beta(\mathbf{m}) := \log_{\mathbf{t}^{\mathbf{m}}} \mathbf{v}^{\mathbf{m}},$$

for some $\mathbf{m} \in (\mathbb{N} \cup \{0\})^{N_0}$ where $\gcd(\mathbf{m}) = 1$ and $\mathbf{m} \neq \mathbf{0}$. Also, for $\mathbf{m} \in (\mathbb{N} \cup \{0\})^{N_0}$ where $\gcd(\mathbf{m}) = 1$ and $\mathbf{m} \neq \mathbf{0}$ and for each $n \in \mathbb{N}$, the number of ways a scaling regularity value $\beta(n\mathbf{m}) = \beta(\mathbf{m})$ is attained with respect to (Φ, \mathbf{p}) at level nM , where $M := \sum \mathbf{m} = \sum_{q=1}^{N_0} m_q$, is given by

$$\binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}}.$$

Moreover, if $\mathbf{m} \in (\mathbb{N} \cup \{0\})^{N_0}$ where $\gcd(\mathbf{m}) = 1$ and $\mathbf{m} \neq \mathbf{0}$, then

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}); s) = \sum_{n=1}^{\infty} \binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}}$$

where $M := \sum \mathbf{m}$, $\operatorname{Re}(s) > f_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}))$, and the corresponding abscissa of convergence $f_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}))$ of $\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}); \cdot)$ is given by

$$f_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m})) = \log_{\mathbf{t}^{\mathbf{m}}} \left(\frac{\mathbf{m}^{\mathbf{m}}}{M^M \mathbf{c}^{\mathbf{m}}} \right).$$

PROOF. Fix $q \in \{1, \dots, N_0\}$. For each $j = 1, \dots, N$ such that $\alpha_j = \beta_q$ we have $p_j = t_q^{\beta_q}$. Thus, $\mathbf{v} := (t_1^{\beta_1}, \dots, t_{N_0}^{\beta_{N_0}})$ is the desired vector.

Let $J \in \mathcal{J}$. By Lemma 5.6, there exist $n \in \mathbb{N}$ and $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$ such that $\gcd(\mathbf{k}) = 1$, $|J| = n \sum \mathbf{k}$, and $A_{\Phi, \mathbf{p}} = \log_{\mathbf{r}^{\mathbf{k}}} \mathbf{p}^{\mathbf{k}}$.

For each $q \in \{1, \dots, N_0\}$, let $m_q := \sum_{j: \alpha_j = \beta_q} k_j$. Define $\mathbf{m} := (m_1, \dots, m_{N_0})$. Then $\sum \mathbf{m} = \sum \mathbf{k}$, $\mathbf{t}^{\mathbf{m}} = \mathbf{r}^{\mathbf{k}}$, and $\mathbf{v}^{\mathbf{m}} = \mathbf{p}^{\mathbf{k}}$. Hence $A_{\Phi, \mathbf{p}}(J) = \beta(\mathbf{m}) := \log_{\mathbf{t}^{\mathbf{m}}} \mathbf{v}^{\mathbf{m}}$. Now, if $\gcd(\mathbf{m}_0) = 1$, then $\beta(\mathbf{m}) = \beta(\mathbf{m}_0)$ if and only if there is $n \in \mathbb{N}$ such that $\mathbf{m} = n\mathbf{m}_0$ since the collection $\{\beta_q\}_{q=1}^{N_0}$ is rationally independent. Moreover, for $M := \sum \mathbf{m}_0$ we have

$$\#\{J \in \mathcal{J}_{n \sum \mathbf{m}_0} : A_{\Phi, \mathbf{p}} = \beta_{\mathbf{m}_0}\} = \binom{nM}{n\mathbf{m}_0} \mathbf{c}^{n\mathbf{m}_0}.$$

Note that each $J \in \mathcal{J}$ will be taken into account since all $\mathbf{m} \in (\mathbb{N} \cup \{0\})^{N_0}$ where $\mathbf{m} \neq \mathbf{0}$ and $\gcd(\mathbf{m}) = 1$ are considered.

Now, if $A_{\Phi, \mathbf{p}}(J) = \beta(\mathbf{m})$, then $r_J = \mathbf{t}^{n\mathbf{m}}$ for some $n \in \mathbb{N}$. Furthermore, for each $n \in \mathbb{N}$

$$\#\{J \in \mathcal{J}_{n \Sigma \mathbf{m}_0} : A_{\Phi, \mathbf{p}} = \beta_{\mathbf{m}_0}\} = \binom{nM}{n\mathbf{m}_0} \mathbf{c}^{n\mathbf{m}_0}.$$

Hence,

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}); s) = \sum_{n=1}^{\infty} \binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}}.$$

In order to determine the abscissa of convergence $f_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}))$, apply Stirling's formula and make use of the function $\psi(s)$ for $s \in \mathbb{R}$ given by

$$\psi(s) = \frac{\mathbf{t}^{s\mathbf{m}} \mathbf{c}^{\mathbf{m}} M^M}{\mathbf{m}^{\mathbf{m}}}.$$

We have $\psi'(s) < 0$ since $0 < t_q < 1$ for $q = 1, \dots, N_0$. Also, $\psi(0) = \mathbf{c}^{\mathbf{m}} M^M / \mathbf{m}^{\mathbf{m}} > 1$ since $\mathbf{m} \neq \mathbf{0}$, $M = \sum \mathbf{m}$, and $c_q > 0$ for each $q = 1, \dots, N_0$. Hence, there is a real unique real number ρ such that $\rho > 0$ and

$$1 = \frac{\mathbf{t}^{\rho\mathbf{m}} \mathbf{c}^{\mathbf{m}} M^M}{\mathbf{m}^{\mathbf{m}}}.$$

The real number ρ will prove to be our abscissa of convergence. Indeed, for a fixed real number s , Stirling's formula yields

$$\begin{aligned} \binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}} &= \frac{(nM)!}{(nm_1)! \cdots (n_m N_0)!} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}} \\ &= \frac{\mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}} M^{nM}}{\mathbf{m}^{n\mathbf{m}}} \cdot \frac{\sqrt{2\pi nM}}{\mathbf{m}^{n\mathbf{m}}} (1 + \varepsilon_n), \end{aligned}$$

where $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. Hence,

$$\left[\binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}} \right]^{1/n} = \frac{\mathbf{c}^{\mathbf{m}} \mathbf{t}^{s\mathbf{m}} M^M}{\mathbf{m}^{\mathbf{m}}} (1 + \delta_n),$$

where $\delta_n \rightarrow 0$ as $n \rightarrow \infty$. The root test implies that the numerical series

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}); s) = \sum_{n=1}^{\infty} \binom{nM}{n\mathbf{m}} \mathbf{c}^{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}}$$

converges for $s > \rho$ and diverges for $s < \rho$. Therefore,

$$f_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m})) = \rho = \log_{\mathbf{t}^{\mathbf{m}}} \left(\frac{\mathbf{m}^{\mathbf{m}}}{\mathbf{c}^{\mathbf{m}} M^M} \right).$$

□

COROLLARY 5.12. *If the conditions of Theorem 5.11 are satisfied and, additionally, if $N_0 = N$, then Proposition 2.18 is recovered for scaling regularity values $\alpha(\mathbf{k})$. Specifically, in this case we have*

$$f_{\mathbf{r}, \mathbf{p}}(\alpha(\mathbf{k})) = \log_{\mathbf{r}^{\mathbf{k}}} \left(\frac{\mathbf{k}^{\mathbf{k}}}{K^K} \right) = \frac{\sum_{j=1}^N (k_j/k) \log k_j/K}{\sum_{j=1}^N (k_j/K) \log r_j} = \dim_H(F(\mathbf{k}/K)),$$

where $F(\mathbf{k}/K)$ is the Besicovitch subset (as given in Definition 2.16) of the self-similar set F .

REMARK 5.13. In addition to the recovery of Proposition 2.18, Theorem 5.11 recovers and allows for the generalization of all of the results found in Section 5 of [9] where the family of partitions is taken to be the *natural* family of partitions (adapted to Φ). This generalization includes a partial recovery of the behavior of the multifractal spectrum discussed in Remark 2.14 when $N_0 = 2$ and $t_1 = t_2$, as in Example 2.15. (The recovery is partial due to the restriction to a countable collection of scaling regularity values; however, taking the concave envelope of the resulting graph yields the entire spectrum.) Moreover, due to the use of scaling regularity in place of coarse Hölder regularity, *our results extend the results of Section 5 of [9] to the setting of arbitrary self-similar measures supported on self-similar subsets of some Euclidean space of any dimension, as opposed to self-similar measures supported on subsets of the unit interval $[0, 1]$.*

REMARK 5.14 (Natural Hausdorff measures). Consider a weighted self-similar system (Φ, \mathbf{p}) , where $\mathbf{p} = (r_1^D, \dots, r_N^D)$ such that D is the unique real (and hence, positive) root of the corresponding Moran equation (2). The resulting self-similar measure μ is the natural Hausdorff measure of the underlying fractal support. Such measures are used to find lower bounds on the Hausdorff dimension of their supports; see [12, Ch. 9]. Moreover, $\alpha = D$ is the only nontrivial scaling regularity value associated with (Φ, \mathbf{p}) . One then says that μ is *monofractal*. On the other hand, if $\mathbf{p} \neq (r_1^D, \dots, r_N^D)$, then there are countably many nontrivial regularity values associated with (Φ, \mathbf{p}) . It follows that μ is truly *multifractal* in this case.

In the case of the ternary Cantor set with $\mathbf{r} = (1/3, 1/3)$ weighted by $\mathbf{p} = (1/2, 1/2)$, the graph of the primitive of the natural Hausdorff measure μ (i.e., the graph of $\mu([0, x])$) is the well-known Devil's staircase; see, [26, Plate 83, p. 83], [13, Ch. 6], [24, §12.1.1] and Example 2.15 above. The primitive of μ is the Cantor–Lebesgue function. Recall that this function is a nondecreasing, surjective, and continuous function from $[0, 1]$ to itself which has derivative zero on $[0, 1] \setminus F$, where F is the Cantor set.

In the next section we close the paper by proposing some preliminary ideas for further results and future work.

6. Further results and future work

The results presented in this paper suggest several interesting problems to pursue. For instance, since our results stem directly from scaling ratios and probability vectors, one may consider interpreting analogous results on an arbitrary metric space along with an appropriate space of measures. Also, the following conjecture, which was originally stated (in a similar but more restrictive setting) in Conjecture 5.8 of [9], is not addressed in the rest of the present paper.

CONJECTURE 6.1. *For a self-similar measure μ and all $t \in [t_{\min}, t_{\max}]$, we have*

$$\hat{f}(t) = f_g(t) = f_s(t),$$

where $\hat{f}(t)$ is the concave envelope of the scaling multifractal spectrum $f(\alpha)$ on $[t_{\min}, t_{\max}]$, and f_g and f_s are the geometric and symbolic Hausdorff multifractal spectra defined in Section 2.

Another problem worthy of study, but not addressed in this paper and yet motivated by the theory of complex of dimensions of fractal strings in [24], is the determination of the full collection of the sets of α -scaling complex dimensions $\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\alpha)$

of a weighted self-similar system (Φ, \mathbf{p}) with respect to the nontrivial regularity values α and, in turn, the determination of the tapestry of complex dimensions $\mathcal{T}_{\mathbf{r}, \mathbf{p}}$. Aside from weighted self-similar systems with regularity values satisfying the conditions of Theorem 5.8 and Corollary 5.9, the α -scaling complex dimensions associated with other regularity values, as in Theorem 5.11, are not known.

Nonetheless, the final two sections provide interesting partial results on such further problems as well as motivation for further research.

6.1. Generalized hypergeometric series. In this section, the scaling zeta functions found in Corollary 5.12 (of Theorem 5.11) are shown to be generalized hypergeometric series (see [2]). Such series have been well studied and well understood, and related work may provide an alternative or supplementary approach to the theory of complex dimensions of self-similar measures.

Let $b > 0$, $\mathbf{k} = (k_1, \dots, k_N) \in (\mathbb{N} \cup \{0\})^N$ for some $N \in \mathbb{N} \setminus \{0, 1\}$, $K := \sum \mathbf{k}$ and $B(\mathbf{k}) := K^K / \mathbf{k}^{\mathbf{k}}$. (It will help to keep Notation 5.5 in mind in the sequel.)

Consider the following Dirichlet series:

$$\zeta_b(\mathbf{k}; s) = \sum_{n=1}^{\infty} \binom{nK}{n\mathbf{k}} b^{-nKs}.$$

This form of Dirichlet series appears, indirectly, in Section 5. Indeed, if the conditions of Theorem 5.11 are satisfied and, additionally, if $N_0 = N$, then we take $b^{-1} = \mathbf{r}^{\mathbf{k}/K} = \mathbf{t}^{\mathbf{m}/M}$ with $K = M$ such that $\mathbf{r} = \mathbf{t}$ is the vector of scaling ratios of the corresponding weighted self-similar system. We get

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}) : s) = \sum_{n=1}^{\infty} \binom{nM}{n\mathbf{m}} \mathbf{t}^{sn\mathbf{m}} = \sum_{n=1}^{\infty} \binom{nK}{n\mathbf{k}} b^{-nKs} = \zeta_b(\mathbf{k}; s).$$

Now, via the formula

$$\binom{nK}{n\mathbf{k}} = \frac{\prod_{j=1}^K (1((j/K) + n)/1(j/K))}{\prod_{q=1}^N \left(\prod_{j=1}^{k_q} (1((j/k_q) + n)/1(j/k_q)) \right)} \cdot \frac{B(\mathbf{k})^n}{n!}$$

we have

$$(20) \quad \zeta_b(\mathbf{k}; s) = -1 + \sum_{n=0}^{\infty} \frac{\prod_{j=1}^K (1((j/K) + n)/1(j/K))}{\prod_{q=1}^N \left(\prod_{j=1}^{k_q} (1((j/k_q) + n)/1(j/k_q)) \right)} \cdot \frac{(B(\mathbf{k})b^{-Ks})^n}{n!}.$$

The series on the right-hand side of (20) is a generalized hypergeometric series of the form ${}_K F_{K-1}$ (see [2] for the precise definition and notation). That is, we have

$$\begin{aligned} \zeta_b(\mathbf{k}; s) &= -1 + \sum_{n=0}^{\infty} \frac{\prod_{j=1}^K (1((j/K) + n)/1(j/K))}{\prod_{q=1}^N \left(\prod_{j=1}^{k_q} (1((j/k_q) + n)/1(j/k_q)) \right)} \cdot \frac{(B(\mathbf{k})b^{-Ks})^n}{n!} \\ &= -1 + {}_K F_{K-1}(1/K, \dots, 1; 1/k_1, \dots, (k_n - 1)/k_N; B(\mathbf{k})b^{-Ks}). \end{aligned}$$

Thus, even though the full families of scaling zeta functions of a self-similar measure in the general case have yet to be determined, the collection of scaling zeta functions which are given by a hypergeometric series is primed for further analysis. As seen in the next section, there are scaling zeta functions which are neither nearly the zeta function of a self-similar system nor a hypergeometric series, even in the case where the distinct scaling ratios are rationally independent.

6.2. A self-similar system with nonlattice and lattice structure. This last section investigates the structure of the scaling zeta functions in a special case where the distinct scaling regularity values are rationally independent but only two of the scaling zeta functions are known.

Consider a weighted self-similar system (Φ, \mathbf{p}) such that Φ satisfies the open set condition and is nonlattice with scaling ratios $\mathbf{r} \in (0, 1)^N$ for some $N \in \mathbb{N}$, $N \geq 3$, satisfying the following conditions: First, there exist $t, t_0 \in (0, 1)$ and $\mathbf{t} := (t, t^2, t_0)$ such that for each $j = 1, \dots, N$, r_j is equal to either t, t^2 , or t_0 . Second, there exists $\mathbf{c} = (c_1, c_2, c_0) \in \mathbb{N}^3$ such that $\sum \mathbf{c} = N$ and

$$\begin{aligned} c_1 &= \#\{j : r_j = t, j \in \{1, \dots, N\}\}, \\ c_2 &= \#\{j : r_j = t^2, j \in \{1, \dots, N\}\}, \end{aligned}$$

and

$$c_0 = \#\{j : r_j = t_0, j \in \{1, \dots, N\}\}.$$

Further, suppose there are a probability vector \mathbf{p} and rationally independent real numbers γ and γ_0 such that for each $j \in \{1, \dots, N\}$, $r_j = t$ implies $p_j = t^\gamma$; $r_j = t^2$ implies $p_j = t^{2\gamma}$; and $r_j = t_0$ implies $p_j = p_0 := t_0^{\gamma_0}$.

Due to Theorem 5.11, in order to determine the distinct scales $l_n(\alpha)$ and the corresponding multiplicities $m_n(\alpha)$ of the sequence of α -scales $\mathcal{L}_{\mathbf{r}, \mathbf{p}}(\alpha)$, it suffices to consider the scaling regularity values $\alpha(\mathbf{k})$ where $\mathbf{k} \in (\mathbb{N} \cup \{0\})^N$, $\mathbf{k} \neq \mathbf{0}$, and $\gcd(\mathbf{k}) = 1$.

Fix \mathbf{k} as above. Let

$$m := \sum_{j:r_j=t} k_j + \sum_{j:r_j=t^2} k_j, \quad m_0 := \sum_{j:r_j=t_0} k_j,$$

and $\mathbf{m} := (m, m_0)$. We have

$$\alpha(\mathbf{k}) := \log_{\mathbf{r}^{\mathbf{k}}} \mathbf{p}^{\mathbf{k}} = \frac{\log t^{m\gamma} t_0^{m_0\gamma_0}}{\log t^m t_0^{m_0}}.$$

Define $\beta(\mathbf{m}) := \log t^{m\gamma} t_0^{m_0\gamma_0} / \log t^m t_0^{m_0}$.

Now, for a fixed \mathbf{m} where $\gcd(\mathbf{m}) = 1$ and each $n \in \mathbb{N}$ we have

$$l_n(\beta(\mathbf{m})) = (t^m t_0^{m_0})^n.$$

The key difficulty here lies in determining the multiplicities $m_n(\beta(\mathbf{m}))$ of the scales $l_n(\beta(\mathbf{m}))$, specifically, due to the fact that some initial scaling ratios are equal to t and others are equal to t^2 . Thus, for instance, the γ -scale t^4 is attained with respect to some vectors $\mathbf{k}_1, \mathbf{k}_2$, and \mathbf{k}_3 where $|\mathbf{k}_1| = 2$, $|\mathbf{k}_2| = 3$, and $|\mathbf{k}_3| = 4$. More specifically, if both nonzero components of \mathbf{k}_1 correspond to the scale t^2 , two of the nonzero components of \mathbf{k}_2 correspond to t and the other corresponds to t^2 , and all four nonzero components of \mathbf{k} correspond to t , then

$$\alpha(\mathbf{k}_1) = \alpha(\mathbf{k}_2) = \alpha(\mathbf{k}_3) = \beta(\mathbf{m}) = \gamma$$

and

$$\mathbf{r}^{\mathbf{k}_1} = \mathbf{r}^{\mathbf{k}_2} = \mathbf{r}^{\mathbf{k}_3} = t^4.$$

That is, a given scale $l_n(\beta(\mathbf{m}))$ can arise in various stages with respect to (Φ, \mathbf{p}) , making the determination of the precise form of the multiplicity $m_n(\beta(\mathbf{m}))$ difficult in general. As a result, and although the full family of scaling zeta functions in the setting of this section have been determined, the abscissae of convergence and the

α -scaling complex dimensions are known for just two nontrivial scaling regularity values.

The following notation is used repeatedly below. Let $\lfloor \cdot \rfloor$ denote the floor function. That is, for $x \in \mathbb{R}$, $\lfloor x \rfloor$ is the integer part of x given by the greatest integer such that $\lfloor x \rfloor \leq x$.

Suppose $\gcd(\mathbf{m}) = 1$ and set $M = \sum \mathbf{m}$. Then, for each $n \in \mathbb{N}$,

$$(21) \quad \begin{aligned} m_n(\beta(\mathbf{m})) &= \#\{J \in \mathcal{J} : r_J = l_n(\beta(\mathbf{m}))\} \\ &= \sum_{j=0}^{\lfloor nM/2 \rfloor} \binom{\sum \mathbf{v}(j)}{\mathbf{v}(j)} \mathbf{c}^{\mathbf{v}(j)}, \end{aligned}$$

with $\mathbf{v}(j) = (v_1(j), v_2(j), v_0)$ where, for $j = 0, \dots, \lfloor nM/2 \rfloor$,

$$v_1(j) := nm - 2 \lfloor nM/2 \rfloor + 2j, \quad v_2(j) := \lfloor nM/2 \rfloor - j, \quad \text{and } v_0 := nm_0.$$

The difficulty in determining the form of the generalization of $m_n(\beta(\mathbf{m}))$ lies in determining the relationship between the components of the generalization of the vectors $\mathbf{v}(j)$. The $\beta(\mathbf{m})$ -scaling zeta function of (Φ, \mathbf{p}) for some \mathbf{m} with $\gcd(\mathbf{m}) = 1$ is given by

$$\zeta_{\mathbf{r}, \mathbf{p}}(\beta(\mathbf{m}); s) = \sum_{n=1}^{\infty} m_n(\beta(\mathbf{m})) (l_n(\beta(\mathbf{m})))^s,$$

where $\operatorname{Re}(s)$ is large enough.

EXAMPLE 6.2. In this example, the closed forms of the scaling zeta functions for just two nontrivial scaling regularity values, namely γ_0 and γ , are known. Both cases are the result of Corollary 5.9, thus the abscissae of convergence and the α -scaling complex dimensions associated with just two nontrivial regularity values can be had.

We have $\gcd(\mathbf{m}) = 1$ and $\beta(\mathbf{m}) = \gamma_0$ if and only if $m = 0$ and $m_0 = 1$. In this case, $l_n(\gamma_0) = t_0^n$, $m_n(\gamma_0) = c_0^n$, and by Corollary 5.9 we have, for $s \in \mathbb{C}$,

$$\zeta_{\mathbf{r}, \mathbf{p}}(\gamma_0; s) = \frac{c_0 t_0^s}{1 - c_0 t_0^s}.$$

Moreover,

$$\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\gamma_0) = \left\{ -\log_{t_0} c_0 + i \frac{2\pi}{\log t_0} z : z \in \mathbb{Z} \right\}.$$

As for scaling regularity γ , we have $\gcd(\mathbf{m}) = 1$ and $\beta(\mathbf{m}) = \gamma$ if and only if $m = 1$ and $m_0 = 0$. In this case, $l_n(\gamma) = t^n$ and

$$m_n(\gamma) = \sum_{j=0}^{\lfloor n/2 \rfloor} \binom{\lfloor \frac{n+1}{2} \rfloor + j}{n - 2 \lfloor \frac{n}{2} \rfloor + 2j} \mathbf{c}^{\mathbf{v}(j)}.$$

However, a closed form of $\zeta_{\mathbf{r}, \mathbf{p}}(\gamma; s)$ is obtained in Corollary 5.9. That is,

$$\zeta_{\mathbf{r}, \mathbf{p}}(\gamma; s) = \frac{c_1 t_1^s + c_2 t_2^{2s}}{1 - c_1 t_1^s - c_2 t_2^{2s}}, \quad s \in \mathbb{C}.$$

Moreover, $\zeta_{\mathbf{r}, \mathbf{p}}(\gamma; s) = \zeta_{\eta}(s)$, where η is the generalized lattice string with scaling ratios $r_1 = t$ and $r_2 = t^2$, weights $m_1 = c_1$ and $m_2 = c_2$, and c_1 gaps $g_1 = t$ and c_2 gaps $g_2 = t^2$. Hence, $\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\gamma) = \mathcal{D}_{\eta}$.

EXAMPLE 6.3. This example develops the full recovery of the Fibonacci string as in Example 3.6 as well as the results of Example 5.16 from [9].

Consider a weighted self-similar system (Φ, \mathbf{p}) such that Φ satisfies the open set condition with scaling ratios $\mathbf{r} = (1/2, 1/4, 1/10)$. Then Φ is nonlattice. Further, suppose the probability vector \mathbf{p} is given by $\mathbf{p} = (1/2, 1/4, 1/4)$ such that the distinct initial scaling regularity values $\gamma = 1$ and $\gamma_0 = \log_{10} 4$ are rationally independent. Moreover, in the notation given by the more general setting above, we have $t = 1/2, t_0 = 1/10$ and $c_1 = c_2 = c_0 = 1$. As in the previous example, the only scaling regularity values for which the scaling zeta functions are known are $\gamma_0 = \log_{10} 4$ and $\gamma = 1$, so only these two cases are discussed in this example.

First consider the simplest case of scaling regularity $\gamma_0 = \log_{10} 4$. The γ_0 -scaling zeta function is given by

$$\zeta_{\mathbf{r}, \mathbf{p}}(\gamma_0; s) = \sum_{n=1}^{\infty} 10^{-ns} = \frac{10^{-s}}{1 - 10^{-s}}.$$

Thus, the set of γ_0 -scaling complex dimensions is given by

$$\mathcal{D}_{\mathbf{r}, \mathbf{p}}(\gamma_0) = \{izp : z \in \mathbb{Z}\},$$

where $p = 2\pi/\log 10$.

In this case of scaling regularity $\gamma = 1$, we nearly recover the geometric zeta function of the Fibonacci string ζ_{Fib} . The 1-scales are given by $l_n(1) = 2^{-n}$ for $n \in \mathbb{N}$ and the multiplicities $m_n(1)$ are given by the Fibonacci numbers. In fact, in light of the formula for the more general $m_n(\beta(\mathbf{m}))$ in (21) and the formulas from Example 3.6, we recover a classic formula which yields the Fibonacci numbers as sums of particular binomial coefficients. Specifically,

$$m_n(1) = \sum_{j=0}^{\lfloor \frac{n}{2} \rfloor} \binom{\lfloor \frac{n+1}{2} \rfloor + j}{n - 2 \lfloor \frac{n}{2} \rfloor + 2j} = F_{n+1},$$

where F_{n+1} is the $(n+1)$ th Fibonacci number. See Example 3.6 above, Example 5.16 of [9], and [24, §2.3.2] for a discussion of the Fibonacci string and its geometric zeta function ζ_{Fib} .

The scaling zeta function $\zeta_{\mathbf{r}, \mathbf{p}}(1; s)$ is then given by

$$\zeta_{\mathbf{r}, \mathbf{p}}(1; s) = \sum_{n=1}^{\infty} F_{n+1} 2^{-ns} = \zeta_{\text{Fib}}(s) - 1 = \frac{2^{-s} + 4^{-s}}{1 - 2^{-s} - 4^{-s}},$$

where ζ_{Fib} is given by (5). Thus, the corresponding complex dimensions, in both the classic sense and with respect to the scaling regularity 1, are given by (7). That is,

$$\mathcal{D}_{\text{Fib}} = \mathcal{D}_{\mathbf{r}, \mathbf{p}}(1) = \{D_{\text{Fib}} + izp : z \in \mathbb{Z}\} \cup \{-D_{\text{Fib}} + i(z + 1/2)p : z \in \mathbb{Z}\},$$

where $\varphi = (1 + \sqrt{5})/2$ is the Golden Ratio, $D_{\text{Fib}} = \log_2 \varphi$, and the oscillatory period is $p = 2\pi/\log 2$.

In closing, we mention that some next steps include the determination of the α -scaling complex dimensions and the full tapestry of complex dimensions $\mathcal{T}_{\mathbf{r}, \mathbf{p}}$ associated with such self-similar measures (cf. [9, §6]). Thereafter, one may study the implications (for the oscillatory behavior of self-similar systems and measures) of counting functions and volume formulas associated with the nontrivial scaling

regularity values, in the spirit of similar notions stemming from the theory of complex dimensions of fractal strings developed throughout [24] (see also [18] and the relevant references therein, for the higher-dimensional case).

References

- [1] V. K. Balakrishnan, *Introductory Discrete Mathematics*, Courier Dover Publications, 1996. (Reprint of the Prentice-Hall, Englewood Cliffs, New Jersey, 1991 edition.)
- [2] G. Bateman and A. Erdelyi, *Higher Transcendental Functions*, vol. 1, New York, McGraw-Hill, 1953.
- [3] A. S. Besicovitch, On the sum of digits of real numbers represented in the dyadic system, *Math. Ann.* **110** (1934) 321–330.
- [4] A. S. Besicovitch and S. J. Taylor, On the complementary intervals of a linear closed set of zero Lebesgue measure, *J. London Math. Soc.* **29** (1954), 449–459.
- [5] G. Brown, G. Michon and J. Peyrière, On the multifractal analysis of measures, *J. Statist. Phys.* **66** (1992), 775–790.
- [6] R. Cawley and R. D. Mauldin, Multifractal decompositions of Moran fractals, *Adv. in Math.* **92** (1992), 196–236.
- [7] F. M. Dekking and W. X. Li, Hausdorff dimensions of subsets of Moran fractals with prescribed group frequency of their codings, *Nonlinearity*, **16** (2003), 1–13.
- [8] R. de Santiago, The Recursive Structure of Lattice Strings, *Master's thesis*, California State Polytechnic University, Pomona, in progress, 2012.
- [9] K. E. Ellis, M. L. Lapidus, M. C. Mackenzie, and J. A. Rock, Partition zeta functions, multifractal spectra, and tapestries of complex dimensions, in *Benoit Mandelbrot: A Life in Many Dimensions* (M. Frame, Ed.), World Scientific, Singapore, in press, 2012. (arXiv:1007.1467v2 [math-ph], 2011).
- [10] H. G. Eggleston, The fractional dimension of a set defined by decimal properties, *Quart. J. Math. Oxford Ser.* **20** (1949), 31–36.
- [11] D. Essouabri and M. L. Lapidus, Analytic continuation of a class of multifractal zeta function, preprint (and work in progress), 2012.
- [12] K. Falconer, *Fractal Geometry – Mathematical foundations and applications*, 2nd ed., John Wiley, Chichester, 2003.
- [13] J. Feder, *Fractals*, Plenum Press, New York, 1988.
- [14] J. E. Hutchinson, Fractals and self-similarity, *Indiana Univ. Math. J.* **30** (1981), 713–747.
- [15] S. Jaffard, Wavelet techniques in multifractal analysis, in: [23], Part 2, pp. 91–151.
- [16] M. Kesseböhmer and S. Kombrink, Fractal curvature measures and Minkowski content for self-conformal subsets of the real line, *Adv. in Math.* **230** (2012), 2474–2512.
- [17] M. L. Lapidus, J. Lévy-Véhel and J. A. Rock, Fractal strings and multifractal zeta functions, *Letters in Mathematical Physics* No. 1, **88** (2009), 101–129. (Special issue dedicated to the memory of Moshe Flato.) (Also: E-print, arXiv:math-ph/0610015v3, 2009.)
- [18] M. L. Lapidus, E. P. J. Pearse and S. Winter, Pointwise tube formulas for fractal sprays and self-similar tilings with arbitrary generators, *Adv. in Math.* **227** (2011), 1349–1398. (Also: E-print, arXiv:1006.3807v3 [math.MG], 2011.)
- [19] M. L. Lapidus, E. P. J. Pearse and S. Winter, Minkowski measurability results for self-similar tilings and fractals with monophase generators, E-print, arXiv:1104.1641v2 [math.MG], 2012.
- [20] M. L. Lapidus and C. Pomerance, The Riemann zeta-function and the one-dimensional Weyl–Berry conjecture for fractal drums, *Proc. London Math. Soc.* (3) **66** (1993), 41–69.
- [21] M. L. Lapidus and C. Pomerance, Counterexamples to the modified Weyl–Berry conjecture on fractal drums, *Math. Proc. Cambridge Philos. Soc.* **119** (1996), 167–178.
- [22] M. L. Lapidus and J. A. Rock, Towards zeta functions and complex dimensions of multifractals, *Complex Variables and Elliptic Equations* No. 6, **54** (2009), 545–559. (Special issue dedicated to Fractal Analysis.) (Also: E-print, Institut des Hautes Etudes Scientifiques, IHES/M/08/34, 2008.)
- [23] M. L. Lapidus and M. van Frankenhuysen (eds.), *Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot*, Proceedings of Symposia in Pure Mathematics **72** (Part 1: *Analysis, Number Theory and Dynamical Systems*. Part 2: *Multifractals, Probability and Statistical Mechanics, and Applications*), Amer. Math. Soc., Providence, RI, 2004.

- [24] M. L. Lapidus and M. van Frankenhuysen, *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and spectra of fractal strings*, 2nd revised and enlarged edition (of the 2006 edition), Springer Monographs in Mathematics, Springer-Verlag, New York, in press, 2012.
- [25] J. Lévy-Véhel and F. Mendivil, Multifractal and higher-dimensional zeta functions, *Nonlinearity* No. 1, **24** (2011), 259–276.
- [26] B. B. Mandelbrot, *The Fractal Geometry of Nature*, revised and enlarged edition (of the 1977 edition), W. H. Freeman, New York, 1983.
- [27] R. D. Mauldin and M. Urbański, *Graph Directed Markov Systems: Geometry and Dynamics of Limit Sets*, Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 2003.
- [28] P. A. P. Moran, Additive functions of intervals and Hausdorff measure, *Math. Proc. Cambridge Philos. Soc.* **42** (1946), 15–23.
- [29] L. Olsen, *Random Geometrically Graph Directed Self-Similar Multifractals*, Pitman Research Notes in Math. Series, vol. 307, Longman Scientific and Technical, London, 1994.
- [30] L. Olsen, A multifractal formalism, *Adv. in Math.* **116** (1996), 82–196.
- [31] L. Olsen, Multifractal geometry, in: *Fractal Geometry and Stochastics II* (Greifswald/Koserow, 1998), *Prog. Prob.* **46** (2000), Birkhäuser, Basel, pp. 3–37.
- [32] L. Olsen, A lower bound for the symbolic multifractal spectrum of a self-similar multifractal with arbitrary overlaps, *Math. Nachr.* No. 10, **282** (2009), 1461–1477.
- [33] J. A. Rock, Zeta Functions, Complex Dimensions of Fractal Strings and Multifractal Analysis of Mass Distributions, *Ph.D. Dissertation*, University of California, Riverside, 2007.
- [34] R. S. Strichartz, A. Taylor, and T. Zhang, Densities of Self-Similar Measures on the Line, *J. Experimental Mathematics* No. 2, **4** (1995), 101–128.

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE POLYTECHNIC UNIVERSITY, POMONA, CALIFORNIA 91768 USA

E-mail address: rdesantiago@csupomona.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CALIFORNIA 92521-0135 USA

E-mail address: lapidus@math.ucr.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, RIVERSIDE, CALIFORNIA 92521-0135 USA

E-mail address: roby@math.ucr.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, CALIFORNIA STATE POLYTECHNIC UNIVERSITY, POMONA, CALIFORNIA 91768 USA

E-mail address: jarock@csupomona.edu