

Extensions of flat functors and theories of presheaf type

Olivia CARMELLO



Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Juin 2014

IHES/M/14/23

Extensions of flat functors and theories of presheaf type

Olivia Caramello*

Institut des Hautes Études Scientifiques
35 route de Chartres 91440, Bures-sur-Yvette, France
olivia@ihes.fr

June 20, 2014

Abstract

We develop a general theory of extensions of flat functors along geometric morphisms of toposes, and apply it to the study of the class of theories whose classifying topos is equivalent to a presheaf topos. As a result, we obtain a characterization theorem providing necessary and sufficient semantic conditions for a theory to be of presheaf type. This theorem subsumes all the previous partial results obtained on the subject and has several corollaries which can be used in practice for testing whether a given theory is of presheaf type as well as for generating new examples of theories belonging to this class. Along the way, we establish a number of other results of independent interest, including developments about colimits in the context of indexed categories, expansions of geometric theories and methods for constructing theories classified by a given presheaf topos.

Contents

1	Introduction	3
2	Indexed colimits in toposes	7
2.1	Background on indexed categories	7
2.2	$\mathcal{E}$ -filtered indexed categories	10
2.3	Indexation of internal diagrams	10
2.4	Colimits and tensor products	11
2.5	$\mathcal{E}$ -final subcategories	17

*Supported by a CARMIN IHÉS-IHP post-doctoral position (as from 1/12/2013) and by a visiting position of the Max Planck Institute for Mathematics (in the period 1/10/2013 - 30/11/2013).

2.6	A characterization of $\mathcal{E}$ -indexed colimits	24
2.7	Explicit calculation of set-indexed colimits	36
3	Extensions of flat functors	44
3.1	General extensions	45
3.2	Extensions along embeddings of categories	46
3.3	Extensions from categories of set-based models to syntactic categories	49
3.4	A general adjunction	55
4	Preliminary results on theories of presheaf type	61
4.1	A syntactic description of finitely presentable models	61
4.2	Objects of homomorphisms	63
4.3	Strong finite presentability	66
4.4	Semantic $\mathcal{E}$ -finite presentability	69
5	Semantic criteria for a theory to be of presheaf type	71
5.1	The characterization theorem	71
5.2	Concrete reformulations	77
5.2.1	Condition (i)	77
5.2.2	Condition (ii)	81
5.2.3	Condition (iii)	84
5.3	Abstract reformulation	93
6	Faithful interpretations of theories of presheaf type	94
6.1	General results	94
6.2	Finitely presentable and finitely generated models	99
6.3	Reformulations of condition (iii) of Theorem 5.1	102
6.4	Quotient theories	106
6.4.1	Presheaf-type quotients and rigid topologies	108
6.4.2	Finding theories classified by a given presheaf topos	111
6.5	Injectivizations of theories	115
6.5.1	Condition (iii) of Theorem 5.1 for injectivizations	118
6.5.2	A criterion for injectivizations	122
7	Expansions of theories	127
7.1	General theory	127
7.2	Expanding a geometric theory to a theory of presheaf type	135
7.3	Faithful expansions	137
8	Examples	140
8.1	Theories whose finitely presentable models are finite	140
8.1.1	The theory of abstract circles	140
8.1.2	The geometric theory of finite sets	143
8.2	The theory of Diers' fields	146

8.3	The theory of algebraic extensions of a given field	150
8.4	Groups with decidable equality	152
8.5	Locally finite groups	154
8.6	Vector spaces	155
8.7	The theory of abelian l -groups with strong unit	155

1 Introduction

Following [2], we say that a geometric theory is of *of presheaf type* if it is classified by a presheaf topos.

A geometric theory $\mathbb{T}$ is of presheaf type if and only if it is classified by the topos $[\mathbf{f.p.T-mod}(\mathbf{Set}), \mathbf{Set}]$, where $\mathbf{f.p.T-mod}(\mathbf{Set})$ is (a skeleton of) the full subcategory of $\mathbf{T-mod}(\mathbf{Set})$ on the finitely presentable $\mathbf{T}$ -models (cf. [5]).

The subject of theories of presheaf type has a long history, starting with the book [19] by Hakim, which first introduced the point of view of classifying toposes in the context of the theory of commutative rings with unit and its quotients. The subsequent pioneering work [25] by Lawvere led to the discovery that any finitary algebraic theory is of presheaf type, classified by the topos of presheaves on the opposite of its category of finitely presentable models (cf. [23]). This result was later generalized to cartesian (or essentially algebraic) theories as well as to universal Horn theories (cf. [3]). At the same time, new examples of non-cartesian theories of presheaf type were discovered (cf. for instance [2] for a long, but by no means exhaustive, list of examples), and partial results in connection to the problem of characterizing the class of theories of presheaf type emerged; for instance, [22], [2] and [32] contain different sets of sufficient conditions for a theory to be of presheaf type.

Theories of presheaf type occupy a central role in Topos Theory for a number of reasons:

- (i) Every small category $\mathcal{C}$ can be seen, up to Cauchy-completion, as the category of finitely presentable models of a theory of presheaf type (namely, the theory of flat functors on $\mathcal{C}^{\text{op}}$);
- (ii) As every Grothendieck topos is a subtopos of some presheaf topos, so every geometric theory is a quotient of some theory of presheaf type (cf. the duality theorem of [9] between subtoposes of the classifying topos of a geometric theory and quotients of the theory);
- (iii) Every finitary algebraic theory (and more generally, any cartesian theory) is of presheaf type;
- (iv) The class of theories of presheaf type contains, besides all cartesian theories, many other interesting mathematical theories pertaining to different fields of mathematics (for instance, the coherent theory of

linear orders or the geometric theory of algebraic extensions of a given field);

- (v) The ‘bridge technique’ of [10] can be fruitfully applied in the context of theories of presheaf type due to the fact that the classifying topos of any such theory admits (at least) two quite different representations, one of semantic nature (namely, set-valued functors on the category of finitely presentable models of the theory) and one of syntactic nature (namely, sheaves on the syntactic site of the theory).

It is therefore important to dispose of effective criteria for testing whether a theory is of presheaf type, as well as of methods for generating new theories of presheaf type.

In this paper, we carry out a systematic investigation of this class of theories, obtaining in particular a characterization theorem providing necessary and sufficient conditions for a theory to be of presheaf type, expressed in terms of the models of the theory in arbitrary Grothendieck toposes. This theorem, whose general statement is quite abstract, admits several ramifications and simpler corollaries which can be effectively applied in practice to test whether a given theory is of presheaf type as well as for generating new examples of theories of presheaf type, also through appropriate ‘modifications’ of given geometric theories. All the partial results and recognition criteria previously obtained on the subject are naturally subsumed by this general result; moreover, the constructive nature of the characterization theorem allows to replace the requirements that the theory should have enough set-based models in the sufficient criteria of [2] and [22] with explicit semantic conditions which can be directly verified without having to invoke any form of the axiom of choice.

In order to establish our characterization theorem, we embark, in the first two sections of the paper, in a general analysis of indexed colimits in toposes and extensions of flat functors along geometric morphisms. In fact, the necessary and sufficient conditions for a theory $\mathbb{T}$ to be of presheaf type provided by the characterization theorem arise precisely from the requirement that for any Grothendieck topos $\mathcal{E}$, the operation of extension of flat functors with values in $\mathcal{E}$ from the opposite of the category of finitely presentable models of $\mathbb{T}$ to the geometric syntactic category of $\mathbb{T}$ should define an equivalence of categories onto the category of $\mathbb{T}$ -models in $\mathcal{E}$, naturally in $\mathcal{E}$. We then investigate the preservation, by ‘faithful interpretations’ of theories, of each of the conditions in the characterization theorem, obtaining results of the form ‘under appropriate conditions, a geometric theory in which a theory of presheaf type faithfully interprets is again of presheaf type’. Finally, we discuss known and new examples of theories of presheaf type in light of the theory developed in the paper.

More specifically, the contents of the paper can be summarized as follows.

In section 2, we investigate $\mathcal{E}$ -indexed colimits of internal diagrams in Grothendieck toposes $\mathcal{E}$, analyzing in particular their behaviour with respect to final $\mathcal{E}$ -indexed functors (cf. section 2.5) and establishing explicit characterizations for a $\mathcal{E}$ -indexed cocone to be colimiting (cf. section 2.6). In section 2.4, we exploit the abstract interpretation of colimits as kinds of tensor products to derive commutation results which play an important role in the subsequent parts of the paper as they allow us to interpret certain set-indexed colimits arising in the context of our main characterization theorem as special kinds of filtered indexed colimits.

In section 3, we investigate the properties of the operation on flat functors induced by a geometric morphism of toposes via Diaconescu's equivalence. We focus in particular on geometric morphisms between presheaf toposes induced by embeddings between small categories, and on geometric morphisms to the classifying topos of a geometric theory induced by a small category of set-based models of the theory. We also establish, in section 3.4, a general 'hom-tensor' adjunction between categories of $\mathcal{E}$ -valued functors (for $\mathcal{E}$ a Grothendieck topos) $[\mathcal{C}, \mathcal{E}]$ and $[\mathcal{D}, \mathcal{E}]$ induced by a functor $P : \mathcal{C} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$, which generalizes the well-known adjunction induced by Kan extensions along a given functor.

In section 4, in order to set up the field for the statement and proof of the characterization theorem, we identify some notable properties of theories of presheaf type, notably including the fact that every finitely presentable model of such a theory is finitely presented - in a strong sense which we make precise in section 4.3 - and admits an entirely syntactic description in terms of the signature of the theory and the notion of provability of geometric sequents over it in the theory (cf. section 4.1). We also show, in section 4.2, that for any geometric theory $\mathbb{T}$ and any Grothendieck topos $\mathcal{E}$, there exists for each pair of $\mathbb{T}$ -models M and N in $\mathcal{E}$, an 'object of $\mathbb{T}$ -model homomorphisms' in $\mathcal{E}$ from M to N which classifies the $\mathbb{T}$ -model homomorphisms in slices of $\mathcal{E}$ between the localizations of M and N in it.

In section 5, we establish our main characterization theorem providing necessary and sufficient conditions for a geometric theory to be classified by a presheaf topos. We first state the result abstractly and then proceed to obtain explicit reformulations of each of the conditions. We also derive some corollaries which allow to verify the satisfaction of the conditions of the theorem in specific situations which naturally arise in practice. Lastly, we show that, once recast in the language of indexed colimits of internal diagrams in toposes, the conditions of the characterization theorem for a given geometric theory $\mathbb{T}$ amount precisely to requirement that every model of $\mathbb{T}$ in any Grothendieck topos $\mathcal{E}$ should be a canonical $\mathcal{E}$ -indexed colimit of a certain $\mathcal{E}$ -filtered diagram of 'constant' finitely presentable models of $\mathbb{T}$ which are $\mathcal{E}$ -finitely presentable.

In section 6, we introduce the notion of faithful interpretation of geometric theories and investigate to what extent the satisfaction of the conditions of

the characterization theorem is preserved by this kind of interpretations. As applications of the general results that we obtain on this topic, we consider in particular the case of quotients of a given geometric theory $\mathbb{T}$ and that of injectivizations (i.e., theories obtained by adding, for each sort over the signature of the theory, a binary predicate which is provably complemented to the equality relation relative to that sort), providing various sufficient conditions for these theories to be of presheaf type if $\mathbb{T}$ is. To this end, we carry out in section 6.2 a general analysis of the relationship between finitely presentable and finitely generated models of a given geometric theory. In section 6.4.2, we treat the problem of finding a geometric theory classified by a given presheaf topos $[\mathcal{K}, \mathbf{Set}]$, and prove a general theorem ensuring that if the category $\mathcal{K}$ can be realized as a full subcategory of the category of finitely presentable models of a theory of presheaf type $\mathbb{T}$, there exists a quotient of $\mathbb{T}$ classified by the topos $[\mathcal{K}, \mathbf{Set}]$, which can be described in most explicit ways in terms of $\mathbb{T}$ and $\mathcal{K}$ provided that some natural conditions are satisfied. We also discuss, in section 6.4.1, the relationship between rigid topologies on the opposite of the category of finitely presentable models of a theory of presheaf type $\mathbb{T}$ and the presheaf-type quotients of $\mathbb{T}$.

In section 7, we investigate expansions of geometric theories from the point of view of the geometric morphisms that they induce between the respective classifying toposes. In particular, we introduce the notion of a localic (resp. hyperconnected) expansion, and show that it naturally corresponds to the notion of localic (resp. hyperconnected) geometric morphism at the level of classifying toposes; as a result, we obtain a logical characterization of the hyperconnected-localic factorization of a geometric morphism. Next, we address the problem of expanding a given geometric theory $\mathbb{T}$ to a theory classified by a presheaf topos of the form $[\mathcal{K}, \mathbf{Set}]$, where $\mathcal{K}$ is a small category of set-based models of $\mathbb{T}$, and describe a general method for defining such expansions.

In section 8, we discuss classical, as well as new, examples of theories of presheaf type from the perspective of the theory developed in the paper. We revisit in particular well-known examples of theories of presheaf type whose finitely presentable models are all finite, notably including the geometric theory of finite sets, and give fully constructive proofs of the fact that Mordijk's theory of abstract circles and Johnstone's theory of *Diers fields* are of presheaf type. Next, we introduce new examples of theories of presheaf type, including the theory of algebraic extensions of a given field, the theory of locally finite groups, the theory of vector spaces with linear independence predicates and the theory of abelian ℓ -groups with strong unit. We also show that the injectivization of the algebraic theory of groups is not of presheaf type and explicitly describe a presheaf completion for it.

2 Indexed colimits in toposes

2.1 Background on indexed categories

Before proceeding further, we need to recall some standard notions and facts from the theory of indexed categories; we refer the reader to [21] (especially sections B1.2, B2.3 and B3) and to [29] for the background.

Given an internal category $\mathbb{C}$ in a topos $\mathcal{E}$, we denote by $\mathbb{C}_1$ its object of arrows, by $\mathbb{C}_0$ its object of objects and by $d_0^{\mathbb{C}}, d_1^{\mathbb{C}} : \mathbb{C}_1 \rightarrow \mathbb{C}_0$ the domain and codomain arrows.

Given a small category $\mathcal{C}$ and a topos $\mathcal{E}$ defined over **Set**, we can always internalize $\mathcal{C}$ into $\mathcal{E}$ by means of the inverse image $\gamma_{\mathcal{E}}^*$ of the unique geometric morphism $\gamma_{\mathcal{E}} : \mathcal{E} \rightarrow \mathbf{Set}$ from $\mathcal{E}$ to **Set**; the resulting internal category in $\mathcal{E}$ will be denoted by the symbol $\underline{\mathbb{C}}$.

Indexed categories will be denoted by underlined letters, with possibly a subscript indicating the indexing category; the fibre at an object E of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}$ will be denoted by the symbol $\mathcal{A}_E$, and the functor $\mathcal{A}_E \rightarrow \mathcal{A}_{E'}$ corresponding to an arrow $e : E \rightarrow E'$ in $\mathcal{E}$ will be denoted by the symbol $\mathcal{A}_e$.

A $\mathcal{E}$ -indexed subcategory $\underline{\mathcal{B}}_{\mathcal{E}}$ of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ consists, for each object every $E \in \mathcal{E}$, of a subcategory $\mathcal{B}_E$ of the category $\mathcal{A}_E$ such that for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$ the functor $\mathcal{A}_{\alpha} : \mathcal{A}_E \rightarrow \mathcal{A}_{E'}$ restricts to the subcategories $\mathcal{B}_E$ and $\mathcal{B}_{E'}$. This clearly defines a $\mathcal{E}$ -indexed category $\underline{\mathcal{B}}_{\mathcal{E}}$ with a $\mathcal{E}$ -indexed inclusion $\underline{\mathcal{B}}_{\mathcal{E}} \hookrightarrow \underline{\mathcal{A}}_{\mathcal{E}}$.

A $\mathcal{E}$ -indexed functor $F : \underline{\mathcal{B}}_{\mathcal{E}} \rightarrow \underline{\mathcal{A}}_{\mathcal{E}}$ is said to be full if for every object E of $\mathcal{E}$ the functor $F_E : \mathcal{B}_E \rightarrow \mathcal{A}_E$ is full. A $\mathcal{E}$ -indexed subcategory $\underline{\mathcal{B}}_{\mathcal{E}}$ of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ is said to be a full $\mathcal{E}$ -indexed category of $\underline{\mathcal{A}}_{\mathcal{E}}$ if the associated $\mathcal{E}$ -indexed inclusion functor is full.

Every Grothendieck topos $\mathcal{E}$ gives rise to a $\mathcal{E}$ -indexed category $\underline{\mathcal{E}}_{\mathcal{E}}$ obtained by indexing $\mathcal{E}$ over itself.

Recall that if $\mathbb{C} = (d_0^{\mathbb{C}}, d_1^{\mathbb{C}} : \mathbb{C}_1 \rightarrow \mathbb{C}_0)$ is an internal category in a topos $\mathcal{E}$, a diagram of shape $\mathbb{C}$ in $\mathcal{E}$ is a pair $(f : F \rightarrow \mathbb{C}_0, \phi : \mathbb{C}_1 \times_{\mathbb{C}_0} F \rightarrow F)$ of arrows in $\mathcal{E}$ satisfying appropriate conditions, where the pullback $\mathbb{C}_1 \times_{\mathbb{C}_0} F$ is taken relatively to the arrow $d_0^{\mathbb{C}} : \mathbb{C}_1 \rightarrow \mathbb{C}_0$ and the arrow $f : F \rightarrow \mathbb{C}_0$:

$$\begin{array}{ccc} \mathbb{C}_1 \times_{\mathbb{C}_0} F & \xrightarrow{\pi_F} & F \\ \downarrow \pi_1 & & \downarrow f \\ \mathbb{C}_1 & \xrightarrow{d_0^{\mathbb{C}}} & \mathbb{C}_0 \end{array} .$$

For a Grothendieck topos $\mathcal{E}$ and an internal category $\mathbb{C}$ in $\mathcal{E}$, we have a $\mathcal{E}$ -indexed category $[\underline{\mathbb{C}}, \mathcal{E}]$, whose underlying category is the category $[\mathbb{C}, \mathcal{E}]$ of diagrams of shape $\mathbb{C}$ in $\mathcal{E}$ and morphisms between them.

Any internal category $\mathbb{C}$ in $\mathcal{E}$ gives naturally rise to a $\mathcal{E}$ -indexed category, which we call the $\mathcal{E}$ -externalization of $\mathbb{C}$ and denote by the symbol $\underline{\mathbb{C}}_{\mathcal{E}}$. The

category $[\mathbb{C}, \mathcal{E}]$ is equivalent (naturally in $\mathcal{E}$) to the category $[\underline{\mathbb{C}}_{\mathcal{E}}, \underline{\mathcal{E}}_{\mathcal{E}}]_{\mathcal{E}}$ of $\mathcal{E}$ -indexed functors $\underline{\mathbb{C}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ and indexed natural transformations between them (by Lemma B2.3.13 in [21]) and also to the category $[\mathcal{C}, \mathcal{E}]$ (by Corollary B2.3.14 in [21]). The equivalence between $[\mathbb{C}, \mathcal{E}]$ and $[\mathcal{C}, \mathcal{E}]$ restricts to an equivalence between the full subcategories $\mathbf{Tors}(\mathbb{C}, \mathcal{E})$ of $\mathbb{C}$ -torsors in $\mathcal{E}$ (as in section B3.2 of [21]) and $\mathbf{Flat}(\mathcal{C}, \mathcal{E})$ of flat functors $\mathcal{C} \rightarrow \mathcal{E}$ (as in chapter VII of [26]). For any internal functor between internal categories in $\mathcal{E}$ or internal diagram F in $\mathcal{E}$, we denote the corresponding $\mathcal{E}$ -indexed functor by the symbol $\underline{F}_{\mathcal{E}}$. For any functor $F : \mathcal{C} \rightarrow \mathcal{E}$, we denote the internal diagram in $[\mathbb{C}, \mathcal{E}]$ corresponding to it under the equivalence $[\mathcal{C}, \mathcal{E}] \simeq [\mathbb{C}, \mathcal{E}]$ by the symbol $\bar{F}$.

The discrete opfibration $p : \mathbb{F} \rightarrow \mathbb{C}$ over $\mathbb{C}$ corresponding to a diagram $(f : F \rightarrow \mathbb{C}_0, \phi : \mathbb{C}_1 \times_{\mathbb{C}_0} F \rightarrow F)$ of shape $\mathbb{C}$ in $\mathcal{E}$ is defined as follows: $\mathbb{F}_0 = F$, $\mathbb{F}_1 = \mathbb{C}_1 \times_{\mathbb{C}_0} F$, $d_0^{\mathbb{F}} = \pi_F : F_1 = \mathbb{C}_1 \times_{\mathbb{C}_0} F \rightarrow F = F_0$, $d_1^{\mathbb{F}} = \phi : F_1 = \mathbb{C}_1 \times_{\mathbb{C}_0} F \rightarrow F = F_0$, $p_0 = f : \mathbb{F}_0 = F \rightarrow \mathbb{C}_0$, $p_1 = \pi_1 : F_1 = \mathbb{C}_1 \times_{\mathbb{C}_0} F \rightarrow \mathbb{C}_1$. The discrete opfibration corresponding to a diagram $F \in [\mathbb{C}, \mathcal{E}]$ will also be denoted by $\pi_F^{opf} : \int^{opf} F \rightarrow \mathbb{C}$.

For internal diagrams $G \in [\mathbb{C}^{op}, \mathcal{E}]$ it is also natural to consider the *discrete fibration* corresponding to G , i.e. the opposite functor $\pi_G^{opf, op} : \int^{opf} G^{op} \rightarrow \mathbb{C}^{op, op} = \mathbb{C}$. We shall denote this functor by $\pi_G^f : \int^f G \rightarrow \mathbb{C}$.

For any internal category $\mathbb{C}$ in a topos $\mathcal{E}$, the category $[\mathbb{C}, \mathcal{E}]$ of diagrams of shape $\mathbb{C}$ in $\mathcal{E}$ is equivalent to the category $\mathbf{DOpf}/\mathbb{C}$ of *discrete opfibrations* over $\mathbb{C}$ (cf. Proposition B2.5.3 [21]); we shall denote this equivalence by

$$\tau_{\mathcal{E}}^{\mathbb{C}} : [\mathbb{C}, \mathcal{E}] \rightarrow \mathbf{DOpf}/\mathbb{C}.$$

A diagram of shape $\mathbb{C}$ in $\mathcal{E}$ lies in the subcategory $\mathbf{Tors}(\mathbb{C}, \mathcal{E})$ of $[\mathbb{C}, \mathcal{E}]$ if and only if the domain of the corresponding discrete opfibration is a filtered internal category in $\mathcal{E}$.

Any internal functor $H : \mathbb{C} \rightarrow \mathbb{D}$ between internal categories $\mathbb{C}$ and $\mathbb{D}$ in a topos $\mathcal{E}$ induces a functor $[\mathbb{D}, \mathcal{E}] \rightarrow [\mathbb{C}, \mathcal{E}]$, denoted $D \mapsto D \circ H$, which corresponds, at the level of $\mathcal{E}$ -indexed categories, to the composition functor with the indexed functor corresponding to H , and, at the level of discrete opfibrations associated to the internal diagrams, as the pullback of them along the functor H . The latter pullbacks in the category of internal categories in $\mathcal{E}$ are computed ‘pointwise’ as pullbacks in $\mathcal{E}$, and they are preserved by the dualizing functor $\mathbb{C} \rightarrow \mathbb{C}^{op}$.

Let us recall from [29] the notion of (indexed) colimit of a $\mathcal{E}$ -indexed functor, where $\mathcal{E}$ is a cartesian category. We shall denote by $!_I$ the unique arrow from an object I of $\mathcal{E}$ to the terminal object 1 of $\mathcal{E}$. Given a $\mathcal{S}$ -indexed functor $\Gamma : \underline{X} \rightarrow \underline{A}$ and an object A of A_1 , we denote by $\Delta_{\underline{A}}^{\underline{X}} A$ the $\mathcal{E}$ -indexed functor $\underline{X} \rightarrow \underline{A}$ assigning to any $E \in \mathcal{E}$ the constant functor on X_E with value $\mathcal{A}_{!_I}(A)$.

Let $\underline{X}$ be any indexed category and $\Gamma : \underline{X} \rightarrow \underline{A}$ any indexed functor.

An indexed cocone $\mu : \Gamma \rightarrow A$ consists of an object A in A_1 together with an indexed natural transformation $\mu : \Gamma \rightarrow \Delta_{\underline{A}}^X A$, i.e. for each I in $\mathcal{E}$ an ordinary cocone $\mu_I : \Gamma_I \rightarrow \Delta(\mathcal{A}_I(A))$ such that for each arrow $\alpha : J \rightarrow I$ in $\mathcal{E}$, $\alpha^* \circ \mu_I = \mu_J \circ \alpha^*$. If $\mu : \Gamma \rightarrow \Delta_{\underline{A}}^X A$ is a universal such cone we say that it is a colimit cone over the indexed functor Γ with vertex A . If, furthermore, for any object I of $\mathcal{E}$, the localization $\mu/I : \Gamma/I \rightarrow (\Delta_{\underline{A}}^X A)/I = \Delta_{\underline{A}/I}^{X/I}(\mathcal{A}_I(A))$ of μ at I is a colimit cone we say that μ is the indexed colimit of Γ/I .

We can describe this notion more explicitly in the particular case of $\mathcal{E}$ -indexed functors with values in the $\mathcal{E}$ -indexed category $\underline{\mathcal{E}}_{\mathcal{E}}$. Let us suppose that $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -indexed category and $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ is an indexed functor. A cocone μ over D consists of an object U of $\mathcal{E}$ and, for each object E of $\mathcal{E}$, of a cocone $\mu_E : D_E \rightarrow \Delta(!_E^*(U))$ over the diagram $D_E : \mathcal{A}_E \rightarrow \mathcal{E}/E$ such that for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$ we have $\alpha^*(\mu_E(c)) = \mu_{E'}(\mathcal{A}_{\alpha}(c))$ for all $c \in \mathcal{A}_E$, that is $\alpha^* \mu_E = \mu_{E'} \mathcal{A}_{\alpha}$ as arrows $D_{E'}(\mathcal{A}_{\alpha}(c)) = \alpha^*(D_E(c)) \rightarrow !_E^*(U) = \alpha^*(!_E^*(U))$ in $\mathcal{E}/E'$, where $\alpha^* : \mathcal{E}/E \rightarrow \mathcal{E}/E'$ is the pullback functor (notice that $D_{E'} \circ \mathcal{A}_{\alpha} = \alpha^* \circ D_E$ since D is a $\mathcal{E}$ -indexed functor).

Note that if $\underline{\mathcal{A}}_{\mathcal{E}}$ is the $\mathcal{E}$ -externalization of an internal category $\mathbb{C}$ in $\mathcal{E}$ and $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ is an indexed functor corresponding to an internal diagram $D \in [\mathbb{C}, \mathcal{E}]$, the discrete opfibration associated to the internal diagram in $[\mathbb{C}, \mathcal{E}]$ corresponding to the localization $D/E : \underline{\mathcal{A}}_{\mathcal{E}}/E \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}/E \cong \underline{\mathcal{E}}/E_{\mathcal{E}/E}$ is the image of the discrete opfibration associated to D under the pullback functor $!_E^* : \mathcal{E} \rightarrow \mathcal{E}/E$ along the unique arrow $!_E : E \rightarrow 1_{\mathcal{E}}$.

The colimit $\text{colim}_{\mathcal{E}}(D)$ in $\mathcal{E}$ of an internal diagram $D \in [\mathbb{C}, \mathcal{E}]$, where $\mathbb{C}$ is an internal category in $\mathcal{E}$, is defined to be the coequalizer of the two arrows $d_0^{\int^{opf} D}, d_1^{\int^{opf} D}$ (recall that $\int^{opf} D$ is the domain of the discrete opfibration over $\mathbb{C}$ corresponding to the diagram D).

For any internal diagram $G \in [\mathbb{F}, \mathcal{E}]$, its colimit $\text{colim}_{\mathcal{E}}(G)$ is isomorphic to the $\mathcal{E}$ -indexed colimit $\underline{\text{colim}}_{\mathcal{E}}(\underline{G}_{\mathcal{E}})$ of the $\mathcal{E}$ -indexed functor $\underline{G}_{\mathcal{E}} : \mathbb{F}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ corresponding to it under the equivalence $[\mathbb{F}, \mathcal{E}] \simeq [\mathbb{F}_{\mathcal{E}}, \underline{\mathcal{E}}_{\mathcal{E}}]_{\mathcal{E}}$.

If $D \in [\mathbb{C}, \mathcal{E}]$ is an internal diagram in $\mathcal{E}$ with colimit $\text{colim}_{\mathcal{E}}(D) = \text{coeq}(d_1^{\int^{opf} D}, d_1^{\int^{opf} D})$, the colimiting $\mathcal{E}$ -indexed cocone $(\text{colim}_{\mathcal{E}}(D), \mu)$ of the corresponding $\mathcal{E}$ -indexed functor $\underline{D}_{\mathcal{E}} : \mathbb{C}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ can be described as follows. Let us denote by c the canonical coequalizer arrow $(\int^{opf} D)_0 \rightarrow \text{coeq}(d_0^{\int^{opf} D}, d_1^{\int^{opf} D})$ in $\mathcal{E}$. For any object E of $\mathcal{E}$ and any object $x : E \rightarrow \mathbb{C}_0$ of the category $\mathbb{C}_{\mathcal{E}}E$, the arrow $\mu_E(x) : D_E(x) = r_x \rightarrow !_E^*(\text{colim}_{\mathcal{E}}(D)) = \text{coeq}(d_0^{\int^{opf} D}, d_1^{\int^{opf} D}) \times E$ is equal to $\langle c \circ z_x, r_x \rangle$, where the arrow z_x is defined by the following pullback diagram:

$$\begin{array}{ccc} R_x & \xrightarrow{z_x} & (\int^{opf} D)_0 \\ \downarrow r_x & & \downarrow (\pi_D^{opf})_0 \\ E & \xrightarrow{x} & \mathbb{C}_0. \end{array}$$

2.2 $\mathcal{E}$ -filtered indexed categories

The following definition will be important in the sequel.

Definition 2.1. Let $\mathcal{E}$ be a Grothendieck topos and $\underline{\mathcal{A}}$ be a $\mathcal{E}$ -indexed category. We say that $\underline{\mathcal{A}}$ is $\mathcal{E}$ -filtered if the following conditions are satisfied:

- (a) For any object E of $\mathcal{E}$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object b_i of the category $\mathcal{A}_{E_i}$;
- (b) For any $E \in \mathcal{E}$ and any objects a and b of the category $\mathcal{A}_E$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object c_i of the category $\mathcal{A}_{E_i}$ and arrows $f_i : \mathcal{A}_{e_i}(a) \rightarrow c_i$ and $g_i : \mathcal{A}_{e_i}(b) \rightarrow c_i$ in the category $\mathcal{A}_{E_i}$;
- (c) For any object E of $\mathcal{E}$ and any two parallel arrows $u, v : a \rightarrow b$ in the category $\mathcal{A}_E$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object c_i of the category $\mathcal{A}_{E_i}$ and an arrow $w_i : \mathcal{A}_{e_i}(b) \rightarrow c_i$ in $\mathcal{A}_{E_i}$ such that $w_i \circ \mathcal{A}_{e_i}(u) = w_i \circ \mathcal{A}_{e_i}(v)$.

The $\mathcal{E}$ -externalization of any internal filtered category in $\mathcal{E}$ is $\mathcal{E}$ -filtered, but it is not true that if the externalization of an internal category $\mathbb{C}$ in $\mathcal{E}$ is $\mathcal{E}$ -filtered then $\mathbb{C}$ is filtered as an internal category in $\mathcal{E}$.

A standard example of indexed $\mathcal{E}$ -filtered categories is provided by indexed categories of elements of flat functors with values in $\mathcal{E}$, in the sense of the following definition.

Definition 2.2. Let $P : \mathcal{C}^{op} \rightarrow \mathcal{E}$ be a functor. The $\mathcal{E}$ -indexed category of elements $\int P_{\mathcal{E}}$ of P assigns to any object E of $\mathcal{E}$ the category $\int P_E$ whose objects are the pairs (c, x) where c is an object of $\mathcal{C}$ and $x : E \rightarrow F(c)$ is an arrow in $\mathcal{E}$, and whose arrows $(c, x) \rightarrow (d, y)$ are the arrows $f : c \rightarrow d$ in $\mathcal{C}$ such that $F(f) \circ y = x$, and to any arrow $e : E' \rightarrow E$ in $\mathcal{E}$ the functor $\int P_e : \int P_E \rightarrow \int P_{E'}$ sending any object (c, x) of $\int P_E$ to the object $(c, x \circ e)$ of $\int P_{E'}$ and acting accordingly on the arrows.

Proposition 2.3. Let $F : \mathcal{C}^{op} \rightarrow \mathcal{E}$ be a flat functor. Then the indexed category $\int F_{\mathcal{E}}$ is $\mathcal{E}$ -filtered.

Proof Straightforward from the characterization of flat functors as filtering functors given in chapter VII of [26]. $\square$

2.3 Indexation of internal diagrams

Given an internal diagram $D \in [\mathbb{C}, \mathcal{E}]$, the corresponding $\mathcal{E}$ -indexed functor $\overline{D}_{\mathcal{E}} : \underline{\mathbb{C}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ can be described as follows. For any object E of $\mathcal{E}$, $D_E : \mathbb{C}_E \rightarrow \mathcal{E}/E$ sends any object $x : E \rightarrow \mathbb{C}_0$ of $\mathbb{C}_E$ to the object $r_x : R_x \rightarrow E$ of $\mathcal{E}/E$ obtained by pulling $(\pi_F^{opf})_0 : (\int^{opf} F)_0 \rightarrow \mathbb{C}_0$ back along x , and any

arrow $h : E \rightarrow \mathbb{C}_1$ of $\mathbb{C}_E$ from $x : E \rightarrow \mathbb{C}_0$ to $x' : E \rightarrow \mathbb{C}_0$ to the arrow $D_E(h) : r_x \rightarrow r_{x'}$ in $\mathcal{E}/E$ defined as follows. Consider the pullback squares

$$\begin{array}{ccc} S & \xrightarrow{u} & (\int^{opf} D)_1 \\ \downarrow s & & \downarrow (\pi_D^{opf})_1 \\ E & \xrightarrow{h} & \mathbb{C}_1 \end{array}$$

and

$$\begin{array}{ccc} R_{x'} & \xrightarrow{z_{x'}} & (\int^{opf} D)_0 \\ \downarrow r_{x'} & & \downarrow (\pi_D^{opf})_0 \\ E & \xrightarrow{x'} & \mathbb{C}_0 . \end{array}$$

Since $d_1^{\mathbb{C}} \circ h = x'$ we have that $(\pi_D^{opf})_0 \circ d_1^{\int^{opf} D} \circ u = x' \circ s$ and hence by the universal property of the first pullback square, there exists a unique arrow $\beta : S \rightarrow R_{x'}$ such that $r_{x'} \circ \beta = s$ and $z_{x'} \circ \beta = d_1^{\int^{opf} D} \circ u$.

As $\pi_F^{opf} : \int^{opf} F \rightarrow \mathbb{C}$ is a discrete opfibration, the diagram

$$\begin{array}{ccc} (\int^{opf} D)_1 & \xrightarrow{d_0^{\int^{opf} D}} & (\int^{opf} D)_0 \\ \downarrow (\pi_D^{opf})_1 & & \downarrow (\pi_D^{opf})_0 \\ \mathbb{C}_1 & \xrightarrow{d_0^{\mathbb{C}}} & \mathbb{C}_0 \end{array}$$

is a pullback. ‘Composing’ it with the first pullback square thus yields a pullback square

$$\begin{array}{ccc} S & \xrightarrow{d_0^{\int^{opf} D} \circ u} & (\int^{opf} D)_0 \\ \downarrow s & & \downarrow (\pi_D^{opf})_0 \\ E & \xrightarrow{d_0^{\mathbb{C}} \circ h} & \mathbb{C}_0 . \end{array}$$

Now, since $(\pi_D^{opf})_0 \circ z_x = x \circ r_x = d_0^{\mathbb{C}} \circ h \circ r_x$, the universal property of this pullback square provides a unique arrow $\gamma : R_x \rightarrow S$ such that $s \circ \gamma = r_x$ and $d_0^{\int^{opf} D} \circ u \circ \gamma = z_x$. We define $D_E(h) : r_x \rightarrow r_{x'}$ in $\mathcal{E}/E$ to be equal to the composite arrow $\beta \circ \gamma : R_x \rightarrow R_{x'}$.

2.4 Colimits and tensor products

For any internal category $\mathbb{F}$ in $\mathcal{E}$, we denote by $\text{coeq}(\mathbb{F})$ the coequalizer of the two arrows $d_0^{\mathbb{F}}$ and $d_1^{\mathbb{F}}$.

From the above discussion it follows that for any $D, P \in [\mathbb{C}, \mathcal{E}]$, $\text{colim}_{\mathcal{E}}(P \circ \pi_D^{\text{opf}}) \cong \text{colim}_{\mathcal{E}}(D \circ \pi_P^{\text{opf}})$. Indeed, if we consider the pullback square

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{q^P} & \int^{\text{opf}} P \\ \downarrow q^D & & \downarrow \pi_P^{\text{opf}} \\ \int^{\text{opf}} D & \xrightarrow{\pi_D^{\text{opf}}} & \mathbb{C} \end{array}$$

in the category $\mathbf{cat}(\mathcal{E})$ of internal categories in $\mathcal{E}$, we have that $\text{colim}_{\mathcal{E}}(P \circ \pi_D^{\text{opf}}) \cong \text{coeq}(\text{dom}(q^D)) = \text{coeq}(\mathbb{R}) = \text{coeq}(\text{dom}(q^P)) \cong \text{colim}_{\mathcal{E}}(D \circ \pi_P^{\text{opf}})$.

Similarly, by exploiting the fact that the operation $\mathbb{F} \rightarrow \text{coeq}(\mathbb{F})$ on internal categories in $\mathcal{E}$ is invariant under the dualization functor $\mathbb{F} \rightarrow \mathbb{F}^{\text{op}}$, we obtain another commutation result, which we shall use in the sequel: for any internal diagram $F \in [\mathbb{C}^{\text{op}}, \mathcal{E}]$ and any internal diagram $P \in [\mathbb{C}, \mathcal{E}]$, we have a natural isomorphism $\text{colim}_{\mathcal{E}}(P \circ \pi_F^f) \cong \text{colim}_{\mathcal{E}}(F \circ \pi_P^f)$. To prove this, we consider the following pullback squares:

$$\begin{array}{ccc} \mathbb{R} & \xrightarrow{q^P} & \int^{\text{opf}} P \\ \downarrow q^F & & \downarrow \pi_P^{\text{opf}} \\ \int^f F & \xrightarrow{\pi_F^f} & \mathbb{C} \end{array} \qquad \begin{array}{ccc} \mathbb{S} & \xrightarrow{r^F} & \int^{\text{opf}} F \\ \downarrow r^P & & \downarrow \pi_F^{\text{opf}} \\ \int^f P & \xrightarrow{\pi_P^f} & \mathbb{C}^{\text{op}} \end{array}$$

We have that

$$\text{colim}_{\mathcal{E}}(P \circ \pi_F^f) \cong \text{coeq}(\text{dom}(q^F)) = \text{coeq}(\mathbb{R}),$$

while

$$\text{colim}_{\mathcal{E}}(F \circ \pi_P^f) = \text{coeq}(\text{dom}(r^P)) = \text{coeq}(\mathbb{S}).$$

But the fact that the dualization functor preserves pullbacks in $\mathbf{cat}(\mathcal{E})$ implies that $\mathbb{S} \simeq \mathbb{R}^{\text{op}}$, whence $\text{coeq}(\mathbb{R}) \cong \text{coeq}(\mathbb{S})$, as required.

Summarizing, we have the following

Proposition 2.4. *Let $\mathbb{C}$ be an internal category in a topos $\mathcal{E}$, P and D internal diagrams in $[\mathbb{C}, \mathcal{E}]$ and F an internal diagram in $[\mathbb{C}^{\text{op}}, \mathcal{E}]$. Then we have natural isomorphisms*

- (i) $\text{colim}_{\mathcal{E}}(P \circ \pi_D^{\text{opf}}) \cong \text{colim}_{\mathcal{E}}(D \circ \pi_P^{\text{opf}})$;
- (ii) $\text{colim}_{\mathcal{E}}(P \circ \pi_F^f) \cong \text{colim}_{\mathcal{E}}(F \circ \pi_P^f)$.

□

Let us now proceed to applying this proposition in the context of a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$, where $\mathcal{C}$ is a small category and $\mathcal{E}$ is a Grothendieck topos, and a functor $P : \mathcal{C} \rightarrow \mathbf{Set}$. To this end, we explicitly describe the discrete opfibration corresponding to the diagram of shape $\mathbb{C}$ in $\mathcal{E}$, where $\mathbb{C}$ is the internalization of $\mathcal{C}$ in $\mathcal{E}$, associated to a functor $G : \mathcal{C} \rightarrow \mathcal{E}$. We denote by $Ob(\mathcal{C})$ the set of objects of $\mathcal{C}$ and by $Arr(\mathcal{C})$ the set of arrows of $\mathcal{C}$.

We have that $F = \coprod_{c \in \mathcal{C}} G(c)$ and that $f : F = \coprod_{c \in \mathcal{C}} G(c) \rightarrow \mathbb{C}_0 = \coprod_{c \in \mathcal{C}} 1_{\mathcal{E}}$ is equal to $\coprod_{c \in \mathcal{C}} !_{G(c)}$, where $!_{G(c)}$ is the unique arrow $G(c) \rightarrow 1_{\mathcal{E}}$ in $\mathcal{E}$ (for any $c \in \mathcal{C}$).

Let $J_f : G(dom(f)) \rightarrow \coprod_{f \in Arr(\mathcal{C})} G(dom(f))$ (resp. $\mu_c : G(c) \rightarrow \coprod_{c \in Ob(\mathcal{C})} G(c)$), $\kappa_f : 1_{\mathcal{E}} \rightarrow \coprod_{f \in Arr(\mathcal{C})} 1_{\mathcal{E}}$, $\lambda_c : 1_{\mathcal{E}} \rightarrow \coprod_{c \in Ob(\mathcal{C})} 1_{\mathcal{E}}$ be the canonical coproduct arrows.

First, let us show that the diagram

$$\begin{array}{ccc} \coprod_{f \in Arr(\mathcal{C})} G(dom(f)) & \xrightarrow{d_0^G} & \coprod_{c \in Ob(\mathcal{C})} G(c) \\ \downarrow \coprod_{f \in Arr(\mathcal{C})} !_{G(dom(f))} & & \downarrow \coprod_{c \in Ob(\mathcal{C})} !_{G(c)} \\ \coprod_{f \in Arr(\mathcal{C})} 1_{\mathcal{E}} & \xrightarrow{d_0^{\mathbb{C}}} & \coprod_{c \in Ob(\mathcal{C})} 1_{\mathcal{E}} . \end{array}$$

is a pullback, where the arrow d_0^G is defined by setting $d_0^G \circ J_f = \mu_{dom(f)}$ (for any $f \in Arr(\mathcal{C})$). We have to prove that, for any object E of $\mathcal{E}$ and arrows $\alpha : E \rightarrow \coprod_{f \in Arr(\mathcal{C})} 1_{\mathcal{E}}$ and $\beta : E \rightarrow \coprod_{c \in Ob(\mathcal{C})} G(c)$ such that $\coprod_{c \in Ob(\mathcal{C})} !_{G(c)} \circ \beta = d_0^{\mathbb{C}} \circ \alpha$, there exists a unique arrow $\gamma : E \rightarrow \coprod_{f \in Arr(\mathcal{C})} !_{G(dom(f))}$ such that $\alpha = \coprod_{f \in Arr(\mathcal{C})} G(dom(f)) \circ \gamma$ and $\beta = d_0^G \circ \gamma$. To this end, consider, for any $c \in Ob(\mathcal{C})$ and $f \in Arr(\mathcal{C})$, the commutative diagram

$$\begin{array}{ccccc} E_{(f,c)} & \xrightarrow{q_f} & E_f & \xrightarrow{\alpha_f} & 1_{\mathcal{E}} \\ \downarrow p_c & & \downarrow y_f & & \downarrow \kappa_f \quad \searrow id \\ E_c & \xrightarrow{z_c} & E & \xrightarrow{\alpha} & \coprod_{f \in Arr(\mathcal{C})} 1_{\mathcal{E}} \\ \downarrow \beta_c & & \downarrow \beta & & \downarrow d_0^{\mathbb{C}} \quad \nearrow \lambda_{dom(f)} \\ G(c) & \xrightarrow{\kappa_c} & \coprod_{c \in Ob(\mathcal{C})} G(c) & \xrightarrow{\coprod_{c \in Ob(\mathcal{C})} !_{G(c)}} & \coprod_{c \in Ob(\mathcal{C})} 1_{\mathcal{E}} \\ & \searrow !_{G(c)} & & \nearrow \lambda_c & \\ & & 1_{\mathcal{E}} & & \end{array}$$

where all the squares except for the lower-right one are pullbacks. The commutativity of the diagram, combined with the fact that distinct coproduct arrows are disjoint from each other, immediately implies that for any pair (f, c) such that $c \neq \text{dom}(f)$, we have $E_{(f,c)} \cong 0_{\mathcal{E}}$. On the other hand, the stability of coproducts under pullbacks implies that $E \cong \coprod_{c \in \text{Ob}(\mathcal{C})} E_c$, and

$$E \cong \coprod_{f \in \text{Arr}(\mathcal{C})} E_f; \text{ whence } E \cong \coprod_{(f,c) \in \text{Arr}(\mathcal{C}) \times \text{Ob}(\mathcal{C})} E_{(f,c)} \cong \coprod_{(f,c) \mid \text{dom}(f)=c} E_{(f,c)},$$

with canonical coproduct arrows $\xi_{(f,c)} = y_f \circ q_f = z_c \circ p_c : E_{(f,c)} \rightarrow E$. We define, for each pair (f, c) such that $c = \text{dom}(f)$, the arrow $\gamma_{(f,c)} : E_{(f,c)} \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} G(\text{dom}(f))$ as the composite $J_f \circ \beta_c \circ p_c$, and set γ equal to the arrow

$$\gamma : E \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} G(\text{dom}(f)).$$

We have to verify that $\alpha = \coprod_{f \in \text{Arr}(\mathcal{C})} !_{G(\text{dom}(f))} \circ \gamma$ and $\beta = d_0^G \circ \gamma$ or, equivalently, that for any pair (f, c) such that $\text{dom}(f) = c$, we have:

$$(1) \quad \alpha \circ \xi_{(f,c)} = \coprod_{f \in \text{Arr}(\mathcal{C})} !_{G(\text{dom}(f))} \circ \gamma_{(f,c)} \text{ and}$$

$$(2) \quad \beta \circ \xi_{(f,c)} = d_0^G \circ \gamma_{(f,c)}.$$

To prove (1), we preliminarily show that for any pair (f, c) such that $c = \text{dom}(f)$, we have $\alpha_f \circ q_f = !_{G(c)} \circ \beta_c \circ p_c$. Since the arrow $\lambda_c = \lambda_{\text{dom}(f)} : 1_{\mathcal{E}} \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} 1_{\mathcal{E}}$ is monic, it is equivalent to prove that $\lambda_{\text{dom}(f)} \circ \alpha_f \circ q_f = \lambda_c \circ !_{G(c)} \circ \beta_c \circ p_c$. But the commutativity of the above diagram yields $\lambda_{\text{dom}(f)} \circ \alpha_f \circ q_f = d_0^{\mathcal{C}} \circ \kappa_f \circ \alpha_f \circ q_f = d_0^{\mathcal{C}} \circ \alpha \circ y_f \circ q_f = \coprod_{c \in \text{Ob}(\mathcal{C})} !_{G(c)} \circ \beta \circ y_f \circ q_f = \coprod_{c \in \text{Ob}(\mathcal{C})} !_{G(c)} \circ \beta \circ z_c \circ p_c = \coprod_{c \in \text{Ob}(\mathcal{C})} !_{G(c)} \circ \mu_c \circ \beta_c \circ p_c = \lambda_c \circ !_{G(c)} \circ \beta_c \circ p_c$, as required.

We thus have $\alpha \circ \xi_{(f,c)} = \alpha \circ y_f \circ q_f = \kappa_f \circ \alpha_f \circ q_f = \kappa_f \circ !_{G(c)} \circ \beta_c \circ p_c = \coprod_{f \in \text{Arr}(\mathcal{C})} !_{G(\text{dom}(f))} \circ \gamma_{(f,c)}$. This proves condition (1).

Further, $\beta \circ \xi_{(f,c)} = \beta \circ z_c \circ p_c = \mu_c \circ \beta_c \circ p_c = \mu_{\text{dom}(f)} \circ \beta_c \circ p_c = d_0^G \circ J_f \circ \beta_c \circ p_c = d_0^G \circ \gamma_{(f,c)}$. This proves condition (2).

Let us define the arrow

$$d_1^G : \coprod_{f \in \text{Arr}(\mathcal{C})} G(\text{dom}(f)) \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} G(c)$$

by setting, for each $f \in \text{Arr}(\mathcal{C})$, $d_1^G \circ J_f = \mu_{\text{cod}(f)} \circ G(f)$.

The discrete opfibration $p : \mathbb{F} \rightarrow \mathbb{C}$ corresponding to G can be described as follows: $\mathbb{F}_0 = \coprod_{c \in \text{Ob}(\mathcal{C})} G(c)$, $\mathbb{F}_1 = \coprod_{f \in \text{Arr}(\mathcal{C})} G(\text{dom}(f))$, the domain and codomain arrows $d_0^{\mathbb{F}}, d_1^{\mathbb{F}} : \mathbb{F}_1 \rightarrow \mathbb{F}_0$ are respectively equal to d_0^G and to

$d_1^G, p_0 : \mathbb{F}_0 \rightarrow \mathbb{C}_0$ is equal to $\coprod_{c \in \text{Ob}(\mathcal{C})} !_{G(c)}$ and $p_1 : \mathbb{F}_1 \rightarrow \mathbb{C}_1$ is equal to $\coprod_{f \in \text{Arr}(\mathcal{C})} !_{G(\text{dom}(f))}$. The composition law in the internal category $\mathbb{F}$ is defined in the obvious way.

We leave to the reader the straightforward task of verifying that this is indeed the discrete opfibration corresponding to the functor G via the composite of the equivalence $[\mathcal{C}, \mathcal{E}] \simeq [\mathbb{C}, \mathcal{E}]$ with the equivalence

$$\tau_{\mathcal{E}}^{\mathbb{C}} : [\mathbb{C}, \mathcal{E}] \rightarrow \mathbf{DOpf}/\mathbb{C}.$$

Recalling that, for any functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$, the discrete fibration $\pi_F^f : \int^f F \rightarrow \mathbb{C}$ associated to it is equal to $\pi_F^{\text{opf}^{\text{op}}} : \int^{\text{opf}^{\text{op}}} F^{\text{op}} \rightarrow \mathbb{C}^{\text{op}^{\text{op}}} = \mathbb{C}$, we deduce the following explicit description of the discrete fibration $\pi_F^f : \int^f F \rightarrow \mathbb{C}$ associated to a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$: $(\int^f F)_0 = \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$, $(\int^f F)_1 = \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$, the domain and codomain arrows $d_0^F, d_1^F : (\int^f F)_1 \rightarrow (\int^f F)_0$ are defined by the conditions $d_0^F \circ J_f = \mu_{\text{cod}(f)}$ and $d_1^F \circ J_f = \mu_{\text{dom}(f)} \circ F(f)$ for all $f \in \text{Arr}(\mathcal{C})$ (where $\mu_c : F(c) \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$ and $J_f : F(\text{cod}(f)) \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$ are the canonical coproduct arrows), $(\pi_F^f)_0 : (\int^f F)_0 \rightarrow \mathbb{C}_0$ is equal to $\coprod_{c \in \text{Ob}(\mathcal{C})} !_{F(c)}$ and $(\pi_F^f)_1 : (\int^f F)_1 \rightarrow \mathbb{C}_1$ is equal to $\coprod_{f \in \text{Arr}(\mathcal{C})} !_{F(\text{cod}(f))}$. The composition law in the internal category $\int^f F$ is defined in the obvious way.

Theorem 2.5. *Let $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ be a functor from the opposite of a small category $\mathcal{C}$ to a Grothendieck topos $\mathcal{E}$ and $P : \mathcal{C} \rightarrow \mathbf{Set}$ be a functor. Then the following three objects are naturally isomorphic:*

- (i) $\text{colim}(F \circ \pi_P^f)$
- (ii) $\text{colim}_{\mathcal{E}}(\overline{F} \circ \pi_{P_{\mathcal{E}}}^f) \cong \text{colim}_{\mathcal{E}}(P_{\mathcal{E}} \circ \pi_{\overline{F}}^f)$ (cf. Proposition 2.4)
- (iii) $\underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \underline{\pi_{\overline{F}}^f})$,

where $P_{\mathcal{E}}$ is the internal diagram in $[\mathbb{C}, \mathcal{E}]$ given by $\overline{\gamma_{\mathcal{E}}^* \circ P}$.

Proof The isomorphism between $\text{colim}_{\mathcal{E}}(P_{\mathcal{E}} \circ \pi_{\overline{F}}^f)$ and $\underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \underline{\pi_{\overline{F}}^f})$ follows from the general fact that for any internal diagram $G \in [\mathbb{D}, \mathcal{E}]$ its colimit $\text{colim}_{\mathcal{E}}(G)$ is isomorphic to the $\mathcal{E}$ -indexed colimit $\underline{\text{colim}}_{\mathcal{E}}(\underline{G_{\mathcal{E}}})$. It thus remains to prove the isomorphism between $\text{colim}(F \circ \pi_P^f)$ and $\text{colim}_{\mathcal{E}}(\overline{F} \circ \pi_{P_{\mathcal{E}}}^f)$. To this end, we recall the following three general facts:

- (1) For any functor $G : \mathcal{D} \rightarrow \mathcal{E}$, its colimit $\text{colim}(G)$ is naturally isomorphic to the colimit $\text{colim}_{\mathcal{E}}(\overline{G})$;
- (2) For any functor $H : \mathcal{D} \rightarrow \mathcal{C}$ between small categories $\mathcal{C}$ and $\mathcal{D}$ and any functor $M : \mathcal{C} \rightarrow \mathcal{E}$, we have a natural isomorphism $\overline{F \circ H} \cong \overline{F} \circ \gamma_{\mathcal{E}}^*(H)$;
- (3) For any geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$, the diagram

$$\begin{array}{ccc}
[\mathbb{C}, \mathcal{E}] & \xrightarrow{\tau_{\mathcal{E}}^{\mathbb{C}}} & \mathbf{DOpf}/\mathbb{C} \\
\downarrow f^*(-) & & \downarrow f^*(-) \\
[f^*(\mathbb{C}), \mathcal{E}] & \xrightarrow{\tau_{\mathcal{F}}^{f^*(\mathbb{C})}} & \mathbf{DOpf}/f^*(\mathbb{C})
\end{array}$$

commutes.

We therefore have that

$$\text{colim}(F \circ \pi_P^f) \cong \text{colim}_{\mathcal{E}}(\overline{F \circ \pi_P^f}) \cong \text{colim}_{\mathcal{E}}(\overline{F} \circ \gamma_{\mathcal{E}}^*(\pi_P^f)) \cong \text{colim}_{\mathcal{E}}(\overline{F} \circ \pi_{P_{\mathcal{E}}}^f),$$

where the first isomorphism follows from (1), the second from (2) and the third from (3). $\square$

We shall indicate the three isomorphic objects of the theorem by the symbol $F \otimes_{\mathcal{C}} P$.

The following lemma is essentially contained in the proof of Giraud's theorem (cf. [18] or, for instance, the Appendix of [26]), but we were not able to find its exact statement in the literature; we thus provide a proof of it for the reader's convenience.

Lemma 2.6. *Let $\mathcal{E}$ be a Grothendieck topos and $\{e_i : E_i \rightarrow E \mid i \in I\}$ an epimorphic family in $\mathcal{E}$. Then the arrow $\coprod_{i \in I} e_i : \coprod_{i \in I} E_i \rightarrow E$ yields an isomorphism $(\coprod_{i \in I} E_i)/R \cong E$, where R is the equivalence relation in $\mathcal{E}$ on the object $\coprod_{i \in I} E_i$ given by the subobject $\coprod_{(i,j) \in I \times I} E_i \times_E E_j \hookrightarrow \coprod_{(i,j) \in I \times I} E_i \times E_j \cong \coprod_{i \in I} E_i \times \coprod_{j \in I} E_j$; in particular, the arrow $\coprod_{i \in I} e_i : \coprod_{i \in I} E_i \rightarrow E$ is the coequalizer in $\mathcal{E}$ of the two canonical arrows $\coprod_{(i,j) \in I \times I} E_i \times_E E_j \rightarrow \coprod_{i \in I} E_i$.*

Proof Let R be the kernel pair of the epimorphism $\coprod_{i \in I} e_i$, that is the pullback of this arrow along itself; then, by the well-known exactness properties of Grothendieck toposes, R is an equivalence relation on $\coprod_{i \in I} E_i$ such that the coequalizer in $\mathcal{E}$ of the two associated projections is isomorphic to q . Now, the fact that pullbacks preserve coproducts in a Grothendieck topos implies that R is isomorphic to the subobject $\coprod_{(i,j) \in I \times I} E_i \times_E E_j \hookrightarrow \coprod_{(i,j) \in I \times I} E_i \times E_j \cong \coprod_{i \in I} E_i \times \coprod_{j \in I} E_j$, as required. $\square$

Remark 2.7. The lemma admits an obvious generalization to arbitrary separating sets for the topos $\mathcal{E}$ (cf. the Appendix of [26]).

Corollary 2.8. Let $a : A \rightarrow E$ and $l : L \rightarrow E$ be objects of the topos $\mathcal{E}/E$, and $\{e_i : E_i \rightarrow E \mid i \in I\}$ an epimorphic family in $\mathcal{E}$. Then a family of arrows $\{f_i : e_i^*(a) \rightarrow e_i^*(l) \mid i \in I\}$ in the toposes $\mathcal{E}/E_i$ defines a (unique) arrow $f : a \rightarrow l$ in $\mathcal{E}/E$ such that $e_i^*(f) = f_i$ for all $i \in I$ if and only if for every $i, j \in I$, $q_i^*(f_i) = q_j^*(f_j)$, where the arrows q_i and q_j are defined by the following pullback square:

$$\begin{array}{ccc} E_{i,j} & \xrightarrow{q_i} & E_i \\ \downarrow q_j & \searrow e_{i,j} & \downarrow e_i \\ E_j & \xrightarrow{q_j} & E \end{array}$$

□

2.5 $\mathcal{E}$ -final subcategories

The following definition will be important in the sequel. We shall borrow the notation from section 2.4.

Definition 2.9. Let $\underline{\mathcal{A}}_{\mathcal{E}}$ be a $\mathcal{E}$ -indexed category and $i : \underline{\mathcal{B}}_{\mathcal{E}} \rightarrow \underline{\mathcal{A}}_{\mathcal{E}}$ be a $\mathcal{E}$ -indexed functor.

- (a) We say that i is $\mathcal{E}$ -final if for every $E \in \mathcal{E}$ and $x \in \mathcal{A}_E$ there exists a non-empty set $\mathcal{E}_E^x$ of triplets $\{(e_i, b_i, f_i) \mid i \in I\}$ such that the family $\{e_i : E_i \rightarrow E \mid i \in I\}$ is epimorphic, b_i is an object of $\mathcal{B}_{E_i}$ and $f_i : \mathcal{A}_{e_i}(x) \rightarrow i_{E_i}(b_i)$ is an arrow in $\mathcal{A}_{E_i}$ (for each $i \in I$) with the property that for any triplets $\{(e_i, b_i, f_i) \mid i \in I\}$ and $\{(e'_j, c_j, f'_j) \mid j \in J\}$ in $\mathcal{E}_E^x$ there exists an epimorphic family $\{g_k^{i,j} : E_k^{i,j} \rightarrow E_{i,j} \mid k \in K_{i,j}\}$ and for each $k \in K_{i,j}$ an object $d_k^{i,j}$ of $\mathcal{B}_{E_k^{i,j}}$ and arrows $r_k^{i,j} : \mathcal{B}_{q_i \circ g_k^{i,j}}(b_i) \rightarrow d_k^{i,j}$ and $s_k^{i,j} : \mathcal{B}_{q_j \circ g_k^{i,j}}(c_j) \rightarrow d_k^{i,j}$ in $\mathcal{B}_{E_k^{i,j}}$ such that $i_{E_k^{i,j}}(r_k^{i,j}) \circ \mathcal{A}_{q_i}(f_i) = i_{E_k^{i,j}}(s_k^{i,j}) \circ \mathcal{A}_{q_j}(f'_j)$.
- (b) We say that a $\mathcal{E}$ -indexed subcategory $\underline{\mathcal{B}}_{\mathcal{E}}$ of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -final subcategory of $\underline{\mathcal{A}}_{\mathcal{E}}$ if the canonical $\mathcal{E}$ -indexed inclusion functor $i : \underline{\mathcal{B}}_{\mathcal{E}} \hookrightarrow \underline{\mathcal{A}}_{\mathcal{E}}$ is $\mathcal{E}$ -final, in other words if for every $E \in \mathcal{E}$ and $x \in \mathcal{A}_E$ there exists a non-empty set $\mathcal{E}_E^x$ of triplets $\{(e_i, b_i, f_i) \mid i \in I\}$ such that the family $\{e_i : E_i \rightarrow E \mid i \in I\}$ is epimorphic, b_i is an object of $\mathcal{B}_{E_i}$ and $f_i : \mathcal{A}_{e_i}(x) \rightarrow b_i$ is an arrow in $\mathcal{A}_{E_i}$ (for each $i \in I$) with the property that for any triplets $\{(e_i, b_i, f_i) \mid i \in I\}$ and $\{(e'_j, c_j, f'_j) \mid j \in J\}$ in $\mathcal{E}_E^x$ there exists an epimorphic family $\{g_k^{i,j} : E_k^{i,j} \rightarrow E_{i,j} \mid k \in K_{i,j}\}$ and for each $k \in K_{i,j}$ an object $d_k^{i,j}$ of $\mathcal{B}_{E_k^{i,j}}$ and arrows $r_k^{i,j} : \mathcal{B}_{q_i \circ g_k^{i,j}}(b_i) \rightarrow d_k^{i,j}$ and $s_k^{i,j} : \mathcal{B}_{q_j \circ g_k^{i,j}}(c_j) \rightarrow d_k^{i,j}$ in $\mathcal{B}_{E_k^{i,j}}$ such that $r_k^{i,j} \circ \mathcal{A}_{q_i}(f_i) = s_k^{i,j} \circ \mathcal{A}_{q_j}(f'_j)$.

- (c) We say that a $\mathcal{E}$ -indexed subcategory $\underline{\mathcal{B}}_{\mathcal{E}}$ of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ is a *$\mathcal{E}$ -strictly final subcategory* of $\underline{\mathcal{A}}_{\mathcal{E}}$ if for any arrow $f : a \rightarrow b$ in $\mathcal{A}_E$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ such that for any $i \in I$, the arrow $\mathcal{A}_{e_i}(f) : \mathcal{A}_{e_i}(a) \rightarrow \mathcal{A}_{e_i}(b)$ lies in $\mathcal{B}_{E_i}$.
- (d) We say that a $\mathcal{E}$ -indexed subcategory $\underline{\mathcal{B}}_{\mathcal{E}}$ of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ is *$\mathcal{E}$ -full* if for every arrow $f : b \rightarrow b'$ in $\mathcal{A}_E$, where b and b' are objects of $\mathcal{B}_E$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ such that for any $i \in I$ the arrow $\mathcal{A}_{e_i}(f)$ lies in $\mathcal{B}_{E_i}$.

Remarks 2.10. (a) Let $\underline{\mathcal{B}}_{\mathcal{E}}$ be a $\mathcal{E}$ -indexed subcategory of $\underline{\mathcal{A}}_{\mathcal{E}}$ with the property that for every object E of $\mathcal{E}$ and any object $a \in \mathcal{A}_E$ $\alpha : E' \rightarrow E$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ such that for any $i \in I$, the object $\mathcal{A}_{e_i}(a)$ lies in $\mathcal{B}_{E_i}$ and for any arrow $f : a \rightarrow b$ in $\mathcal{A}_E$ where $a, b \in \mathcal{B}_E$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ such that for any $i \in I$, the arrow $\mathcal{A}_{e_i}(f) : \mathcal{A}_{e_i}(a) \rightarrow \mathcal{A}_{e_i}(b)$ lies in $\mathcal{B}_{E_i}$. Then $\underline{\mathcal{B}}_{\mathcal{E}}$ is a $\mathcal{E}$ -strictly final subcategory of $\underline{\mathcal{A}}_{\mathcal{E}}$.

- (b) Every $\mathcal{E}$ -strictly final subcategory is a $\mathcal{E}$ -final subcategory.
- (c) If i is a $\mathcal{E}$ -full embedding of a $\mathcal{E}$ -indexed category $\underline{\mathcal{B}}_{\mathcal{E}}$ into a $\mathcal{E}$ -filtered $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ then i is $\mathcal{E}$ -final if and only if for every $E \in \mathcal{E}$ and $x \in \mathcal{A}_E$ there exists a non-empty epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object b_i of $\mathcal{B}_{E_i}$ and an arrow $f_i : \mathcal{A}_{e_i}(x) \rightarrow b_i$ of $\mathcal{A}_{E_i}$.

Proposition 2.11. Let $\underline{\mathcal{B}}_{\mathcal{E}}$ be a $\mathcal{E}$ -full $\mathcal{E}$ -indexed subcategory of a $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$. Then $\underline{\mathcal{B}}_{\mathcal{E}}$ is $\mathcal{E}$ -filtered if and only if $\underline{\mathcal{A}}_{\mathcal{E}}$ is $\mathcal{E}$ -filtered.

Proof The proof is entirely analogous to the classical one and left to the reader. $\square$

The following result represents a $\mathcal{E}$ -indexed version of the classical theorem formalizing the behaviour of colimits with respect to final subcategories.

Theorem 2.12. Let $i : \underline{\mathcal{B}}_{\mathcal{E}} \hookrightarrow \underline{\mathcal{A}}_{\mathcal{E}}$ be a $\mathcal{E}$ -final functor and $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ a $\mathcal{E}$ -indexed functor. Then D admits a colimit (resp. a $\mathcal{E}$ -indexed colimit) if and only if $D \circ i$ admits a colimit (resp. a $\mathcal{E}$ -indexed colimit), and the two colimits are equal.

Proof First, let us show that any cocone λ over $D \circ i$ with vertex V can be (uniquely) extended to a cocone $\tilde{\lambda}$ over D . For any object a of $\mathcal{A}_E$, we have to define an arrow $\tilde{\lambda}_E(a) : D_E(a) \rightarrow !^*_E(V)$ in $\mathcal{E}/E$. By our hypotheses, there exists an epimorphic family $\mathcal{E} = \{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and a family of arrows $\{f_i : \mathcal{A}_{e_i}(a) \rightarrow i_{E_i}(b_i) \mid i \in I\}$, where b_i lies in $\mathcal{B}_{E_i}$. Consider, for each $i \in I$, the arrow $\lambda_{E_i}(b_i) \circ D_{E_i}(f_i) : D_{E_i}(\mathcal{A}_{e_i}(a)) = e_i^*(D_E(a)) \rightarrow !^*_{E_i}(V) =$

$e_i^*(!_E^*(V))$. By applying the condition in the definition of $\mathcal{E}$ -final functor to the pair $((e_i, f_i, b_i), (e_i, f_i, b_i))$ and exploiting the fact that λ is a cocone over $D \circ i$, we obtain that $q_i^*(\lambda_{E_i}(b_i) \circ D_{E_i}(f_i)) = q_j^*(\lambda_{E_j}(b_j) \circ D_{E_j}(f_j))$ (here we use the notation of Corollary 2.8). By Corollary 2.8, the family of arrows $\lambda_{E_i}(\mathcal{A}_{e_i}(a))$ (for $i \in I$) thus induces a unique arrow $u_{\mathcal{E}} = D_E(a) \rightarrow !_E^*(V)$. To be able to set $\tilde{\lambda}_E(a)$ equal to this arrow, we have to show that such definition does not depend on the choice of the family $\{(e_i, f_i)\}$. But this follows similarly as above, by applying the condition in the definition of $\mathcal{E}$ -final functor and invoking the fact that λ is a cocone over $D \circ i$.

Notice that if $a \in \mathcal{B}_E$ then $\tilde{\lambda}_E(a) = \lambda_E(a)$, and that for any cocone ξ over D , $\xi = \xi \circ i$.

Now that we have showed that for any $a \in \mathcal{A}_E$ the definition of the arrow $\tilde{\lambda}_E(a) : D_E(a) \rightarrow !_E^*(V)$ in the topos $\mathcal{E}/E$ is well-posed, it remains to prove that the assignment $a \rightarrow \tilde{\lambda}_E(a)$ defines a cocone over the diagram D_E with vertex $!_E^*(V)$, i.e. that for any arrow $f : a \rightarrow b$ in $\mathcal{A}_E$, $\tilde{\lambda}_E(b) \circ D_E(f) = \tilde{\lambda}_E(a)$ as arrows in $\mathcal{E}/E$. Further, we have to show that the assignment $E \rightarrow \tilde{\lambda}_E$ defines a $\mathcal{E}$ -indexed cone on the $\mathcal{E}$ -indexed functor D , i.e. that for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$ and any $a \in \mathcal{A}_E$ we have $\alpha^*(\tilde{\lambda}_E(a)) = \tilde{\lambda}_{E'}(\mathcal{A}_\alpha(a))$. This can be easily deduced from the fact that λ is an indexed cocone over $D \circ i$.

The second part of the theorem, for indexed colimits, follows from the first part by noticing that any localization of an indexed $\mathcal{E}$ -final functor with respect to a localization functor $\mathcal{E} \rightarrow \mathcal{E}/E$ is a $\mathcal{E}/E$ -final functor. So it will be sufficient to show the first part.

To complete the proof of the theorem, it remains to show that for any cocone (U, μ) over D , (U, μ) is colimiting over D if and only if $(U, \mu \circ i)$ is colimiting over D .

Suppose that D has a colimiting cocone μ with vertex U . We want to prove that $\mu \circ i$ is a colimiting cocone for the functor $D \circ i$. Let λ be a cocone over the diagram $D \circ i$ with vertex V ; then, as we have just proved, $\tilde{\lambda}$ is a cocone over D with vertex V ; therefore (U, μ) factors through $(V, \tilde{\lambda})$, by an arrow $z : U \rightarrow V$ in $\mathcal{E}$. Clearly, z yields in particular a factorization of $(U, \mu \circ i)$ across (V, λ) . The uniqueness of the factorization of $(U, \mu \circ i)$ across (V, λ) follows from the fact that for any two cocones ξ and χ over $D \circ i$, if ξ factors through χ by an arrow w then $\tilde{\xi}$ factors through $\tilde{\chi}$ by the same arrow. This proves that if (U, μ) is a colimiting cocone over the diagram D then $(U, \mu \circ i)$ is a colimiting cocone over the diagram $D \circ i$.

Conversely, suppose that $(U, \mu \circ i)$ is a colimiting cocone over the diagram $D \circ i$. $(Z, \tilde{\chi})$ is a colimiting cocone over the diagram D . For any cocone (Z, χ) over D , we have that $(Z, \chi \circ i)$ is a cocone over $D \circ i$; therefore $(U, \mu \circ i)$ factors uniquely through $(Z, \chi \circ i)$ or equivalently (by similar arguments as above), $(Z, \tilde{\chi})$ factors uniquely through (U, μ) . Therefore (U, μ) is a colimiting cocone over D , as required. $\square$

Let $\underline{\mathbf{Set}}_{\mathcal{E}}$ be the $\mathcal{E}$ -indexed category given by: $\mathbf{Set}_E = \mathbf{Set}$ for all $E \in \mathcal{E}$ and $\mathbf{Set}_{\alpha} = 1_{\mathbf{Set}}$ (where $1_{\mathbf{Set}}$ is the identical functor on $\mathbf{Set}$) for all arrows α in $\mathcal{E}$. There is a $\mathcal{E}$ -indexed functor

$$\gamma_{\mathcal{E}} : \underline{\mathbf{Set}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$$

defined by: $\gamma_{\mathcal{E}E} = \gamma_{\mathcal{E}/E}^* : \mathbf{Set} \rightarrow \mathcal{E}/E$ (for any object E of $\mathcal{E}$).

We shall consider $\mathcal{E}$ -indexed functors obtained by composing the $\mathcal{E}$ -indexed functor $\gamma_{\mathcal{E}} : \underline{\mathbf{Set}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ with a $\mathcal{E}$ -indexed functor $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathbf{Set}}_{\mathcal{E}}$.

Notice that a $\mathcal{E}$ -indexed functor $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathbf{Set}}_{\mathcal{E}}$ consists of a functor $D_E : \mathcal{A}_E \rightarrow \mathbf{Set}$ for each object E of $\mathcal{E}$ such that for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$, $D_{E'} \circ \mathcal{A}_{\alpha} = D_E$.

Let us give an explicit description of the cocones over the $\mathcal{E}$ -indexed functor $\gamma_{\mathcal{E}} \circ D$. Specializing the general definition, we obtain that a cocone (U, μ) over $\gamma_{\mathcal{E}} \circ D$ consists of an object U of $\mathcal{E}$ and of an arrow $\mu_E(a) : D_E(a) \rightarrow !_E^*(U)$ in $\mathcal{E}/E$ (for each objects E of $\mathcal{E}$ and a of $\mathcal{A}_E$) such that for any arrow $f : a \rightarrow b$ in $\mathcal{A}_E$, $\mu_{E'}(b) \circ \gamma_{\mathcal{E}/E}^*(D_E(f)) = \mu_E(a)$ and for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$, $\alpha^*(\mu_E(a)) = \mu_{E'}(\mathcal{A}_{\alpha}(a))$.

By using the well-known adjunction between $\gamma_{\mathcal{E}}^*$ and the global sections functor $\mathcal{E} \rightarrow \mathbf{Set}$, we can alternatively present the above set of data as follows: a cocone (U, μ) over $\gamma_{\mathcal{E}} \circ D$ consists of an object U of $\mathcal{E}$ and of a function $\mu_E(a) : D_E(a) \rightarrow \text{Hom}_{\mathcal{E}}(E, U)$ (for each objects E of $\mathcal{E}$ and a of $\mathcal{A}_E$) such that for any arrow $f : a \rightarrow b$ in $\mathcal{A}_E$, $\mu_{E'}(b) \circ D_E(f) = \mu_E(a)$ and for any arrow $\alpha : E' \rightarrow E$ in $\mathcal{E}$, $\text{Hom}_{\mathcal{E}}(\alpha, U) \circ \mu_E(a) = \mu_{E'}(\mathcal{A}_{\alpha}(a))$ (notice that the domains of these two arrows are the same since $D_E(a) = D_{E'}(\mathcal{A}_{\alpha}(a))$).

Theorem 2.13. *Let $F : \mathcal{C}^{op} \rightarrow \mathcal{E}$ be a functor. Then the $\mathcal{E}$ -indexed subcategory $\int F_{\mathcal{E}}$ of $\int F_{\mathcal{E}}^f$ is strictly final.*

Proof By definition of $\int F_{\mathcal{E}}^f$, for any object E of $\mathcal{E}$ the category $\int F_E^f$ has as objects the arrows $x : E \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$ and as arrows $x \rightarrow x'$ the arrows $z : E \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$ such that $d_0^F \circ z = x$ and $d_1^F \circ z = x'$. For any $E \in \mathcal{E}$, the category $\int F_E$ thus identifies as a subcategory of the category $\int F_E^f$, through the assignment sending any pair (c, x) , where $x : E \rightarrow F(c)$, to the arrow $\mu_c \circ x : E \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$ and any arrow $(c, x) \rightarrow (c', x')$ to the arrow $J_f \circ x' : E \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$ (here we use the notations of section 2.4). It is clear that these assignments make $\int F_{\mathcal{E}}$ into a $\mathcal{E}$ -indexed subcategory of $\int F_{\mathcal{E}}^f$.

To prove our thesis, we shall apply the criterion of Remark 2.10(a). To show that the $\mathcal{E}$ -indexed subcategory $\int F_{\mathcal{E}}$ satisfies the first condition in the

remark, we observe that for any object $x : E \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$ of the category $\int F^f_E$, if we consider the pullbacks

$$\begin{array}{ccc} E_c & \xrightarrow{x_c} & F(c) \\ \downarrow e_c & & \downarrow \mu_c \\ E & \xrightarrow{x} & \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \end{array}$$

of x along the coproduct arrows $\mu_c : F(c) \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$, we obtain an epimorphic family $\{e_c : E_c \rightarrow E \mid c \in \text{Ob}(\mathcal{C})\}$ such that $\int F^f_{e_c}(x)$ lies in the subcategory $\int F_{E_c}$.

It remains to prove that the second condition of Remark 2.10(a) is satisfied. Let us suppose that (c, x) and (c', x') are objects of the subcategory $\int F_E$, and suppose that $\alpha : E \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$ is an arrow such that $d_0^F \circ \alpha = \mu_c \circ x$ and $d_1^F \circ \alpha = \mu_{c'} \circ x'$:

$$\begin{array}{ccc} E & \xrightarrow{\alpha} & \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \\ \downarrow x & & \downarrow d_0^F \\ F(c) & \xrightarrow{\mu_c} & \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \end{array} \quad \begin{array}{ccc} E & \xrightarrow{\alpha} & \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \\ \downarrow x' & & \downarrow d_1^F \\ F(c') & \xrightarrow{\mu_{c'}} & \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \end{array} .$$

Consider, for each $f \in \text{Arr}(\mathcal{C})$, the following pullback square:

$$\begin{array}{ccc} E_f & \xrightarrow{\alpha_f} & F(\text{cod}(f)) \\ \downarrow e_f & & \downarrow J_f \\ E & \xrightarrow{\alpha} & \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \end{array} .$$

The commutativity of the three diagrams above, combined with the fact that distinct coproduct arrows are disjoint, implies that for any arrow f of $\mathcal{C}$ such that $\text{dom}(f) \neq c$ or $\text{cod}(f) \neq c'$ we have $E_f \cong 0_{\mathcal{E}}$. We can therefore restrict our attention to the arrows f such that $\text{dom}(f) = c$ and $\text{cod}(f) = c'$. For any such arrow f , we have $\alpha_f = x' \circ e_f$ (equivalently, since $\mu_{c'} = \mu_{\text{cod}(f)}$ is monic, $\mu_{c'} \circ \alpha_f = \mu_{c'} \circ x' \circ e_f$). Indeed, $\mu_{c'} \circ \alpha_f = d_1^F \circ J_f \circ \alpha_f = d_1^F \circ \alpha \circ e_f = \mu_{c'} \circ x' \circ e_f$. Therefore $\alpha \circ e_f = J_f \circ x' \circ e_f$; indeed, $\alpha \circ e_f = J_f \circ \alpha_f = J_f \circ x' \circ e_f$. Lastly, we observe that the arrow f defines an arrow $(c, x \circ e_f) \rightarrow (c', x' \circ e_f)$ in the subcategory $(\int F)_{E_f}$, i.e.

$F(f) \circ x' \circ e_f = x \circ e_f$. Indeed, the arrow $\mu_{\text{dom}(f)} = \mu_{c'}$ is monic and we have $\mu_{\text{dom}(f)} \circ F(f) \circ x' \circ e_f = d_0^F \circ J_f \circ x' \circ e_f = J_f \circ \alpha \circ e_f = \mu_c \circ x \circ e_f$.

The epimorphic family $\{e_f : E_f \rightarrow E \mid \text{dom}(f) = c \text{ and } \text{cod}(f) = c'\}$ thus satisfies the property that the arrow $(\int F_{\mathcal{E}}^f)_{e_f}(\alpha)$ lies in the subcategory $(\int F)_{E_f}$. Our proof is therefore complete. $\square$

Theorem 2.14. *Let $\mathcal{C}$ be a small category, $\mathcal{E}$ a Grothendieck topos, $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ and $P : \mathcal{C} \rightarrow \mathbf{Set}$ functors and $P_{\mathcal{E}}$ the internal diagram in $[\mathbb{C}, \mathcal{E}]$ given by $\overline{\gamma_{\mathcal{E}}^* \circ P}$. Then the restriction of the functor $(P_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f) \cong \overline{P_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f}$ to the $\mathcal{E}$ -indexed subcategory $\int \overline{F}_{\mathcal{E}}$ of $\int F_{\mathcal{E}}^f$ is naturally isomorphic to $\gamma_{\mathcal{E}} \circ z_{\mathcal{E}}$, where $z_{\mathcal{E}} : \int \overline{F}_{\mathcal{E}} \rightarrow \mathbf{Set}_{\mathcal{E}}$ is the $\mathcal{E}$ -indexed functor defined by: $z_{\mathcal{E}E}((c, x)) = P(c)$ and $z_{\mathcal{E}E}(f) = P(f)$ (for any $E \in \mathcal{E}$, object (c, x) and arrow f in the category $\int F_E$).*

Proof We shall exhibit an isomorphism

$$\overline{P_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f}((c, x)) \cong (\gamma_{\mathcal{E}} \circ z_{\mathcal{E}})_E((c, x)),$$

natural in $E \in \mathcal{E}$ and $(c, x) \in \int F_E$.

Consider the following pullback diagram in $\mathbf{cat}(\mathcal{E})$:

$$\begin{array}{ccc} \overline{F} \otimes_{\mathbb{C}} P & \xrightarrow{t_P} & \int^{opf} P_{\mathcal{E}} \\ \downarrow t_{\overline{F}} & & \downarrow \pi_{P_{\mathcal{E}}}^{opf} \\ \int^f \overline{F} & \xrightarrow{\pi_{\overline{F}}} & \mathbb{C}. \end{array}$$

Since coproducts are stable under pullback in a topos, we have that:

$$(\overline{F} \otimes_{\mathbb{C}} P)_0 = \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \times \gamma_{\mathcal{E}}^*(P(c)),$$

$$(\overline{F} \otimes_{\mathbb{C}} P)_1 = \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))),$$

$$t_{F0} : (F \otimes_{\mathbb{C}} P)_0 = \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \times \gamma_{\mathcal{E}}^*(P(c)) \rightarrow (\int^f F)_0 = \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$$

is equal to the arrow $\coprod_{c \in \text{Ob}(\mathcal{C})} \pi_{F(c)}$, where $\pi_{F(c)}$ is the canonical projection $F(c) \times \gamma_{\mathcal{E}}^*(P(c)) \rightarrow F(c)$,

$$t_{\overline{F}1} : (\overline{F} \otimes_{\mathbb{C}} P)_1 = \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))) \rightarrow (\int^f F)_1 = \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f))$$

is equal to the arrow $\coprod_{f \in \text{Arr}(\mathcal{C})} \pi'_{F(\text{cod}(f))}$, where $\pi'_{F(\text{cod}(f))}$ is the canonical projection $F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))) \rightarrow F(\text{cod}(f))$ and the domain and codomain arrows

$$d_0^{\overline{F} \otimes_{\mathbb{C}} P}, d_1^{\overline{F} \otimes_{\mathbb{C}} P} : (\overline{F} \otimes_{\mathbb{C}} P)_1 \rightarrow (\overline{F} \otimes_{\mathbb{C}} P)_0$$

are defined by the following conditions:

$$d_0^{\overline{F} \otimes_{\mathbb{C}} P} \circ W_f = Z_{\text{cod}(f)} \circ \langle \pi'_{F(\text{cod}(f))}, \gamma_{\mathcal{E}}^*(P(f)) \circ \mu_{\gamma_{\mathcal{E}}^*(P(\text{dom}(f)))} \rangle$$

and

$$d_1^{\overline{F} \otimes_{\mathbb{C}} P} \circ W_f = Z_{\text{dom}(f)} \circ \langle F(f) \circ \pi'_{F(\text{cod}(f))}, \mu_{\gamma_{\mathcal{E}}^*(P(\text{dom}(f)))} \rangle,$$

where $\mu_{\gamma_{\mathcal{E}}^*(P(\text{dom}(f)))}$ is the canonical projection

$$F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))) \rightarrow \gamma_{\mathcal{E}}^*(P(\text{dom}(f)))$$

and

$$W_f : F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))) \rightarrow \coprod_{f \in \text{Arr}(\mathcal{C})} F(\text{cod}(f)) \times \gamma_{\mathcal{E}}^*(P(\text{dom}(f))),$$

$$Z_c : F(c) \times \gamma_{\mathcal{E}}^*(P(c)) \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \times \gamma_{\mathcal{E}}^*(P(c))$$

are the canonical coproduct arrows (respectively for $f \in \text{Arr}(\mathcal{C})$ and $c \in \text{Ob}(\mathcal{C})$).

Now, the internal diagram $P_{\mathcal{E}} \circ \pi_{\overline{F}}^f$ corresponds precisely to the discrete opfibration $t_{\overline{F}}$ and hence the $\mathcal{E}$ -indexed functor $P_{\mathcal{E}} \circ \pi_{\overline{F}}^f$ sends any generalized element $x : E \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c)$ to the object of $\mathcal{E}/E$ given by the pullback of $t_{\overline{F}_0}$ along it. In particular, since in a topos any diagram of the form

$$\begin{array}{ccc} A_i & \longrightarrow & \coprod_{i \in I} A_i \\ \downarrow f_i & & \downarrow \coprod_{i \in I} f_i \\ B_i & \longrightarrow & \coprod_{i \in I} B_i, \end{array}$$

where the horizontal arrows are the canonical coproduct arrows, is a pull-back and products commute with pullbacks, the functor $(P_{\mathcal{E}} \circ \pi_{\overline{F}}^f)_E$ sends any object (c, x) of the category $\int F_E$ to the object given by the canonical projection $E \times \gamma_{\mathcal{E}}^*(P(c)) \rightarrow E$. But this object is canonically isomorphic to $\gamma_{\mathcal{E}/E}^*(P(c))$ in $\mathcal{E}/E$, which is the value at (c, x) of the functor $(\gamma_{\mathcal{E}} \circ z_{\mathcal{E}})_E$. Our proof is therefore complete. $\square$

Let us now explicitly describe the $\mathcal{E}$ -indexed colimiting cocone on the restriction to the $\mathcal{E}$ -indexed subcategory $\int \overline{F}_{\mathcal{E}}$ of the $\mathcal{E}$ -indexed diagram $\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f$ (cf. section 2.1 for the background). The vertex of this colimiting cocone is the codomain of the coequalizer $w : (\overline{F} \otimes_{\mathbb{C}} P)_0 \rightarrow \underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f)$ of the pair of arrows $d_0^{\overline{F} \otimes_{\mathbb{C}} P}, d_1^{\overline{F} \otimes_{\mathbb{C}} P} : (\overline{F} \otimes_{\mathbb{C}} P)_1 \rightarrow (\overline{F} \otimes_{\mathbb{C}} P)_0$. For any object (c, x) of the category $\int F_E$, the colimit arrow $(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f)_E((c, x)) \cong \gamma_{\mathcal{E}/E}^*(P(c)) \rightarrow \underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f)$ is given by the composite $!_E^*(w) \circ h_{(c,x)}$, where $h_{(c,x)}$ is the arrow $!_E^*(Z_c) \circ \langle x \times 1_{\gamma_{\mathcal{E}}^*(P(c))}, \pi_E \rangle : \gamma_{\mathcal{E}/E}^*(P(c)) \cong E \times \gamma_{\mathcal{E}}^*(P(c)) \rightarrow \coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \times \gamma_{\mathcal{E}}^*(P(c)) \times E \cong !_E^*(\coprod_{c \in \text{Ob}(\mathcal{C})} F(c) \times \gamma_{\mathcal{E}}^*(P(c)))$; in other words, the function $\xi_{(c,x)} : P(c) \rightarrow \text{Hom}_{\mathcal{E}}(E, \underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f))$ corresponding to it assigns to every element $a \in P(c)$ the arrow $w \circ Z_c \circ \langle x, r_a \circ !_E \rangle$, where $r_a : 1_{\mathcal{E}} \rightarrow \gamma_{\mathcal{E}}^*(P(c)) \cong \coprod_{a \in P(c)} 1_{\mathcal{E}}$ is the coproduct arrow corresponding to the element a and $!_E$ is the unique arrow $E \rightarrow 1_{\mathcal{E}}$ in $\mathcal{E}$.

It is immediate to verify that, under the isomorphism

$$\underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f) \cong \text{colim}(F \circ \pi_P^f)$$

of Theorem 2.5, the functions $\xi_{(c,x)} : P(c) \rightarrow \underline{\text{colim}}_{\mathcal{E}}(\underline{P_{\mathcal{E}}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f) \cong \text{colim}(F \circ \pi_P^f)$ admit the following description in terms of the colimiting arrows $\chi_{(c,a)} : F(c) \rightarrow \text{colim}(F \circ \pi_P^f)$: for any $a \in P(c)$, $\xi_{(c,x)}(a) = \chi_{(c,a)} \circ x$.

2.6 A characterization of $\mathcal{E}$ -indexed colimits

In this section we establish necessary and sufficient conditions for a $\mathcal{E}$ -indexed cocone on a small diagram in $\mathcal{E}$ to be a ($\mathcal{E}$ -indexed) colimit cone.

Before stating and proving the main theorem of this section, we need to introduce some relevant definitions and a few technical lemmas.

Given a $\mathcal{E}$ -indexed functor $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ and an object R of the topos $\mathcal{E}$, we denote by $\mathcal{I}_R^D$ the set of pairs of the form (x, y) , where x is an object of the category $\mathcal{A}_R$ and y is an arrow $1_{\mathcal{E}/R} \rightarrow D_R(x)$ in the topos $\mathcal{E}/R$. On such a set we consider the equivalence relation $\mathcal{R}_R^D$ generated by the pairs of the form $((x, y), (x', y'))$, where there exists an arrow $f : x \rightarrow x'$ in the category $\mathcal{A}_R$ such that $y' = D_R(f) \circ y$.

The following remarks are useful in connection with the application of the localization technique.

Remarks 2.15. (a) For any $E \in \mathcal{E}$, we have a natural bijection $\mathcal{I}_E^D \cong \mathcal{I}_{1_{\mathcal{E}/E}}^{D/E}$, and the relation $\mathcal{R}_E^D$ corresponds to the relation $\mathcal{R}_{1_{\mathcal{E}/E}}^{D/E}$ under this bijection.

- (b) For any arrow $f : F \rightarrow E$ and any object $x \in \mathcal{A}_E$, $D_F(\mathcal{A}_f(x)) = f^*(D_E(x))$ (this follows from the fact that D is a $\mathcal{E}$ -indexed functor). We shall denote the canonical arrow $\text{dom}(D_F(\mathcal{A}_f(x))) \rightarrow \text{dom}(D_E(x))$ by the symbol r_f .
- (c) For any object x of $\mathcal{A}_E$ and any arrow $y : (f : F \rightarrow E) \rightarrow D_E(x)$ in the topos $\mathcal{E}/E$, there exists a unique arrow $y_f : (1_F : F \rightarrow E) \rightarrow D_F(\mathcal{A}_f(x))$ in the topos $\mathcal{E}/F$ such that $r_f \circ y_f = y$ (where r_f is considered here as an arrow in $\mathcal{E}/E$ in the obvious way);
- (d) For any $\mathcal{E}$ -indexed cocone μ over D with vertex U , any object x of $\mathcal{A}_E$ and any arrow $y : (f : F \rightarrow E) \rightarrow D_E(x)$ in the topos $\mathcal{E}/E$, we have $\mu_E(x) \circ y = (1_U \times f) \circ \mu_F(\mathcal{A}_f(x)) \circ y_f$. Moreover, we have pullback squares

$$\begin{array}{ccc} D_F(\mathcal{A}_f(x)) & \xrightarrow{r_f} & D_E(x) \\ \downarrow \mu_F(\mathcal{A}_f(x)) & & \downarrow \mu_E(x) \\ U \times F & \xrightarrow{1_U \times f} & U \times E \\ \downarrow \pi_F & & \downarrow \pi_E \\ F & \xrightarrow{f} & E \end{array}$$

in $\mathcal{E}$, where π_E and π_F are the obvious canonical projections. In particular, for any other pair (x', y') , $\mu_E(x) \circ y = \mu_E(x') \circ y'$ if and only if $\mu_F(\mathcal{A}_f(x)) \circ y_f = \mu_F(\mathcal{A}_f(x')) \circ y'_f$;

- (e) If $\underline{\mathcal{A}}_{\mathcal{E}}$ is the $\mathcal{E}$ -externalization of an internal category $\mathbb{C}$ in $\mathcal{E}$, x is a generalized element $E \rightarrow \mathbb{C}_0$ and y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$, there exists a unique arrow $y_f : (1_F : F \rightarrow F) \rightarrow D_F(x \circ f)$ in $\mathcal{E}/F$ such that $z_{x \circ f} \circ y_f = z_x \circ y$ (this arrow is provided by the universal property of the pullback square defining $D_F(x \circ f)$), which coincides with the arrow y_f defined in Remark 2.15(c);
- (f) If a pair $((x, y), (x', y'))$ belongs to $\mathcal{R}_R^D$ then for any indexed cocone λ over D with vertex V , $\lambda_R(x) \circ y = \lambda_R(x') \circ y'$. Indeed, for any pair $((x, y), (x', y'))$ with the property that there exists an arrow $f : x \rightarrow x'$ in the category $\mathcal{A}_R$ such that $y' = D_E(f) \circ y$, we have that $\lambda_R(x') \circ y' = \lambda_R(x') \circ D_E(f) \circ y = \lambda_R(x) \circ y$ (since λ_R is a cocone).

Lemma 2.16. *Let $D \in [\mathbb{C}, \mathcal{E}]$ be an internal diagram, $x : 1 \rightarrow \mathbb{C}_0$ and $f : 1 \rightarrow \mathbb{C}_1$ generalized elements such that $d_0^{\mathbb{C}} \circ f = x$ and $m : F \rightarrow D_1(x)$ an arrow in $\mathcal{E}$. Then the unique arrow $\chi_{m,f} : F \rightarrow (f^{opf} D)_1$ in $\mathcal{E}$ such that $d_0^{f^{opf} D} \circ \chi_{m,f} = z_x \circ m$ and $(\pi_D^{opf})_1 \circ \chi_{m,f} = f \circ !_F$ (provided by the universal property of the pullback square corresponding to the domains of the discrete opfibration associated to D) satisfies the property that $z_x \circ m = d_0^{f^{opf} D} \circ \chi_{m,f}$*

(by definition) and $z_{x'} \circ D_1(f) \circ m = d_1^{\int^{opf} D} \circ \chi_{m,f}$, where x' is equal to $d_1^{\mathbb{C}} \circ f$ and $d_0^{\int^{opf} D}, d_1^{\int^{opf} D}$ are respectively the domain and codomain arrows $(\int^{opf} D)_1 \rightarrow (\int^{opf} D)_0$.

Proof By definition of $D_1(f)$, we have that $z_{x'} \circ D_1(f) = d_1^{\int^{opf} D} \circ r \circ \xi$, where r is defined by the pullback square

$$\begin{array}{ccc} H & \xrightarrow{r} & (\int^{opf} D)_1 \\ \downarrow !_H & & \downarrow \pi_D^{opf}{}_1 \\ 1_{\mathcal{E}} & \xrightarrow{f} & \mathbb{C}_1 \end{array}$$

and ξ is the unique arrow $D_1(x) \rightarrow H$ such that $z_x = d_0^{\int^{opf} D} \circ r \circ \xi$, provided by the universal property of the pullback square

$$\begin{array}{ccc} H & \xrightarrow{d_0^{\int^{opf} D}} & (\int^{opf} D)_0 \\ \downarrow s & & \downarrow (\pi_D^{opf})_0 \\ 1_{\mathcal{E}} & \xrightarrow{d_0^{\mathbb{C}} \circ f} & \mathbb{C}_0. \end{array}$$

It thus remains to prove that $\chi_{m,f} = r \circ \xi \circ m$. But this immediately follows from the universal property of the pullback

$$\begin{array}{ccc} \int^{opf} D_1 & \xrightarrow{d_0^{\int^{opf} D}} & (\int^{opf} D)_0 \\ \downarrow \pi_D^{opf}{}_1 & & \downarrow \pi_D^{opf}{}_0 \\ \mathbb{C}_1 & \xrightarrow{d_0^{\mathbb{C}}} & \mathbb{C}_0, \end{array}$$

since $d_0^{\int^{opf} D} \circ \chi_{m,f} = d_0^{\int^{opf} D} \circ (r \circ \xi \circ m)$ and $\pi_D^{opf}{}_1 \circ \chi_{m,f} = \pi_D^{opf}{}_1 \circ (r \circ \xi \circ m)$. $\square$

Lemma 2.17. *Let D be an internal diagram in $[\mathbb{C}, \mathcal{E}]$ and χ be an arrow $1_{\mathcal{E}} \rightarrow (\int^{opf} D)_1$. Let $x = (\pi_D^{opf})_0 \circ d_0^{\int^{opf} D} \circ \chi$, $x' = (\pi_D^{opf})_0 \circ d_1^{\int^{opf} D} \circ \chi$ and $f = (\pi_D^{opf})_1 \circ \chi$. Then $d_0^{\mathbb{C}} \circ f = x$, $d_1^{\mathbb{C}} \circ f = x'$ and $D_1(f) \circ \chi_0 = \chi_1$, where χ_0 is the unique arrow $1_{\mathcal{E}} \rightarrow D_1(x)$ such that $z_x \circ \chi_0 = d_0^{\int^{opf} D} \circ \chi$ (which exists by the universal property of the pullback square defining $D_1(x)$) and χ_1 is the unique arrow $1_{\mathcal{E}} \rightarrow D_1(x')$ such that $z_{x'} \circ \chi_1 = d_1^{\int^{opf} D} \circ \chi$ (which exists by the universal property of the pullback square defining $D_1(x')$).*

Proof The first two identities follow straightforwardly from the definition. It remains to prove that $D_1(f) \circ \chi_0 = \chi_1$. Let us refer to the proof of Lemma 2.16 for notation. By the universal property of the pullback square defining $D_1(x')$, it is equivalent to verify that $z_{x'} \circ D_1(f) \circ \chi_0 = z_{x'} \circ \chi_1$. But $z_{x'} \circ D_1(f) \circ \chi_0 = d_1^{\int^{opf} D} \circ r \circ \xi \circ \chi_0$, while $z_{x'} \circ \chi_1 = d_1^{\int^{opf} D} \circ \chi$; so to prove the desired equality it suffices to show that $r \circ \xi \circ \chi_0 = \chi$. This follows from the fact that $d_0^{\int^{opf} D} \circ r \circ \xi \circ \chi_0 = z_x \circ \chi_0 = d_0^{\int^{opf} D} \circ \chi$ by virtue of the universal property of the pullback square given by the domain of the discrete opfibration associated to D . $\square$

Proposition 2.18. *Let $D \in [\mathbb{C}, \mathcal{E}]$ be an internal diagram, $x, x' : 1 \rightarrow \mathbb{C}_0$ generalized elements and $y : 1_{\mathcal{E}} \rightarrow D_1(x)$, $y' : 1_{\mathcal{E}} \rightarrow D_1(x')$ arrows in $\mathcal{E}$. Then $(z_x \circ y, z_{x'} \circ y')$ belongs to the equivalence relation on $\text{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, (\int^{opf} D)_0)$ generated by the pairs of the form $(d_0^{\int^{opf} D} \circ a, d_1^{\int^{opf} D} \circ a)$ for some arrow $a : 1_{\mathcal{E}} \rightarrow (\int^{opf} D)_1$ if and only if $((x, y), (x', y'))$ belongs to $\mathcal{R}_{1_{\mathcal{E}}}^D$.*

Proof Let us first prove the ‘if’ direction. We have that $((x, y), (x', y'))$ belongs to $\mathcal{R}_{1_{\mathcal{E}}}^D$ if and only if there exists a finite sequence

$$(x_0, y_0) = (x, y), \dots, (x_n, y_n) = (x', y')$$

such that for any $i \in \{0, \dots, n-1\}$

- (1) either there exists $f_i : 1 \rightarrow \mathbb{C}_1$ such that $d_0^{\mathbb{C}} \circ f_i = x_i$, $d_1^{\mathbb{C}} \circ f_i = x_{i+1}$ and $D_1(f_i) \circ y_i = y_{i+1}$ or
- (2) there exists $g_i : 1 \rightarrow \mathbb{C}_1$ such that $d_0^{\mathbb{C}} \circ g_i = x_{i+1}$, $d_1^{\mathbb{C}} \circ g_i = x_i$ and $D_1(g_i) \circ y_{i+1} = y_i$.

In case (1), Lemma 2.16 implies (by taking m to be y_i and f to be f_i) that $z_{x_i} \circ y_i = d_0^{\int^{opf} D} \circ \chi_{y_i, f_i}$ and $z_{x_{i+1}} \circ y_{i+1} = d_1^{\int^{opf} D} \circ \chi_{y_i, f_i}$ (here the notation is that of the lemma). In case (2), Lemma 2.16 implies (by taking m to be y_{i+1} and f to be g_i) that $z_{x_{i+1}} \circ y_{i+1} = d_0^{\int^{opf} D} \circ \chi_{y_{i+1}, g_i}$ and $z_{x_i} \circ y_i = d_1^{\int^{opf} D} \circ \chi_{y_{i+1}, g_i}$.

From this it clearly follows that $(z_x \circ y, z_{x'} \circ y')$ belongs to the equivalence relation T on $\text{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, (\int^{opf} D)_0)$ generated by the pairs of the form $(d_0^{\int^{opf} D} \circ a, d_1^{\int^{opf} D} \circ a)$ for some arrow $a : 1_{\mathcal{E}} \rightarrow (\int^{opf} D)_1$, as required.

Let us now prove the ‘only if’ direction. If $(z_x \circ y, z_{x'} \circ y')$ belongs to the equivalence relation T defined above, there exists a finite sequence $\chi_0, \dots, \chi_n$ of arrows $1_{\mathcal{E}} \rightarrow (\int^{opf} D)_1$ and of arrows e_i (which are either $d_0^{\int^{opf} D}$ or $d_1^{\int^{opf} D}$ - we denote by e_i^{op} the arrow $d_1^{\int^{opf} D}$ if e_i is $d_0^{\int^{opf} D}$ and the arrow $d_0^{\int^{opf} D}$

if e_i is $d_1^{\int^{opf} D}$ such that $e_0 \circ \chi_0 = z_x \circ y$, $e_{i+1} \circ \chi_{i+1} = e_i^{\text{op}} \circ \chi_i$ for all $i \in \{0, \dots, n-1\}$ and $e_n \circ \chi_n = z_{x'} \circ y'$.

To deduce our thesis, it suffices to apply Lemma 2.17 noticing that if $e_i \circ \chi = z_{x''} \circ y''$ then $\chi_0 = y''$ if e_i is $d_0^{\int^{opf} D}$ and $\chi_1 = y''$ if e_i is $d_1^{\int^{opf} D}$ (where χ_0 and χ_1 are the arrows defined in the statement of Lemma 2.17) since $z_{x''} \circ \chi_0 = d_0^{\int^{opf} D} \circ \chi = z_{x''} \circ y''$ (if e_i is $d_0^{\int^{opf} D}$) and $z_{x''} \circ \chi_1 = d_1^{\int^{opf} D} \circ \chi = z_{x''} \circ y''$ (if e_i is $d_1^{\int^{opf} D}$). $\square$

Theorem 2.19. *Let $D : \mathcal{A}_{\mathcal{E}} \rightarrow \mathcal{E}_{\mathcal{E}}$ be a $\mathcal{E}$ -indexed functor, where $\mathcal{A}_{\mathcal{E}}$ is equivalent to the $\mathcal{E}$ -externalization of an internal category in $\mathcal{E}$. Then a cocone μ over D with vertex U is an indexed colimiting cocone for D if and only if the following conditions are satisfied:*

- (i) *For any object F of $\mathcal{E}$ and arrow $h : F \rightarrow U$ in the topos $\mathcal{E}$, there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object $x_i \in \mathcal{A}_{F_i}$ and an arrow $\alpha_i : 1_{\mathcal{E}/F_i} \rightarrow D_{F_i}(x_i)$ in the topos $\mathcal{E}/F_i$ such that $\langle h \circ f_i, 1_{F_i} \rangle = \mu_{F_i}(x_i) \circ \alpha_i$ as arrows $1_{\mathcal{E}/F_i} \rightarrow F_i^*(U)$ in $\mathcal{E}/F_i$;*
- (ii) *For any pairs (x, y) and (x', y') , where x and x' are objects of $\mathcal{A}_{\mathcal{E}}$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$ and y' is an arrow $(f : F \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, we have $\mu_E(x) \circ y = \mu_E(x') \circ y'$ if and only if there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ such that the pair $((\mathcal{A}_{f \circ f_i}(x), f_i^*(y)), (\mathcal{A}_{f \circ f_i}(x'), f_i^*(y'))$ belongs to the relation $\mathcal{R}_{F_i}^D$.*

Proof First, let us prove that, under the assumption that $\mathcal{A}_{\mathcal{E}}$ is the $\mathcal{E}$ -externalization of an internal category $\mathbb{C}$ in $\mathcal{E}$, the colimit cone for D satisfies the two conditions of the theorem.

Under this assumption, condition (i), applied to the colimiting cocone μ for D , rewrites as follows: for any object F of $\mathcal{E}$ and arrow $h : F \rightarrow \text{colim}(D)$ in $\mathcal{E}$, there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a generalized element $x_i : F_i \rightarrow \mathbb{C}_0$ and an arrow $\alpha_i : 1_{\mathcal{E}/F_i} \rightarrow D_{F_i}(x_i)$ in the topos $\mathcal{E}/F_i$ such that $\langle h \circ f_i, 1_{F_i} \rangle = c \circ z_{x_i} \circ \alpha_i$ as arrows $(f_i : 1_{\mathcal{E}/F_i} \rightarrow F_i^*(\text{colim}(D)))$ in $\mathcal{E}/F_i$ (where the notation is that of section 2.1).

Consider the following pullback square:

$$\begin{array}{ccc} F' & \xrightarrow{f'} & F \\ \downarrow h' & & \downarrow h \\ (\int^{opf} D)_0 & \xrightarrow{c} & \text{colim}(D) . \end{array}$$

Since the arrow c is an epimorphism, the arrow $f' : F' \rightarrow F$ is an epimorphism. This arrow will form, by itself, the single element of an epimorphic

family satisfying condition (i). We set x' equal to the composite $(\pi_D^{opf})_0 \circ h'$ (recall that $(\pi_D^{opf})_0 : (\int^{opf} D)_0 \rightarrow \mathbb{C}_0$ is the object component of the discrete opfibration associated to D). Consider the following pullback square defining $D_{F'}(x')$:

$$\begin{array}{ccc} D_{F'}(x') & \xrightarrow{u} & (\int^{opf} D)_0 \\ \downarrow v & & \downarrow (\pi_D^{opf})_0 \\ F' & \xrightarrow{x'} & \mathbb{C}_0. \end{array}$$

By the universal property of this pullback square, there exists a unique arrow $\alpha_{x'} : F' \rightarrow D_{F'}(x')$ in $\mathcal{E}$ such that $u \circ \alpha_{x'} = h'$ and $v \circ \alpha_{x'} = 1_{F'}$. So $\alpha_{x'}$ is an arrow $1_{\mathcal{E}/F'} \rightarrow D_F(x')$ in the topos $\mathcal{E}/F'$, and $\langle h \circ f', 1_{F'} \rangle = \langle c \circ u \circ \alpha_{x'}, v \circ \alpha_{x'} \rangle$, as required.

Under the assumption that $\underline{\mathcal{A}}_{\mathcal{E}}$ is the $\mathcal{E}$ -externalization of an internal category $\mathbb{C}$ in $\mathcal{E}$, condition (ii), applied to the colimiting cocone μ for D , rewrites as follows: for any pairs (x, y) and (x', y') , where x and x' is a generalized elements $E \rightarrow \mathbb{C}_0$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$ and y' is an arrow $(f : F \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, we have $c \circ z_x \circ y = c \circ z_{x'} \circ y'$ if and only if there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ such that the pair $((x \circ f \circ f_i, y \circ f_i), (x' \circ f \circ f_i, y' \circ f_i))$ belongs to the relation $\mathcal{R}_{F_i}^D$.

Thanks to the localization technique, we can suppose $E = 1_{\mathcal{E}}$ without loss of generality. Indeed, condition (ii) for the diagram D , the cocone μ and the object E is equivalent to condition (ii) for the diagram D/E , the cocone μ/E and the object $1_{\mathcal{E}/E}$, and μ is, by our assumption, an indexed colimiting cocone and hence stable under localization.

By Lemma 2.29, we have that $c \circ z_x \circ y = c \circ z_{x'} \circ y'$ if and only if there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ such that for each $i \in I$ $(z_x \circ y \circ f_i, z_{x'} \circ y' \circ f_i)$ belongs to the equivalence relation on the set $Hom_{\mathcal{E}}(F_i, (\int^{opf} D)_0)$ generated by the set of pairs of the form $(d_0^{\int^{opf} D} \circ a, d_1^{\int^{opf} D} \circ a)$, for a generalized element $a : F_i \rightarrow (\int^{opf} D)_0$ in $\mathcal{E}$.

By Remark 2.15(e), we have that $(z_x \circ y \circ f_i, z_{x'} \circ y' \circ f_i) = (z_{x \circ f \circ f_i} \circ (y \circ f_i), z_{x' \circ f \circ f_i} \circ (y' \circ f_i))$. Our thesis then follows from Proposition 2.18, applied to the toposes $\mathcal{E}/F_i$ and the pairs $((x \circ f \circ f_i, y \circ f_i), (x' \circ f \circ f_i, y' \circ f_i))$.

Let us now prove that the conditions of the theorem are sufficient for μ to be a colimiting cocone. For this part of the theorem, we shall not need to assume the $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$ to be small.

Since conditions (i) and (ii) are both stable under localization, it suffices to prove that μ is a universal colimiting cocone over the diagram D . Suppose that λ is an indexed cocone over D with vertex V . We have to prove that there exists a unique arrow $l : U \rightarrow V$ in $\mathcal{E}$ such that for any $E \in \mathcal{E}$ and object x of $\mathcal{A}_E$, $(l \times 1_E) \circ \mu_E(x) = \lambda_E(x)$. To define the arrow l we shall

define a function $L : \text{Hom}_{\mathcal{E}}(E, U) \rightarrow \text{Hom}_{\mathcal{E}}(E, V)$ natural in $E \in \mathcal{E}$. Given $h \in \text{Hom}_{\mathcal{E}}(E, U)$, by condition (i) in the statement of the theorem, there exists an epimorphic family $\mathcal{E} = \{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object x_i of $\mathcal{A}_{E_i}$ and an arrow $\alpha_i : 1_{\mathcal{E}/E_i} \rightarrow D_{E_i}(x_i)$ in $\mathcal{E}/E_i$ such that $\langle h \circ e_i, 1_{E_i} \rangle = \mu_{E_i}(x_i) \circ \alpha_i$. The family of arrows $\{\lambda_{E_i}(x_i) \circ \alpha_i \mid i \in I\}$ satisfies the hypotheses of Corollary 2.8. Indeed, using the notation of the corollary, we have that $q_i^*(\lambda_{E_i}(x_i) \circ \alpha_i) = q_j^*(\lambda_{E_j}(x_j) \circ \alpha_j)$ for all $i, j \in I$; this can be easily proved by using condition (ii) of the theorem, the fact that the same identity holds for the cocone μ (in place of λ) and Remark 2.15(d). Therefore there exists a unique arrow $L_h^{\mathcal{E}} : E \rightarrow V$ such that $\langle L_h^{\mathcal{E}} \circ e_i, 1_{E_i} \rangle = \lambda_{E_i}(x_i) \circ \alpha_i$ for all $i \in I$. In order to be able to define $L(h)$ as equal to $L_h^{\mathcal{E}}$ we need to check that for any other epimorphic family $\mathcal{E}' = \{e'_j : E'_j \rightarrow E \mid j \in J\}$ in $\mathcal{E}$, we have $L_h^{\mathcal{E}} = L_h^{\mathcal{E}'}$. To this end, consider the fibered product of the epimorphic families $\mathcal{E}$ and $\mathcal{E}'$, that is the family of arrows $p_{i,j} = e_i \circ f_i = e'_j \circ f'_j : F_{i,j} \rightarrow E$ (for $i, j \in I \times J$), where the arrows f_i and f'_j are defined by the following pullback square:

$$\begin{array}{ccc} F_{i,j} & \xrightarrow{f_i} & E_i \\ \downarrow f'_j & & \downarrow e_i \\ E_j & \xrightarrow{e'_j} & E \end{array}$$

It clearly suffices to verify that for any $i, j \in I$, we have $L_h^{\mathcal{E}} \circ p_{i,j} = L_h^{\mathcal{E}'} \circ p_{i,j}$. Now, we have that $\langle L_h^{\mathcal{E}} \circ e_i, 1_{E_i} \rangle = \lambda_{E_i}(x_i) \circ \alpha_i$, while $\langle L_h^{\mathcal{E}'} \circ e'_j, 1_{E'_j} \rangle = \lambda_{E'_j}(x'_j) \circ \beta_j$. Applying respectively f_i^* and f'_j^* to these two identities, we obtain that $\langle L_h^{\mathcal{E}} \circ p_{i,j}, 1_{F_{i,j}} \rangle = \lambda_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i)$ and $\langle L_h^{\mathcal{E}'} \circ p_{i,j}, 1_{F_{i,j}} \rangle = \lambda_{F_{i,j}}(x'_{i,j}) \circ f'_j{}^*(\beta_j)$, where $x_{i,j} = \mathcal{A}_{f_i}(x_i)$ and $x'_{i,j} = \mathcal{A}_{f'_j}(x'_j)$, from which it follows that the condition $L_h^{\mathcal{E}} \circ p_{i,j} = L_h^{\mathcal{E}'} \circ p_{i,j}$ is equivalent to the condition $\lambda_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i) = \lambda_{F_{i,j}}(x'_{i,j}) \circ f'_j{}^*(\beta_j)$. This condition can be proved by using condition (ii) of the theorem, the fact that the same identity holds with the cocone μ in place of λ and Remark 2.15(d). More specifically, the identities $\langle h \circ e_i, 1_{E_i} \rangle = \mu_{E_i}(x_i) \circ \alpha_i$ and $\langle h \circ e'_j, 1_{E'_j} \rangle = \mu_{E'_j}(x'_j) \circ \beta_j$ imply, applying respectively f_i^* and $f'_j{}^*$ to them, the identity $\mu_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i) = \mu_{F_{i,j}}(x'_{i,j}) \circ f'_j{}^*(\beta_j)$. From condition (ii) of the theorem, it thus follows that for any $i, j \in I \times J$, there exists an epimorphic family $\{g_k^{i,j} : G_k^{i,j} \rightarrow F_{i,j} \mid k \in K_{i,j}\}$ such that the pair $((\mathcal{A}_{g_k^{i,j}}(x_{i,j}), f_i^*(\alpha_i) \circ g_k^{i,j}), (\mathcal{A}_{g_k^{i,j}}(x'_{i,j}), f'_j{}^*(\beta_j) \circ g_k^{i,j}))$ belongs to the relation $\mathcal{R}_{G_k^{i,j}}^D$. This in turn implies (by Remark 2.15(d)) that $\lambda_{G_k^{i,j}}(\mathcal{A}_{g_k^{i,j}}(x_{i,j})) \circ (f_i^*(\alpha_i) \circ g_k^{i,j}) = \lambda_{G_k^{i,j}}(\mathcal{A}_{g_k^{i,j}}(x'_{i,j})) \circ (f'_j{}^*(\beta_j) \circ g_k^{i,j})$. But $\lambda_{G_k^{i,j}}(\mathcal{A}_{g_k^{i,j}}(x_{i,j})) \circ (f_i^*(\alpha_i) \circ g_k^{i,j}) = (V \times g_k^{i,j})^*(\lambda_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i))$ and $\lambda_{G_k^{i,j}}(\mathcal{A}_{g_k^{i,j}}(x'_{i,j})) \circ (f'_j{}^*(\beta_j) \circ g_k^{i,j}) = (V \times g_k^{i,j})^*(\lambda_{F_{i,j}}(x'_{i,j}) \circ f'_j{}^*(\beta_j))$, whence $(V \times g_k^{i,j})^*(\lambda_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i)) = (V \times g_k^{i,j})^*(\lambda_{F_{i,j}}(x'_{i,j}) \circ f'_j{}^*(\beta_j))$, equivalently

(since the $g_k^{i,j}$ are jointly epimorphic) $\lambda_{F_{i,j}}(x_{i,j}) \circ f_i^*(\alpha_i) = \lambda_{F_{i,j}}(x'_{i,j}) \circ f_j'^*(\beta_j)$, as required.

It is clear that the assignment $h \rightarrow L(h)$ defined above is natural in $E \in \mathcal{E}$; it therefore remains to show that the resulting arrow $l : U \rightarrow V$ satisfies the required property that for any object $E \in \mathcal{E}$ and any object x of the category $\mathcal{A}_E$, $(l \times 1_E) \circ \mu_E(x) = \lambda_E(x)$. Take s to be the canonical arrow $s : D_E(x) \rightarrow E$ in $\mathcal{E}$ and set $x' = \mathcal{A}_s(x)$, $E' = D_E(x)$; then $x \in \mathcal{A}_{E'}$ and, considered the pullback square

$$\begin{array}{ccc} D_{E'}(x') & \xrightarrow{r} & D_E(x) \\ \downarrow t & & \downarrow s \\ E' & \xrightarrow{s} & E, \end{array}$$

the unique arrow $\alpha : E' \rightarrow D_{E'}(x')$ such that $r \circ \alpha = 1_{E'}$ and $t \circ \alpha = 1_{E'}$ satisfies the property that $\langle h, 1_{E'} \rangle = \mu_{E'}(x') \circ \alpha$. So the epimorphic family $\{1_{E'} : E' \rightarrow E'\}$ satisfies condition (i) of the theorem with respect to the arrow $h = \pi_U \circ \mu_E(x)$, where $\pi_U : U \times E \rightarrow U$ is the canonical projection, and hence $l \circ h = L(h) = \lambda_{E'}(x') \circ \alpha$; therefore, $(l \times 1_E) \circ \mu_E(x) = \lambda_E(x)$, as required.

The proof of the theorem is now complete. $\square$

Remark 2.20. (a) The proof of the theorem shows that the sufficiency of the conditions of the theorem holds more in general for any (i.e., not necessarily small) $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$;

(b) The conditions in the statement of the theorem are both stable under localization; that is, if the cone μ over the diagram D satisfies them then for any $E \in \mathcal{E}$ the cone μ/E over the diagram D/E does.

Proposition 2.21. *Let $\underline{\mathcal{B}}_{\mathcal{E}}$ be a $\mathcal{E}$ -final subcategory of an indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$, $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ a $\mathcal{E}$ -indexed functor and μ a $\mathcal{E}$ -indexed cocone over D . Then μ satisfies the conditions of Theorem 2.19 with respect to the diagram D if and only if $\mu \circ i$ satisfies them with respect to the diagram $D \circ i$, where i is the canonical embedding $\underline{\mathcal{B}}_{\mathcal{E}} \hookrightarrow \underline{\mathcal{A}}_{\mathcal{E}}$.*

Proof Suppose that μ satisfies the conditions of Theorem 2.19 with respect to the diagram D . The fact that $\mu \circ i$ satisfies condition (i) of the theorem with respect to the diagram $D \circ i$ immediately follows from the fact that μ does with respect to the diagram D , by using the fact that for each object $x_i \in \mathcal{A}_{F_i}$ there exists an epimorphic family $\{g_k^i : G_k^i \rightarrow F_i \mid k \in K_i\}$ such that $\mathcal{A}_{g_k^i}(x_i)$ lies in $\mathcal{B}_{G_k^i}$ and for any arrow $\alpha_i : 1_{\mathcal{E}/F_i} \rightarrow D_{F_i}(x_i)$ in the topos $\mathcal{E}/F_i$, its image under the pullback functor g_k^{i*} is an arrow $1_{\mathcal{E}/G_k^i} \rightarrow D_{G_k^i}(\mathcal{A}_{g_k^i}(x_i))$ in the topos $\mathcal{E}/G_k^i$. Let us now show that $\mu \circ i$ satisfies condition (ii) with respect to the diagram $D \circ i$.

Let us first establish the following fact (*): for any pairs (x, y) and (x', y') in $\mathcal{I}_R^D$ (see the beginning of this section for the notation), if $((x, y), (x', y')) \in \mathcal{R}_R^D$, there exists an epimorphic family $\{r_i : R_i \rightarrow R \mid i \in I\}$ in $\mathcal{E}$ such that $((\mathcal{A}_{r_i}(x), r_i^*(y)), (\mathcal{A}_{r_i}(x'), r_i^*(y')))) \in \mathcal{R}_{R_i}^{D \circ i}$.

If $((x, y), (x', y')) \in \mathcal{R}_R^D$, there exists a finite sequence

$$(x, y) = (x_0, y_0), \dots, (x_n, y_n) = (x', y')$$

of pairs in $\mathcal{I}_R^D$ such that for any $j \in \{0, \dots, n-1\}$, either there exists an arrow $f_j : x_j \rightarrow x_{j+1}$ in $\mathcal{A}_R$ such that $y_{j+1} = D_R(f_j) \circ y_j$ or there exists an arrow $f_j : x_{j+1} \rightarrow x_j$ in $\mathcal{A}_R$ such that $y_j = D_R(f_j) \circ y_{j+1}$. By applying the definition of $\mathcal{E}$ -final subcategory a finite number of times (using the fact that the fibered product of epimorphic families is again an epimorphic family), we can find an epimorphic family $\{r_i : R_i \rightarrow R \mid i \in I\}$ in $\mathcal{E}$ such that for every $i \in I$ and any $j \in \{0, \dots, n-1\}$, the arrow $\mathcal{A}_{r_i}(f_j)$ lies in $\mathcal{B}_{R_i}$. From this it immediately follows that $((\mathcal{A}_{r_i}(x), r_i^*(y)), (\mathcal{A}_{r_i}(x'), r_i^*(y')))) \in \mathcal{R}_{R_i}^{D \circ i}$, as required.

Using fact (*), the proof of the fact that $\mu \circ i$ satisfies condition (ii) of the theorem with respect to the diagram $D \circ i$ follows straightforwardly from the fact that μ does with respect to the diagram D .

Conversely, let us suppose that $\mu \circ i$ satisfies the conditions of the theorem with respect to $D \circ i$ and deduce that μ does with respect to D . The fact that the validity of condition (i) for $\mu \circ i$ with respect to $D \circ i$ implies the validity of condition (i) for μ with respect to D is obvious. Concerning condition (ii) for (μ, D) , this can be deduced from condition (ii) for $(\mu \circ i, D \circ i)$ by using the fact that, given (x, y) and (x', y') as in the statement of the condition for (μ, D) , there exists an epimorphic family $\{e_j : E_j \rightarrow E \mid j \in J\}$ such that $\mathcal{A}_{e_j}(x)$ and $\mathcal{A}_{e_j}(x')$ lie in $\mathcal{B}_{E_j}$ for all $j \in J$.

The proof of the proposition is now complete. $\square$

Remark 2.22. The proposition shows in particular that (the necessity of the conditions of) Theorem 2.19 not only holds for $\mathcal{E}$ -indexed categories $\underline{\mathcal{A}}_{\mathcal{E}}$ which are $\mathcal{E}$ -small (i.e., which are equivalent to the $\mathcal{E}$ -externalization of an internal category in $\mathcal{E}$), but for any $\mathcal{E}$ -indexed category which is a $\mathcal{E}$ -final subcategory of a $\mathcal{E}$ -small $\mathcal{E}$ -indexed category (for instance, to the $\mathcal{E}$ -indexed categories of the form $\int F_{\mathcal{E}}$, for a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ - cf. Theorem 2.13).

Proposition 2.23. *Let $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ an indexed functor, where $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -filtered category, R an object of $\mathcal{E}$, $(x, y), (x', y')$ pairs in $\mathcal{I}_R^D$. Then there exists an epimorphic family $\{r_i : R_i \rightarrow R \mid i \in I\}$ in $\mathcal{E}$ such that for all $i \in I$, $((\mathcal{A}_{r_i}(x), r_i^*(y)), (\mathcal{A}_{r_i}(x'), r_i^*(y')))) \in \mathcal{R}_{R_i}^D$ if and only if there exists an epimorphic family $\{e_j : E_j \rightarrow R \mid j \in J\}$ in $\mathcal{E}$ with the property that for any $j \in J$ there exist arrows $f_j : \mathcal{A}_{e_j}(x) \rightarrow z$ and $g_j : \mathcal{A}_{e_j}(x') \rightarrow z$ in the category $\mathcal{A}_{E_j}$ such that $D_{E_j}(f_j) \circ y = D_{E_j}(g_j) \circ y'$.*

Proof The ‘if’ direction is obvious, so it remains to prove the ‘only if’ one. It clearly suffices to prove, by induction on the length of a sequence $(x_1, y_1), \dots, (x_n, y_n)$ of pairs in $\mathcal{I}_R^D$ with the property that for any $i \in \{1, \dots, n-1\}$, either there exists an arrow $f_j : x_j \rightarrow x_{j+1}$ in $\mathcal{A}_R$ such that $y_{j+1} = D_R(f_j) \circ y_j$ or there exists an arrow $f_j : x_{j+1} \rightarrow x_j$ in $\mathcal{A}_R$ such that $y_j = D_R(f_j) \circ y_{j+1}$, that there exists an epimorphic family $\{r_i : R_i \rightarrow R \mid i \in I\}$ in $\mathcal{E}$ such that for all $i \in I$, $((\mathcal{A}_{r_i}(x_1), r_i^*(y_1)), (\mathcal{A}_{r_i}(x_n), r_i^*(y_n))) \in \mathcal{R}_{R_i}^D$. The case $n = 1$ is obvious. Suppose now that the claim is valid for n and prove that it holds for $n + 1$. There exists an epimorphic family $\{e_j : E_j \rightarrow R \mid j \in J\}$ in $\mathcal{E}$ such that for any $j \in J$ there exist arrows $f_j : \mathcal{A}_{e_j}(x_1) \rightarrow z$ and $g_j : \mathcal{A}_{e_j}(x_n) \rightarrow z$ in the category $\mathcal{A}_{E_j}$ such that $D_{E_j}(f_j) \circ y_n = D_{E_j}(g_j) \circ y'_n$. On the other hand, there exists either an arrow $g : x_n \rightarrow x_{n+1}$ in $\mathcal{A}_R$ such that $D_R(g) \circ y_n = y_{n+1}$ or an arrow $h : x_{n+1} \rightarrow x_n$ in $\mathcal{A}_R$ such that $D_R(g) \circ y_{n+1} = y_n$. In the latter case, the thesis follows straightforwardly; so we can concentrate on the case when there exists an arrow $g : x_n \rightarrow x_{n+1}$ in $\mathcal{A}_R$ such that $D_R(g) \circ y_n = y_{n+1}$.

By using the definition of $\mathcal{E}$ -final subcategory, we obtain an epimorphic family $\{e_j : E_j \rightarrow R \mid j \in J\}$ in $\mathcal{E}$ and for each $j \in J$ two arrows $f_j : \mathcal{A}_{e_j}(x_1) \rightarrow z$ and $g_j : \mathcal{A}_{e_j}(x_n) \rightarrow z$ such that $D_{E_j}(f_j) \circ e_j^*(y_1) = D_{E_j}(g_j) \circ e_j^*(y_n)$. Similarly, we obtain, for each $j \in J$, an epimorphic family $\{f_k^j : F_k^j \rightarrow E_j \mid k \in K_j\}$ in $\mathcal{E}$ and for each $k \in K_j$ arrows $m_k^j : \mathcal{A}_{f_k^j}(z) \rightarrow w_k^j$ and $n_k^j : \mathcal{A}_{f_k^j}(\mathcal{A}_{e_j}(x_{n+1})) \rightarrow w_k^j$ such that $D_{F_k^j}(m_k^j) \circ D_{F_k^j}(\mathcal{A}_{f_k^j}(g_j)) \circ (f_k^j \circ e_j)^*(y_n) = D_{F_k^j}(n_k^j) \circ D_{F_k^j}(\mathcal{A}_{f_k^j}(g)) \circ (f_k^j \circ e_j)^*(y_{n+1})$. Now, by definition of $\mathcal{E}$ -final subcategory, for each $j \in J$ and $k \in K_j$ there exists an epimorphic family $\{g_l^{k,j} : G_l^{k,j} \rightarrow F_k^j \mid l \in L_{k,j}\}$ and for each $l \in L_{k,j}$ an arrow $p_l^{k,j} : \mathcal{A}_{g_l^{k,j}}(w_k^j) \rightarrow v_l^{k,j}$ such that $p_l^{k,j} \circ m_k^j \circ \mathcal{A}_{f_k^j}(g_j) = p_l^{k,j} \circ n_k^j \circ \mathcal{A}_{f_k^j}(\mathcal{A}_{e_j}(g))$. For any $j \in J, k \in K_j$ and $l \in L_{k,j}$, set $a_{j,k,l} : \mathcal{A}_{e_j \circ f_k^j \circ g_l^{k,j}}(x_1) \rightarrow v_l^{k,j}$ equal to $p_l^{k,j} \circ \mathcal{A}_{g_l^{k,j}}(m_k^j) \circ \mathcal{A}_{g_l^{k,j} \circ f_k^j}(f_j)$ and $b_{j,k,l} : \mathcal{A}_{e_j \circ f_k^j \circ g_l^{k,j}}(x_n) \rightarrow v_l^{k,j}$ equal to $p_l^{k,j} \circ \mathcal{A}_{g_l^{k,j}}(n_k^j) \circ \mathcal{A}_{g_l^{k,j} \circ f_k^j \circ e_j}(g)$. It is readily seen that $D_{G_l^{k,j}}(a_{j,k,l}) \circ (e_j \circ f_k^j \circ g_l^{k,j})^*(y_1) = D_{G_l^{k,j}}(b_{j,k,l}) \circ (e_j \circ f_k^j \circ g_l^{k,j})^*(y_{n+1})$, whence the required condition is satisfied (by taking as epimorphic family $\{e_j \circ f_k^j \circ g_l^{k,j} \mid j \in J, k \in K_j, l \in L_{k,j}\}$). $\square$

Combining the proposition with Theorem 2.19, we immediately obtain the following result.

Corollary 2.24. *Let $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ an indexed functor, where $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -filtered $\mathcal{E}$ -final subcategory of a small $\mathcal{E}$ -indexed category. Then a cocone μ over D with vertex U is a indexed colimiting cocone for D if and only if the following conditions are satisfied:*

- (i) For any object E of $\mathcal{E}$ and arrow $h : F \rightarrow U$ in the topos $\mathcal{E}$, there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object $x_i \in \mathcal{A}_{F_i}$ and an arrow $\alpha_i : 1_{\mathcal{E}/F_i} \rightarrow D_{F_i}(x_i)$ in the topos $\mathcal{E}/F_i$ such that $\langle h \circ f_i, 1_{F_i} \rangle = \mu_{F_i}(x_i) \circ \alpha_i$ as arrows $1_{\mathcal{E}/F_i} \rightarrow F_i^*(U)$ in $\mathcal{E}/F_i$;
- (ii) For any pairs (x, y) and (x', y') , where x and x' are objects of $\mathcal{A}_E$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$ and y' is an arrow $(f : F \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, we have that $\mu_E(x) \circ y = \mu_E(x') \circ y'$ if and only if there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ arrows $g_i : \mathcal{A}_{f \circ f_i}(x) \rightarrow z_i$ and $h_i : \mathcal{A}_{f \circ f_i}(x') \rightarrow z_i$ in the category $\mathcal{A}_{F_i}$ such that $D_{F_i}(g_i) \circ f_i^*(y) = D_{F_i}(h_i) \circ f_i^*(y')$.

Moreover, if μ is colimiting for D then the following ‘joint embedding property’ holds: for any pairs (x, y) and (x', y') , where x and x' are objects of $\mathcal{A}_E$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$ and y' is an arrow $(f : F \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, there exists an epimorphic family $\{e_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ arrows $g_i : \mathcal{A}_{f \circ e_i}(x) \rightarrow z_i$ and $h_i : \mathcal{A}_{f \circ e_i}(x') \rightarrow z_i$ in the category $\mathcal{A}_{F_i}$ such that $D_{F_i}(g_i) \circ f_i^*(y) = D_{F_i}(h_i) \circ f_i^*(y')$ and the diagram

$$\begin{array}{ccc}
D_{F_i}(\mathcal{A}_{f \circ e_i}(x)) & \xrightarrow{\xi_{(F_i, \mathcal{A}_{f \circ e_i}(x))}} & \\
D_{F_i}(g_i) \searrow & & \nearrow \xi_{(E_i, z_i)} \\
& D_{F_i}(z_i) & \longrightarrow \text{colim}(D) \times F_i \\
D_{F_i}(h_i) \nearrow & & \nwarrow \xi_{(F_i, \mathcal{A}_{f \circ e_i}(x'))} \\
D_{F_i}(\mathcal{A}_{f \circ e_i}(x')) & \xrightarrow{\xi_{(F_i, \mathcal{A}_{f \circ e_i}(x'))}} &
\end{array} ,$$

where $\xi_{(E, x)} : D_E(x) \rightarrow \text{colim}(D) \times E$ are the colimit arrows, commutes.

- Remark 2.25.** (a) In the statement of the corollary, we can suppose without loss of generality the object $f : F \rightarrow E$ of the topos $\mathcal{E}/E$ to be equal to the terminal $1_E : E \rightarrow E$. Indeed, by Remark 2.15, for any object x of $\mathcal{A}_E$ and any arrow $y : (f : F \rightarrow E) \rightarrow D_E(x)$ in the topos $\mathcal{E}/E$, $\mu_E(x) \circ y = \mu_E(x') \circ y'$ if and only if $\mu_F(\mathcal{A}_f(x)) \circ y_f = \mu_F(\mathcal{A}_f(x')) \circ y_{f'}$.
- (b) For any pairs (x, y) and (x', y') , where x and x' are objects of $\mathcal{A}_E$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$, y' is an arrow $(f : F \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, and any epimorphic family $\{g_j : G_j \rightarrow F \mid j \in J\}$ in $\mathcal{E}$, condition (ii) of Corollary 2.24 is satisfied by the pair $((x, y), (x', y'))$ if and only if it is satisfied by the pair $((x, y \circ g_j), (x', y' \circ g_j))$ for all $j \in J$.

Let us now apply Corollary 2.24 in the context of a flat functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ and a functor $P : \mathcal{C} \rightarrow \mathbf{Set}$, where $\mathcal{C}$ is a small category and $\mathcal{E}$ is a Grothendieck topos. Consider the restriction of the functor $(\overline{P_{\mathcal{E}} \circ \pi_{\overline{F}}^f}) \cong \overline{P_{\mathcal{E}} \circ \pi_{\overline{F}}^f}$ (where $P_{\mathcal{E}}$ is the internal diagram $[\mathbb{C}, \mathcal{E}]$ given by $\gamma_{\mathcal{E}}^* \circ \overline{P}$)

to the $\mathcal{E}$ -indexed subcategory $\int F_{\mathcal{E}}$ of $\int F_{\mathcal{E}}^f$. Recall that, by Theorem 2.14, the $\mathcal{E}$ -indexed functor $(\underline{P}_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f)$ is naturally isomorphic to the composite functor $\gamma_{\mathcal{E}} \circ z_{\mathcal{E}}$, where $z_{\mathcal{E}} : \int F_{\mathcal{E}} \rightarrow \mathbf{Set}_{\mathcal{E}}$ is the $\mathcal{E}$ -indexed functor defined by: $z_{\mathcal{E}E}((c, x)) = P(c)$ and $z_{\mathcal{E}E}(f) = P(f)$ (for any $E \in \mathcal{E}$, object (c, x) and arrow f in the category $\int F_E$).

As observed in section 2.5, a $\mathcal{E}$ -indexed cocone μ over this diagram with vertex U can be identified with a family of functions $\mu_{(c,x)} : P(c) \rightarrow \text{Hom}_{\mathcal{E}}(E, U)$ indexed by the pairs (c, x) consisting of an object c of $\mathcal{C}$ and a generalized element $x : E \rightarrow F(c)$ satisfying the following properties:

- (i) For any generalized element $x : E \rightarrow F(c)$ and any arrow $f : d \rightarrow c$ in $\mathcal{C}$, $\mu_{(c,x)} \circ P(f) = \mu_{(c, F(f) \circ x)}$;
- (ii) For any generalized element $x : E \rightarrow F(c)$ and any arrow $e : E' \rightarrow E$, $\mu_{(c, x \circ e)} = \text{Hom}_{\mathcal{E}}(e, U) \circ \mu_{(c,x)}$.

The following theorem characterizes the cocones μ which are colimiting.

Theorem 2.26. *Let $\mathcal{C}$ be a small category, $\mathcal{E}$ a Grothendieck topos, $P : \mathcal{C} \rightarrow \mathbf{Set}$ a functor and $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. Then a $\mathcal{E}$ -indexed cocone $\mu = \{\mu_{(c,x)} : P(c) \rightarrow \text{Hom}_{\mathcal{E}}(E, U) \mid c \in \mathcal{C}, x : E \rightarrow F(c) \text{ in } \mathcal{E}\}$ with vertex U over the diagram given by the restriction of the functor $(\underline{P}_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f) \cong \underline{P}_{\mathcal{E}} \circ \pi_{\overline{F}_{\mathcal{E}}}^f$ (where $\underline{P}_{\mathcal{E}}$ is the internal diagram $[\mathcal{C}, \mathcal{E}]$ given by $\gamma_{\mathcal{E}}^* \circ \overline{P}$) to the $\mathcal{E}$ -indexed subcategory $\int F_{\mathcal{E}}$ is colimiting if and only if the following conditions are satisfied:*

- (i) *For any generalized element $x : E \rightarrow U$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$, for each index $i \in I$ an object c_i in $\mathcal{C}$, a generalized element $x_i : E_i \rightarrow F(c_i)$ and an element $y_i \in P(c_i)$ such that $\mu_{(c_i, x_i)}(y_i) = x \circ e_i$;*
- (ii) *For any pairs (a, x) and (b, x') , where a and b are objects of $\mathcal{C}$ and $x : E \rightarrow F(a)$, $x' : E \rightarrow F(b)$ are generalized elements, and any elements $y \in P(a)$ and $y' \in P(b)$, $\mu_{(a,x)}(y) = \mu_{(b,x')}(y')$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ in $\mathcal{E}$ and for each index $i \in I$ an object c_i of $\mathcal{C}$, arrows $f_i : a \rightarrow c_i$, $g_i : b \rightarrow c_i$ in $\mathcal{C}$ and a generalized element $x_i : E_i \rightarrow F(c_i)$ such that $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$ for all $i \in I$ and $P(f_i)(y) = P(g_i)(y')$.*

Moreover, if μ is colimiting the following ‘joint embedding property’ holds: for any pairs (a, x) and (b, x') , where a and b are objects of $\mathcal{C}$ and $x : E \rightarrow F(a)$, $x' : E \rightarrow F(b)$ are generalized elements, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ an object c_i of $\mathcal{C}$, arrows $f_i : a \rightarrow c_i$, $g_i : b \rightarrow c_i$ in $\mathcal{C}$ and a generalized element $x_i : E_i \rightarrow F(c_i)$

such that $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$ for all $i \in I$ and (by Lemma 3.10) the following diagram commutes:

$$\begin{array}{ccc}
P(a) & \xrightarrow{\mu_{(a,x)}} & \text{Hom}_{\mathcal{E}}(E, U) \\
& \searrow P(f) & \searrow \text{Hom}_{\mathcal{E}}(e_i, M) \\
& & P(c_i) \xrightarrow{\mu_{(c_i, x_i)}} \text{Hom}_{\mathcal{E}}(E_i, U) \\
& \nearrow P(g) & \nearrow \text{Hom}_{\mathcal{E}}(e_i, U) \\
P(b) & \xrightarrow{\mu_{(b, x')}} & \text{Hom}_{\mathcal{E}}(E, U)
\end{array}$$

In fact, the epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ can be taken to be the pullback of the family of arrows $\langle F(f), F(g) \rangle : F(c) \rightarrow F(a) \times F(b)$ (for all spans $(f : a \rightarrow c, g : b \rightarrow c)$ in the category $\mathcal{C}$) along the arrow $\langle x, x' \rangle : E \rightarrow F(a) \times F(b)$.

Proof The theorem can be deduced from Corollary 2.24 by reasoning as follows. Concerning condition (i), its equivalence with condition (i) of Corollary 2.24 follows from Remark 2.25(a) and the fact that for any object (z, c) of the category $\int F_E$ and any arrow $\alpha : 1_{\mathcal{E}/E} \rightarrow D_E((z, c))$ in the topos $\mathcal{E}/E$, denoting by $!_{f_r} : f_r : F_r \rightarrow E \rightarrow 1_{\mathcal{E}/E}$ the pullback of the coproduct arrow $u_r : 1_{\mathcal{E}/E} \rightarrow \gamma_{\mathcal{E}/E}^*(P(c)) \cong D_E((z, c))$ in $\mathcal{E}/E$ (for each $r \in P(c)$), the arrows $!_{f_r}$ define, for $r \in P(c)$, an epimorphic family in $\mathcal{E}/E$ on the object $1_{\mathcal{E}/E}$; indeed, this shows that we can suppose without loss of generality α to factor through one of the coproduct arrows u_r and hence to correspond to an element $y \in P(c)$.

Concerning condition (ii), by Remark 2.25 we can suppose without loss of generality that the pairs (x, y) and (x', y') , where x (resp. x') is an object of $\int F_E$, y is an arrow $(f : F \rightarrow E) \rightarrow D_E(x)$ in $\mathcal{E}/E$ and y' is an arrow $(f' : F' \rightarrow E) \rightarrow D_E(x')$ in $\mathcal{E}/E$, have respectively the form $((x, c), y)$, where $y = u_r \circ !_{f_r : F_r \rightarrow E}$ and $((x', c'), y')$, where $y' = u_{r'} \circ !_{f_{r'} : F_{r'} \rightarrow E}$ (with the above notation), for some $r, r' \in P(c)$. Thus we have that $\mu_E(x) \circ y = \mu_E(x') \circ y'$ if and only if $(\text{Hom}_{\mathcal{E}}(f_r, U) \circ \mu_{(x, c)})(r) = (\text{Hom}_{\mathcal{E}}(f_{r'}, U) \circ \mu_{(x', c')})(r')$; but $(\text{Hom}_{\mathcal{E}}(f_r, U) \circ \mu_{(x, c)})(r) = \mu_{(c, x \circ f_r)}(r)$ and $(\text{Hom}_{\mathcal{E}}(f_{r'}, U) \circ \mu_{(x', c')})(r') = \mu_{(c', x' \circ f_{r'})}(r')$, whence $(\text{Hom}_{\mathcal{E}}(f_r, U) \circ \mu_{(x, c)})(r) = (\text{Hom}_{\mathcal{E}}(f_{r'}, U) \circ \mu_{(x', c')})(r')$ if and only if $\mu_{(c, x \circ f_r)}(r) = \mu_{(c', x' \circ f_{r'})}(r')$. $\square$

2.7 Explicit calculation of set-indexed colimits

Recall from [26] (cf. p. 355) that the colimit of a functor $H : \mathcal{I} \rightarrow \mathcal{E}$ with values in a Grothendieck topos $\mathcal{E}$ can always be realized as the coequalizer $q : \coprod_{i \in I} H(i) \rightarrow \text{colim}(H)$ of the pair of arrows

$$a, b : \coprod_{u: i \rightarrow j \text{ in } \mathcal{I}} H(i) \rightarrow \coprod_{i \in I} H(i)$$

defined by:

$$a \circ \lambda_u = \kappa_i$$

and

$$b \circ \lambda_u = \kappa_j \circ H(u)$$

(for every arrow $u : i \rightarrow j$ in $\mathcal{I}$), where $\lambda_u : H(\text{dom}(u)) \rightarrow \coprod_{u:i \rightarrow j \text{ in } \mathcal{I}} H(i)$ (for any $u : i \rightarrow j$ in $\mathcal{I}$) and $\kappa_i : H(i) \rightarrow \coprod_{i \in I} H(i)$ (for any $i \in \mathcal{I}$) are the canonical coproduct arrows.

Given two arrows $s, t : E \rightarrow \coprod_{i \in I} H(i)$, for every objects $i, j \in \mathcal{I}$, we can consider the following pullback diagrams:

$$\begin{array}{ccc} E_i^s & \xrightarrow{a_i^s} & H(i) \\ \downarrow p_i^s & & \downarrow \kappa_i \\ E & \xrightarrow{s} & \coprod_{i \in I} H(i) \end{array} \quad \begin{array}{ccc} E_j^t & \xrightarrow{a_j^t} & H(j) \\ \downarrow p_j^t & & \downarrow \kappa_j \\ E & \xrightarrow{t} & \coprod_{i \in I} H(i) \end{array}$$

Let us also consider, for each $i, j \in \mathcal{J}$, the pullback square

$$\begin{array}{ccc} E_{i,j}^{s,t} & \xrightarrow{q_{i,j}^s} & E_i^s \\ \downarrow q_{i,j}^t & & \downarrow p_i^s \\ E_j^t & \xrightarrow{p_j^t} & E \end{array}$$

and denote by $r_{i,j}^{s,t} : E_{i,j}^{s,t} \rightarrow E$ the arrow $p_i^s \circ q_{i,j}^s = p_j^t \circ q_{i,j}^t$. Notice that the family of arrows $\{r_{i,j}^{s,t} \mid i, j \in \mathcal{I}\}$ is epimorphic.

We shall use these notations throughout the section.

The following lemma provides a characterization of the coequalizer of the functor H , which will be useful to us later on.

Lemma 2.27. *Let $p : \coprod_{i \in I} H(i) \rightarrow A$ be an epimorphism. Then p is isomorphic to the canonical map $q : \coprod_{i \in I} H(i) \rightarrow \text{colim}(H)$ if and only if for any object E of $\mathcal{E}$, any objects $i, j \in \mathcal{I}$ and any generalized elements $z : E \rightarrow H(i)$ and $w : E \rightarrow H(j)$, $(\kappa_i \circ z, \kappa_j \circ w) \in R_E$ if and only if $p \circ z = p \circ w$.*

Proof As in a topos every epimorphism is the coequalizer of its kernel pair, any two epimorphisms $\alpha : B \rightarrow C$ and $\beta : B \rightarrow D$ in $\mathcal{E}$ are isomorphic in $B/\mathcal{E}$ if and only if their kernel pairs are isomorphic as subobjects of $B \times B$. In particular, p is isomorphic to q if and only if the kernel pair of p is isomorphic to R .

The kernel pair S of p satisfies the property that for any generalized elements $s, t : E \rightarrow \coprod_{i \in I} H(i)$, $(s, t) \in S_E$ if and only if $p \circ s = p \circ t$. It thus remains to prove that the following two conditions are equivalent:

- (1) for any generalized elements $s, t : E \rightarrow \coprod_{i \in I} H(i)$, $p \circ s = p \circ t$ if and only if $(s, t) \in R_E$;
- (2) for any object E of $\mathcal{E}$, any objects $i, j \in \mathcal{I}$ and any generalized elements $z : E \rightarrow H(i)$ and $w : E \rightarrow H(j)$, $p \circ \kappa_i \circ z = p \circ \kappa_j \circ w$ if and only if $(\kappa_i \circ z, \kappa_j \circ w) \in R_E$.

It is clear that (1) implies (2). To prove the converse direction, we observe that, as we remarked above, $(s, t) \in R_E$ if and only if for every $i, j \in \mathcal{I}$ $(\kappa_i \circ a_i^s \circ q_{i,j}^s, \kappa_j \circ a_j^t \circ q_{i,j}^t) \in R_{E_{i,j}^{s,t}}$; but the latter condition is equivalent, under assumption (2), to the requirement that for every $i, j \in \mathcal{I}$, $p \circ \kappa_i \circ a_i^s \circ q_{i,j}^s = p \circ \kappa_j \circ a_j^t \circ q_{i,j}^t$, where $s \circ r_{i,j}^{s,t} = \kappa_i \circ a_i^s \circ q_{i,j}^s$ and $t \circ r_{i,j}^{s,t} = \kappa_j \circ a_j^t \circ q_{i,j}^t$, that is, as the family of arrows $r_{i,j}^{s,t}$ is epimorphic, to the condition $p \circ s = p \circ t$, as required. $\square$

Lemma 2.28. *Let B be an object of a topos $\mathcal{E}$. Then an equivalence relation in $\mathcal{E}$ on B can be identified with an assignment $E \rightarrow R_E$ of an equivalence relation on the set $\text{Hom}_{\mathcal{E}}(E, B)$ satisfying the following properties:*

- (i) *For any arrow $f : E \rightarrow E'$ in $\mathcal{E}$, if $(h, k) \in R_{E'}$ then $(h \circ f, k \circ f) \in R_E$;*
- (ii) *For any epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and any arrows $f, g \in \text{Hom}_{\mathcal{E}}(E, B)$, $(f \circ e_i, g \circ e_i) \in R_{E_i}$ for all $i \in I$ implies $(f, g) \in R_E$.*

Proof Clearly, for any object A of $\mathcal{C}$, the subobjects of A can be identified with the $J_{\mathcal{E}}$ -closed sieves on A , or equivalently with the $c_{J_{\mathcal{E}}}$ -closed subobjects of the representable $\text{Hom}_{\mathcal{E}}(-, A)$, where $c_{J_{\mathcal{E}}}$ is the closure operation on subobjects corresponding to the canonical Grothendieck topology $J_{\mathcal{E}}$ on $\mathcal{E}$. Applying this remark to $A = B \times B$ and noticing that the concept of equivalence relation is cartesian and hence preserved and reflected by cartesian fully faithful functors, in particular by the Yoneda embedding $\mathcal{E} \rightarrow [\mathcal{E}^{\text{op}}, \mathbf{Set}]$, we immediately deduce the thesis. $\square$

Lemma 2.28 implies that $(s, t) \in R_E$ if and only if for every $i, j \in \mathcal{I}$ $(\kappa_i \circ a_i^s \circ q_{i,j}^s, \kappa_j \circ a_j^t \circ q_{i,j}^t) \in R_{E_{i,j}^{s,t}}$; indeed, $s \circ r_{i,j}^{s,t} = \kappa_i \circ a_i^s \circ q_{i,j}^s$ and $t \circ r_{i,j}^{s,t} = \kappa_j \circ a_j^t \circ q_{i,j}^t$.

This shows that, in order to completely describe the relation R , it suffices to consider elements of the form $\kappa_i \circ z$ and $\kappa_j \circ w$, where $z : E \rightarrow H(i)$ and $w : E \rightarrow H(j)$, and characterize when they belong to R_E .

Lemma 2.29. *Let $\mathcal{E}$ be a topos, $f, g : A \rightarrow B$ arrows in $\mathcal{E}$ with coequalizer $q : B \rightarrow C$ and $R \rightrightarrows B \times B$ the kernel pair of q . Then for any object E of $\mathcal{E}$ and any elements $h, k \in \text{Hom}_{\mathcal{E}}(E, B)$, $\langle h, k \rangle : E \rightarrow B \times B$ factors through R if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ such that for each $i \in I$ $(h \circ e_i, k \circ e_i)$ belongs to the equivalence relation on the set $\text{Hom}_{\mathcal{E}}(E_i, B)$ generated by the set of pairs of the form $(f \circ a, g \circ a)$, for a generalized element $a : E_i \rightarrow A$ in $\mathcal{E}$.*

Proof For any object E of $\mathcal{E}$, consider the equivalence relation $R_E^{f,g}$ on the set $\text{Hom}_{\mathcal{E}}(E, B)$ consisting of the pairs (h, k) with the property that there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ such that for each $i \in I$ $(h \circ e_i, k \circ e_i)$ belongs to the equivalence relation on the set $\text{Hom}_{\mathcal{E}}(E_i, B)$ generated by the set of pairs of the form $(f \circ a, g \circ a)$, for a generalized element $a : E_i \rightarrow A$ in $\mathcal{E}$. The assignment $E \rightarrow R_E^{f,g}$ clearly satisfies the conditions of Lemma 2.28 and hence defines an equivalence relation on B in $\mathcal{E}$. To prove that $R = R^{f,g}$, we argue as follows. Denoting by $\pi_1, \pi_2 : R^{f,g} \rightarrow B$ the canonical projections, we have that for any $E \in \mathcal{E}$, $\text{Hom}(E, q)$ coequalizes the arrows $\text{Hom}(E, \pi_1)$ and $\text{Hom}(E, \pi_2)$; therefore q coequalizes π_1 and π_2 in $\mathcal{E}$. To prove that q is actually the coequalizer of π_1 and π_2 in $\mathcal{E}$, we observe that any arrow $p : B \rightarrow D$ such that $p \circ \pi_1 = p \circ \pi_2$ satisfies $p \circ f = p \circ g$ and hence, by definition of q , there exists a unique factorization of p through q . As q is the coequalizer of $R^{f,g}$ and in a topos all equivalence relations are the kernel pairs of their coequalizers, we conclude that $R^{f,g} = R$, as required. $\square$

Remark 2.30. Under the hypotheses of the lemma, R can be characterized as the equivalence relation on B generated by the arrows f and g , i.e. as the smallest equivalence relation on B containing the image of the arrow $\langle f, g \rangle : A \rightarrow B \times B$. Indeed, clearly R contains the image of this arrow, and if T is an equivalence relation on B containing the image of the arrow $\langle f, g \rangle$ then the coequalizer z of T factors through q and hence the kernel pair of q , namely R , is contained in the kernel pair of z , namely T .

Recall that the internal language $\Sigma_{\mathcal{E}}$ of a topos $\mathcal{E}$ is the first-order signature consisting of one sort $\ulcorner E \urcorner$ for any object E of $\mathcal{E}$, one function symbol $\ulcorner f \urcorner : \ulcorner A \urcorner \rightarrow \ulcorner B \urcorner$ for any arrow $f : A \rightarrow B$ in $\mathcal{E}$ and a relation symbol $\ulcorner R \urcorner \rightrightarrows \ulcorner A_1 \urcorner, \dots, \ulcorner A_n \urcorner$ for each subobject $R \rightrightarrows A_1 \times \dots \times A_n$ in $\mathcal{E}$. There is a ‘tautological’ $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$ in $\mathcal{E}$, obtained by interpreting each sort as the corresponding object, each function symbol as the corresponding arrow and each relation symbol as the corresponding subobject.

We shall use the notation $x^{\ulcorner C \urcorner}$ to denote a variable x of sort C , and will omit the superscript when it can be unambiguously inferred from the context.

Lemma 2.29 can be reformulated in logical terms as follows.

Lemma 2.31. *Let $\mathcal{E}$ be a topos, $f, g : A \rightarrow B$ arrows in $\mathcal{E}$ with coequalizer $q : B \rightarrow C$ and R the kernel pair of q . Let $(z^B, z'^B) \in G_{f,g}$ be an abbreviation for the formula*

$$z = z' \vee (\exists x)(\ulcorner f \urcorner(x) = z \wedge \ulcorner g \urcorner(x) = z') \vee (\exists x')(\ulcorner f \urcorner(x') = z' \wedge \ulcorner g \urcorner(x') = z)$$

over $\Sigma_{\mathcal{E}}$. Then the geometric bi-sequent

$$(\ulcorner R \urcorner(y, y') \dashv\vdash_{y^B, y'^B} \bigvee_{n \in \mathbb{N}} (\exists z_1 \dots \exists z_n (z_1 = x \wedge z_n = x' \wedge \bigwedge_{1 \leq i < n} (z_{i-1}, z_i) \in G_{f,g})))$$

is valid in the $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$.

Proof The interpretation of the formula $(z^B, z'^B) \in G_{f,g}$ is the symmetric and reflexive closure of the relation $S_{f,g}$ on B given by the image of the arrow $\langle f, g \rangle : A \rightarrow B \times B$, that is the union of the diagonal subobject $\Delta : B \rightarrow B \times B$ of $B \times B$, $S_{f,g} \rightarrow B \times B$ and the subobject $\tau \circ S_{f,g}$ of $B \times B$ given by the composite of $S_{f,g}$ with the exchange isomorphism $\tau : B \times B \rightarrow B \times B$. The interpretation of the formula $\bigvee_{n \in \mathbb{N}} (\exists z_1 \dots \exists z_n (z_1 = x \wedge z_n = x' \wedge \bigwedge_{1 \leq i < n} (z_{i-1}, z_i) \in G_{f,g}))$ thus coincides with the equivalence relation on B generated by $S_{f,g}$, that is with R (cf. Remark 2.30). $\square$

Given a functor $H : \mathcal{I} \rightarrow \mathcal{E}$ from a small category $\mathcal{I}$ to a Grothendieck topos $\mathcal{E}$ and an object E of $\mathcal{E}$, we define $\mathcal{I}_H^E$ as the set of pairs (i, x) , where i is an object of $\mathcal{I}$ and x is a generalized element $E \rightarrow H(i)$. In the sequel we shall occasionally identify a generalized element $x : E \rightarrow H(i)$ with its composite $\kappa_i \circ x$ with the coproduct arrow κ_i .

The following proposition can be established by means of arguments similar to those employed in the proof of Lemma 2.29.

Proposition 2.32. *Let $H : \mathcal{I} \rightarrow \mathcal{E}$ be a functor from a small category $\mathcal{I}$ to a Grothendieck topos $\mathcal{E}$. Then the equivalence relation R on the object $\coprod_{i \in I} H(i)$ given by the kernel pair of the canonical arrow $\coprod_{i \in I} H(i) \rightarrow \text{colim}(H)$ satisfies the following property: for any object E of $\mathcal{E}$ and any pairs (i, x) and (i', x') in $\mathcal{I}_H^E$, we have that $(\kappa_i \circ x, \kappa_{i'} \circ x') \in R_E$ if and only if there exists an epimorphic family $\{e_k : E_k \rightarrow E \mid k \in K\}$ in $\mathcal{E}$ such that for every $k \in K$, the pair $((i, x \circ e_k), (i', x' \circ e_k))$ belongs to the equivalence relation on the set $\mathcal{I}_H^{E_k}$ generated by the pairs of the form $((j, y), (j', H(f) \circ y))$, where $f : j \rightarrow j'$ is an arrow in $\mathcal{I}$ and y is a generalized element $E_k \rightarrow H(j)$.*

Proof Let us consider the pair of arrows $a, b : \coprod_{u: i \rightarrow j \text{ in } \mathcal{I}} H(i) \rightarrow \coprod_{i \in I} H(i)$ defined at the beginning of section 2.7. The canonical arrow $\coprod_{i \in I} H(i) \rightarrow \text{colim}(H)$ is the coequalizer of a and b .

For any object E of $\mathcal{E}$, let R_E be the relation on the set $\text{Hom}_{\mathcal{E}}(E, \coprod_{i \in I} H(i))$ consisting of the pairs (x, x') with the property that for any $i, j \in I$ there exists an epimorphic family $\{e_k^{(i,j)} : E_k^{(i,j)} \rightarrow E_{i,j}^{x,x'} \mid k \in K_{(i,j)}\}$ in $\mathcal{E}$ such that for each $k \in K_{(i,j)}$, $(x \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)}, x \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)})$ belongs to the equivalence relation $T_k^{(i,j)}$ on the set $\mathcal{I}_H^{E_k^{(i,j)}}$ generated by the set of pairs of the form $((n, y), (m, H(z) \circ y))$ (for $n \in I$, $y : E_k^{(i,j)} \rightarrow H(n)$ and $z : n \rightarrow m$ in I).

It is readily seen that the assignment $E \rightarrow R_E$ satisfies the hypotheses of Lemma 2.28 (notice that the fibered product of two epimorphic families is again an epimorphic family), whence it defines an equivalence relation R in $\mathcal{E}$ on the object $\coprod_{i \in I} H(i)$ such that for any generalized elements $x, x' : E \rightarrow \coprod_{i \in I} H(i)$, $\langle x, x' \rangle$ factors through R if and only if $(x, x') \in R_E$.

The arrow $\langle a, b \rangle$ factors through R since for every $i, j \in I$, $(a \circ r_{i,j}^{s,t}, b \circ r_{i,j}^{s,t})$ belongs to the equivalence relation $T_k^{(i,j)}$.

By Remark 2.30, R thus contains the kernel pair of q . The converse inclusion follows from the fact that for any $(x, x') \in R_E$, $q \circ x = q \circ x'$; indeed, if $\{e_k^{(i,j)} : E_k^{(i,j)} \rightarrow E_{i,j}^{x,x'} \mid k \in K_{(i,j)}\}$ in $\mathcal{E}$ is an epimorphic family such that for each $k \in K_{(i,j)}$, $(x \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)}, x \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)})$ belongs to the equivalence relation $T_k^{(i,j)}$ and hence we have $q \circ x \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)} = q \circ x' \circ r_{i,j}^{x,x'} \circ e_k^{(i,j)}$ for all $k \in K_{(i,j)}$, that is $q \circ x = q \circ x'$. $\square$

Remark 2.33. The proposition could be alternatively deduced as a consequence of Lemma 2.29, but the proof of such implication would be more involved than the direct one that we have given above.

In the case of filtered indexing categories, the description of colimits given above simplifies. We shall be concerned in particular with functors of the form $F \circ \pi_P$, where F is a flat functor with values in a Grothendieck topos and π_P is the fibration associated to a set-valued functor P .

Recall from chapter VII of [26] that a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ is flat if and only if it is filtering, i.e. the following conditions are satisfied:

- (i) For any object E of $\mathcal{E}$ there exists some epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index i an object b_i of $\mathcal{C}$ and a generalized element $E_i \rightarrow F(b_i)$ in $\mathcal{E}$;
- (ii) For any two objects c and d in $\mathcal{C}$ and any generalized element $(x, y) : E \rightarrow F(c) \times F(d)$ in $\mathcal{E}$ there is an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index i an object b_i of $\mathcal{C}$ with arrows $u_i : c \rightarrow b_i$ and $v_i : d \rightarrow b_i$ in $\mathcal{C}$ and a generalized element $z_i : E_i \rightarrow F(b_i)$ in $\mathcal{E}$ such that $\langle F(u_i), F(v_i) \rangle \circ z_i = \langle x, y \rangle \circ e_i$;

- (iii) For any two parallel arrows $u, v : d \rightarrow c$ in $\mathcal{C}$ and any generalized element $x : E \rightarrow F(c)$ in $\mathcal{E}$ for which $F(u) \circ x = F(v) \circ x$, there is an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index i an arrow $w_i : c \rightarrow b_i$ and a generalized element $y_i : E_i \rightarrow F(b_i)$ such that $w_i \circ u = w_i \circ v$ and $F(w_i) \circ y_i = x \circ e_i$.

Notice that can suppose $E = 1_{\mathcal{E}}$ in condition (i) without loss of generality and if all the arrows in the category $\mathcal{C}$ are monic then condition (iii) rewrites as follows: for any two parallel arrows $u, v : D \rightarrow C$ in $\mathcal{C}$, either $u = v$ or the equalizer of $F(u)$ and $F(v)$ is zero.

Proposition 2.34. *Let $\mathcal{C}$ be a small category, $\mathcal{E}$ a Grothendieck topos, $P : \mathcal{C} \rightarrow \mathbf{Set}$ a functor and $F : \mathcal{C}^{op} \rightarrow \mathcal{E}$ a flat functor. Let $\pi_P : \int P \rightarrow \mathcal{C}^{op}$ be the canonical projection from the category of elements of P to $\mathcal{C}^{op}$. Then*

$$\text{colim}(F \circ \pi_P) \cong \left(\coprod_{(c,x) \in \int P} F(c) \right) / R,$$

where R is the equivalence relation in $\mathcal{E}$ defined by saying that, for any objects (c, x) and (d, y) of the category $\int P$, the geometric bi-sequent

$$\bigvee_{\substack{c \xrightarrow{f} a \xrightarrow{g} d \\ P(f)(x) = P(g)(y)}} (\ulcorner R^{\ulcorner \xi_{(c,x)} \urcorner}(u), \ulcorner \xi_{(d,y)} \urcorner(v) \rrcorner \Vdash_{u^{F(c)}, v^{F(d)}} (\exists \xi^{F(a)}) (\ulcorner F(f) \urcorner(\xi) = u \wedge \ulcorner F(g) \urcorner(\xi) = v))$$

is valid in the $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$.

In particular, for any objects $(c, x), (c, y)$ of the category $\int P$, the geometric bi-sequent

$$(\ulcorner R^{\ulcorner \xi_{(c,x)} \urcorner}(u), \ulcorner \xi_{(c,y)} \urcorner(u) \rrcorner \Vdash_{u^{F(c)}} \bigvee_{\substack{c \xrightarrow{f} a \\ P(f)(x) = P(f)(y)}} (\exists \xi^{F(a)}) (\ulcorner F(f) \urcorner(\xi) = u))$$

is valid in the $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$.

Proof Let $\coprod_{(c,x) \in \int P} F(c)$ be the coproduct in $\mathcal{E}$ of the functor $F \circ \pi_P$, with canonical coproduct arrows $\xi_{(c,x)} : F(c) \rightarrow \coprod_{(c,x) \in \int P} F(c)$, and let T be the equivalence relation on $\coprod_{(c,x) \in \int P} F(c)$ corresponding to the quotient $\coprod_{(c,x) \in \int P} F(c) \rightarrow \text{colim}(F \circ \pi_P)$. Then we have $\text{colim}(F \circ \pi_P) \cong (\coprod_{(c,x) \in \int P} F(c)) / R$, and by Lemma 2.29, the relation R is the equivalence relation on the coproduct $\coprod_{(c,x) \in \int P} F(c)$ generated by the image of the arrow $\langle a, b \rangle$ where $s, t : \coprod_{z : (c,x) \rightarrow (d,y) \text{ in } \int P} F(c) \rightarrow \coprod_{(c,x) \in \int P} F(c)$ are the arrows defined above.

Let $\lambda_z : F(c) \rightarrow \coprod_{z:(c,x) \rightarrow (d,y) \text{ in } \int P} F(c)$ be the canonical coproduct arrows (for each arrow $z : (c, x) \rightarrow (d, y)$ in the category $\int P$). Notice that for any arrow $z : (c, x) \rightarrow (d, y)$ in $\int P$ the sequent $\top \vdash_{u'F(c)} (\ulcorner a \urcorner (\ulcorner \lambda_z \urcorner (u')) , \ulcorner b \urcorner (\ulcorner \lambda_z \urcorner (u')))) \in E_{(c,x),(d,y)}$ is valid in $\mathcal{S}_{\mathcal{E}}$.

The fact that the right-to-left implication of the bi-sequent in the statement of the proposition holds in $\mathcal{S}_{\mathcal{E}}$ is clear.

For any objects $(c, x), (d, y)$ in $\int P$, let the expression $(u^{F(c)}, v^{F(d)}) \in E_{(c,x),(d,y)}$ be an abbreviation of the formula
$$\bigvee_{\substack{c \xrightarrow{f} a \xrightarrow{g} d \mid \\ P(f)(x) = P(g)(y)}} (\exists \xi^{F(a)}) (\ulcorner F(f) \urcorner (\xi) = u \wedge \ulcorner F(g) \urcorner (\xi) = v).$$

The validity of the left-to-right implication will clearly follow from Lemma 2.31 once we have shown that for any objects $(c, x), (d, y), (e, z)$ in $\int P$ the sequent

$$(u, v) \in E_{(c,x),(d,y)} \wedge (v, w) \in E_{(d,y),(e,z)} \vdash_{u^{F(c)}, v^{F(d)}, z^{F(e)}} (u, w) \in E_{(c,x),(e,z)}$$

is valid in the $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$.

For simplicity we shall give the proof in the case $\mathcal{E} = \mathbf{Set}$, but all of our arguments are formalizable in the internal logic of the topos and hence are valid in general. In fact, the lift of the proof from the set-theoretic to the topos-theoretic setting is made possible by the following logical characterization of flatness for a functor; specifically, the fact that a functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ is flat can be expressed in terms of the validity of the following sequents in the structure $\mathcal{S}_{\mathcal{E}}$:

$$\begin{aligned} & (\top \vdash_{\square} \bigvee_{c \in \text{Ob}(\mathcal{C})} (\exists x^{F(c)}) (x = x)); \\ & (\top \vdash_{x^{F(a)}, y^{F(b)}} \bigvee_{a \xrightarrow{f} c \xleftarrow{g} b} (\exists z^{F(c)}) (\ulcorner F(f) \urcorner (z) = x \wedge \ulcorner F(g) \urcorner (z) = y)) \end{aligned}$$

for any objects a, b of $\mathcal{C}$;

$$(\ulcorner F(f) \urcorner (x) = \ulcorner F(g) \urcorner (x) \vdash_{x^{F(a)}} \bigvee_{h:a \rightarrow c \mid h \circ f = h \circ g} (\exists z^{F(c)}) (\ulcorner F(h) \urcorner (z) = x))$$

for any pair of arrows $f, g : b \rightarrow a$ in $\mathcal{C}$ with common domain and codomain.

Now, given objects $(c, x), (d, y), (e, z) \in \int P$ and elements $u \in F(c), v \in F(d), w \in F(e)$, suppose that $(u, v) \in E_{(c,x),(d,y)}$ and $(v, w) \in E_{(d,y),(e,z)}$; we want to prove that $(u, w) \in E_{(c,x),(e,z)}$.

As $(u, v) \in E_{(c,x),(d,y)}$, there exist arrows $f : c \rightarrow a$ and $g : d \rightarrow a$ in $\mathcal{C}$ and an element $\xi \in F(a)$ such that $F(f)(\xi) = u$, $F(g)(\xi) = v$, $P(f)(x) = P(g)(y)$. Similarly, as $(v, w) \in E_{(d,y),(e,z)}$, there exist arrows $h : d \rightarrow b$ and $k : e \rightarrow b$ in $\mathcal{C}$ and an element $\chi \in F(b)$ such that $F(h)(\chi) = v$, $F(k)(\chi) = w$ and $P(h)(y) = P(k)(z)$.

Using condition (ii) in the definition of flat functor, we obtain the existence of an object m of $\mathcal{C}$, arrows $r : a \rightarrow m$ and $t : b \rightarrow m$ in $\mathcal{C}$ and an

element $\epsilon \in F(m)$ such that $F(r)(\epsilon) = \xi$ and $F(t)(\epsilon) = \chi$. Now, consider the arrows $r \circ g, t \circ h : d \rightarrow m$; we have that $F(r \circ g)(\epsilon) = F(t \circ h)(\epsilon)$ and hence by condition (iii) in the definition of flat functor there exists an arrow $s : m \rightarrow n$ in $\mathcal{C}$ such that $s \circ r \circ g = s \circ t \circ h$ and an element $\alpha \in F(n)$ such that $F(s)(\alpha) = \epsilon$.

Consider the arrows $s \circ r \circ f$ and $s \circ t \circ k$. They satisfy the necessary conditions to ensure that $(u, w) \in E_{(c,x),(e,z)}$, i.e. $F(s \circ r \circ f)(\alpha) = u$, $F(s \circ t \circ k)(\alpha) = w$, $P(s \circ r \circ f)(x) = P(s \circ t \circ k)(z)$. Indeed, $F(s \circ r \circ f)(\alpha) = F(f)(F(r)(F(s)(\alpha))) = F(f)(F(r)(\epsilon)) = F(f)(\xi) = u$, $F(s \circ t \circ k)(\alpha) = F(k)(F(t)(F(s)(\alpha))) = F(k)(F(t)(\epsilon)) = F(k)(\chi) = w$ and $P(s \circ r \circ f)(x) = P(s \circ r)(P(f)(x)) = P(s \circ r)(P(g)(y)) = P(s \circ r \circ g)(y) = P(s \circ t \circ h)(y) = P(s \circ t)(P(h)(y)) = P(s \circ t)(P(k)(z)) = P(s \circ t \circ k)(z)$, as required.

The second part of the proposition follows from the first by an easy application of condition (iii) in the definition of flat functor. $\square$

The following proposition represents the translation of Proposition 2.34 in the categorical language of generalized elements.

Proposition 2.35. *Let $\mathcal{C}$ be a small category, $\mathcal{E}$ a Grothendieck topos with a separating set S , $P : \mathcal{C} \rightarrow \mathbf{Set}$ a functor and $F : \mathcal{C}^{op} \rightarrow \mathcal{E}$ a flat functor. Let $\pi_P : \int P \rightarrow \mathcal{C}^{op}$ be the canonical projection from the category of elements of P to $\mathcal{C}^{op}$. Then the equivalence relation R defined by the formula*

$$\text{colim}(F \circ \pi_P) \cong \left(\coprod_{(c,x) \in \int P} F(c) \right) / R$$

admits the following characterization: for any objects $(c, x), (d, y)$ of $\int P$ and generalized elements $u : E \rightarrow F(c)$, $v : E \rightarrow F(d)$ (where $E \in S$), $(\xi_{(c,x)} \circ u, \xi_{(d,y)} \circ v) \in R_E$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ an object $a_i \in \mathcal{C}$, a generalized element $h_i : E_i \rightarrow F(b_i)$ and two arrows $f_i : c \rightarrow a_i$ and $f'_i : d \rightarrow a_i$ in $\mathcal{C}$ such that $P(f_i)(x) = P(f'_i)(y)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle u, v \rangle \circ e_i$.

In particular, for any objects (c, x) and (c, y) of $\int P$ and any generalized element $u : E \rightarrow F(c)$ (where $E \in S$), $(\xi_{(c,x)} \circ u, \xi_{(c,y)} \circ v) \in R_E$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ an object $a_i \in \mathcal{C}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and an arrow $f_i : c \rightarrow a_i$ in $\mathcal{C}$ such that $P(f_i)(x) = P(f_i)(y)$ and $F(f_i) \circ h_i = u \circ e_i$.

$\square$

3 Extensions of flat functors

In this section, we investigate the operation on flat functors induced by a geometric morphism of toposes. This will be relevant for the characterization of the class of theories classified by a presheaf topos addressed in section 5.

3.1 General extensions

Let $(\mathcal{C}, J)$ and $(\mathcal{D}, K)$ be Grothendieck sites, and $u : \mathbf{Sh}(\mathcal{D}, K) \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ a geometric morphism. This morphism induces, via Diaconescu's equivalence, a functor

$$\xi_{\mathcal{E}} : \mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) \rightarrow \mathbf{Flat}_J(\mathcal{C}, \mathcal{E}),$$

where we write $\mathbf{Flat}_Z(\mathcal{R}, \mathcal{E})$ for the category of flat Z -continuous functors from $\mathcal{R}$ to $\mathcal{E}$.

This functor can be described explicitly as follows. For any flat K -continuous functor $F : \mathcal{D} \rightarrow \mathcal{E}$ with corresponding geometric morphism $f_F : \mathcal{E} \rightarrow \mathbf{Sh}(\mathcal{D}, K)$, the flat functor $\tilde{F} := \xi_{\mathcal{E}}(F) : \mathcal{C} \rightarrow \mathcal{E}$ is given by $f_F^* \circ u^* \circ l_{\mathcal{C}}$, where $l_{\mathcal{C}} : \mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ is the composite of the Yoneda embedding $y_{\mathcal{C}} : \mathcal{C} \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ with the associated sheaf functor $[\mathcal{C}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}, J)$.

For any natural transformation $\alpha : F \rightarrow G$ between flat functors in $\mathbf{Flat}_K(\mathcal{D}, \mathcal{E})$, the corresponding natural transformation $f_{\alpha}^* : f_F^* \rightarrow f_G^*$, applied to the functor $u^* \circ l_{\mathcal{C}}$, gives rise to a natural transformation $\tilde{\alpha} : \tilde{F} \rightarrow \tilde{G}$ which is precisely $\xi_{\mathcal{E}}(\alpha)$.

We can express $\tilde{F}$ directly in terms of F by using a colimit construction, as follows.

For any $c \in \mathcal{C}$, the K -sheaf $u^*(l_{\mathcal{C}}(c)) : \mathcal{D}^{\text{op}} \rightarrow \mathbf{Set}$, considered as an object of $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, can be canonically expressed as a colimit of representables, indexed over its category of elements $\mathcal{A}_c$; specifically, $u^*(l_{\mathcal{C}}(c)) \cong \text{colim}(y_{\mathcal{D}} \circ \pi_c)$ in $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, where $\pi_c : \mathcal{A}_c \rightarrow \mathcal{D}$ is the canonical projection functor. As the associated sheaf functor $a_K : [\mathcal{D}^{\text{op}}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{D}, K)$ preserves colimits, the functor $u^*(l_{\mathcal{C}}(c))$ is isomorphic in $\mathbf{Sh}(\mathcal{D}, K)$ to the colimit of the functor $l_{\mathcal{D}} \circ \pi_c$. Therefore $\tilde{F}(c) = (f_F^* \circ u^* \circ l_{\mathcal{C}})(c) = f_F^*(\text{colim}(l_{\mathcal{D}} \circ \pi_c)) \cong \text{colim}(f_F^* \circ l_{\mathcal{D}} \circ \pi_c) = \text{colim}(F \circ \pi_c)$.

For any $(d, z) \in \mathcal{A}_c$, we write $\kappa_{(d, z)}^F : F(d) \rightarrow \tilde{F}(c)$ for the canonical colimit arrow.

Proposition 3.1. (i) *For any natural transformation $\alpha : F \rightarrow G$ between flat functors $F, G : \mathcal{D} \rightarrow \mathcal{E}$ in $\mathbf{Flat}_K(\mathcal{D}, \mathcal{E})$, the natural transformation $\tilde{\alpha} : \tilde{F} \rightarrow \tilde{G}$ is characterized by the following condition: for any object (d, z) of the category $\mathcal{A}_c$ the diagram*

$$\begin{array}{ccc} \tilde{F}(c) & \xrightarrow{\tilde{\alpha}(c)} & \tilde{G}(c) \\ \kappa_{(d, z)}^F \uparrow & & \uparrow \kappa_{(d, z)}^G \\ F(d) & \xrightarrow{\alpha(d)} & G(d) \end{array}$$

commutes.

(ii) *The functor $\xi_{\mathcal{E}} : \mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) \rightarrow \mathbf{Flat}_J(\mathcal{C}, \mathcal{E})$ is natural in $\mathcal{E}$; that is,*

for any geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$, the diagram

$$\begin{array}{ccc} \mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) & \xrightarrow{\xi_{\mathcal{E}}} & \mathbf{Flat}_J(\mathcal{C}, \mathcal{E}) \\ \downarrow f^* \circ - & & \downarrow f^* \circ - \\ \mathbf{Flat}(\mathcal{D}, \mathcal{F}) & \xrightarrow{\xi_{\mathcal{F}}} & \mathbf{Flat}(\mathcal{C}, \mathcal{F}) \end{array}$$

commutes.

(iii) For any arrow $f : c \rightarrow c'$ in $\mathcal{C}$ and any object (d, z) in $\mathcal{A}_c$, the diagram

$$\begin{array}{ccc} \tilde{F}(c) & \xrightarrow{\tilde{F}(f)} & \tilde{F}(c') \\ \uparrow \kappa_{(d,z)}^F & \nearrow \kappa_{(d, u^*(l_{\mathcal{C}}(f))(z))}^F & \\ F(d) & & \end{array}$$

commutes.

Proof (i) The given square commutes as it is the naturality square for the natural transformation $f_{\alpha}^* : f_F^* \rightarrow f_G^*$ with respect to the canonical colimit arrow $y_{\mathcal{D}}(d) \rightarrow u^*(l_{\mathcal{C}}(c))$ in $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$ corresponding to the object (d, z) of $\mathcal{A}_c$.

(ii) This follows as an immediate consequence of the fact that Diaconescu's equivalences are natural in $\mathcal{E}$.

(iii) Given a natural transformation $\beta : P \rightarrow P'$ of presheaves P and P' in $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, denoting by $\kappa^P : \int P \rightarrow \mathcal{D}$ and $\kappa^{P'} : \int P' \rightarrow \mathcal{D}$ the canonical projection functors and by $\kappa_{(d,z)}^P : y_{\mathcal{D}}d \rightarrow P$, $\kappa_{(d',z')}^{P'} : y_{\mathcal{D}}d' \rightarrow P$ the canonical colimit arrows in $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$ (for $(d, z) \in \int P$ and $(d', z') \in \int P'$), we clearly have that $\beta \circ \kappa_{(d,z)}^P = k_{(d, \beta(d)(z))}^{P'}$. To obtain our thesis it suffices to apply this result to the natural transformation $\beta : u^*(l_{\mathcal{C}}(f)) : u^*(l_{\mathcal{C}}(c)) \rightarrow u^*(l_{\mathcal{C}}(c'))$; indeed, by applying the functor f_F^* to the resulting equality we obtain precisely the identity in the statement of the proposition. $\square$

3.2 Extensions along embeddings of categories

Let $\mathcal{D}$ be a subcategory of a small category $\mathcal{C}$. Then the inclusion functor $i : \mathcal{D} \hookrightarrow \mathcal{C}$ induces a geometric morphism $E(i) : [\mathcal{D}, \mathbf{Set}] \rightarrow [\mathcal{C}, \mathbf{Set}]$ and hence, by Diaconescu's equivalence, a functor

$$\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{D}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E}) .$$

We can describe $\tilde{F}$ directly in terms of F as follows. For any $c \in \mathcal{C}$, the functor $E(i)^*(y_{\mathcal{C}}(c)) : \mathcal{D} \rightarrow \mathbf{Set}$ coincides with the functor $\text{Hom}_{\mathcal{C}}(c, -) : \mathcal{D} \rightarrow \mathbf{Set}$. Its category of elements $\mathcal{A}_c$ has as objects the pairs (d, h) where

d is an object of $\mathcal{D}$ and h is an arrow $h : c \rightarrow d$ in $\mathcal{C}$ and as arrows $(d, h) \rightarrow (d', h')$ the arrows $k : d' \rightarrow d$ in $\mathcal{D}$ such that $k \circ h' = h$ in $\mathcal{C}$. As f_F^* preserves all small colimits and $\text{Hom}_{\mathcal{C}}(c, -)$ is the colimit in $[\mathcal{D}, \mathbf{Set}]$ of the composition of the canonical projection $\pi_c : \mathcal{A}_c \rightarrow \mathcal{D}^{\text{op}}$ with the Yoneda embedding $\mathcal{D}^{\text{op}} \rightarrow [\mathcal{D}, \mathbf{Set}]$, the object $\tilde{F}(c)$ can be identified with the colimit of the composite functor $F \circ \pi_c : \mathcal{A}_c \rightarrow \mathcal{E}$. For any object d of $\mathcal{D}$, the pair $(d, 1_d)$ defines an object of the category $\mathcal{A}_d$ and hence a canonical colimit arrow $\chi_d : F(d) \rightarrow \tilde{F}(d)$, also denoted by χ_d^F . For any object (d, h) of $\mathcal{A}_c$, the colimit arrow $\kappa_{(d, h)} : F(d) \rightarrow \tilde{F}(c)$ is equal to the composite $\tilde{F}(h) \circ \chi_d$.

Remarks 3.2. (i) The functor $\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{D}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E})$ coincides with the (restriction to the subcategories of flat functors) of the left Kan extension functor $[\mathcal{D}^{\text{op}}, \mathcal{E}] \rightarrow [\mathcal{C}^{\text{op}}, \mathcal{E}]$ along the inclusion $\mathcal{D}^{\text{op}} \hookrightarrow \mathcal{C}^{\text{op}}$ (cf. Remark 3.13 below).

(ii) For any object $d \in \mathcal{D}$ the diagram

$$\begin{array}{ccc} \tilde{F}(d) & \xrightarrow{\tilde{\alpha}(d)} & \tilde{G}(d) \\ \chi_d^F \uparrow & & \uparrow \chi_d^G \\ F(d) & \xrightarrow{\alpha(d)} & G(d) \end{array}$$

commutes (cf. Proposition 3.1(a)).

(iii) For any object $d \in \mathcal{D}$, the arrow $\chi_d : F(d) \rightarrow \tilde{F}(d)$ is monic. Indeed, it can be identified with the image of the canonical subfunctor inclusion $\text{Hom}_{\mathcal{D}}(d, -) \hookrightarrow \text{Hom}_{\mathcal{C}}(d, -)$ under the inverse image f_F^* of the geometric morphism f_F . From this, in view of Remark 3.2(a), it follows at once that the functor $\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{D}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E})$ is faithful.

(iv) For any flat functors $F, G : \mathcal{D}^{\text{op}} \rightarrow \mathcal{E}$ and natural transformation $\beta : \tilde{F} \rightarrow \tilde{G}$, $\beta = \tilde{\alpha}$ for some natural transformation $\alpha : F \rightarrow G$ if and only if for every $d \in \mathcal{D}$ the diagram

$$\begin{array}{ccc} \tilde{F}(d) & \xrightarrow{\beta(d)} & \tilde{G}(d) \\ \chi_d^F \uparrow & & \uparrow \chi_d^G \\ F(d) & \xrightarrow{\alpha(d)} & G(d) \end{array}$$

commutes. One direction follows from Remark 3.2(a). To prove the other one we observe that $\beta(c) = \tilde{\alpha}(c)$ for all $c \in \mathcal{C}$ if and only if for every object $(a, z) \in \mathcal{A}_c$, $\beta(c) \circ \kappa_{(a, z)} = \tilde{\alpha}(c) \circ \kappa_{(a, z)}$. Now, $\beta(c) \circ \kappa_{(a, z)} = \beta(c) \circ \tilde{F}(z) \circ \chi_a^F = \tilde{G}(z) \circ \beta(a) \circ \chi_a^F = \tilde{G}(z) \circ \chi_a^G \circ \alpha(a) = \kappa_{(a, z)}^G \circ \alpha(a) = \tilde{\alpha}(c) \circ \kappa_{(a, z)}^F$ (where the third equality follows from the naturality of β and the last from Proposition 3.1(a)), as required.

- (v) Let d be an object of $\mathcal{D}$ and $y_{\mathcal{D}}d = \text{Hom}_{\mathcal{D}}(-, d) : \mathcal{D}^{\text{op}} \rightarrow \mathbf{Set}$ the representable on $\mathcal{D}$ associated to d . Then $\xi_{\mathcal{E}}(y_{\mathcal{D}}d) = y_{\mathcal{C}}d$, where $y_{\mathcal{C}}d = \text{Hom}_{\mathcal{C}}(-, d)$ is the representable on $\mathcal{C}$ associated to d (considered here as an object of $\mathcal{C}$). Indeed, the flat functor $y_{\mathcal{D}}d$ corresponds to the geometric morphism whose inverse image is the evaluation functor $ev_d^{\mathcal{D}} : [\mathcal{D}, \mathbf{Set}] \rightarrow \mathbf{Set}$ at the object d , and the composite $ev_d^{\mathcal{D}} \circ E(i)^*$ coincides with the evaluation functor $ev_d^{\mathcal{C}} : [\mathcal{C}, \mathbf{Set}] \rightarrow \mathbf{Set}$ at the object d , which corresponds to the flat functor $y_{\mathcal{C}}d$.
- (vi) Let $\mathcal{E}$ be a Grothendieck topos and $\gamma_{\mathcal{E}} : \mathcal{E} \rightarrow \mathbf{Set}$ the unique geometric morphism from $\mathcal{E}$ to $\mathbf{Set}$. Then by Proposition 3.1(ii) and Remark 3.2(v), for any object d of $\mathcal{D}$, the functor $\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{D}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E})$ sends the flat functor $\gamma_{\mathcal{E}}^* \circ y_{\mathcal{D}}d$ to the functor $\gamma_{\mathcal{E}}^* \circ y_{\mathcal{C}}d$.

Proposition 3.3. *Under the natural equivalences $e_{\mathcal{C}} : \text{Ind-}\mathcal{C} \simeq \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathbf{Set})$ and $e_{\mathcal{D}} : \text{Ind-}\mathcal{D} \simeq \mathbf{Flat}(\mathcal{D}^{\text{op}}, \mathbf{Set})$, the functor $\xi_{\mathbf{Set}}$ corresponds to the functor $\text{Ind-}i : \text{Ind-}\mathcal{D} \rightarrow \text{Ind-}\mathcal{C}$.*

Proof Let us first recall the definition of the functor $\text{Ind-}i : \text{Ind-}\mathcal{D} \rightarrow \text{Ind-}\mathcal{C}$. For any flat functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$, $\text{Ind-}i(F)$ is the functor given by the colimit in $\text{Ind-}\mathcal{C}$ of the composite functor $i \circ \pi_F$, where $\pi_F : \int F \rightarrow \mathcal{D}$ is the canonical projection from the category of elements $\int F$ of F to $\mathcal{D}$. For any $c \in \mathcal{C}$, we thus have $\text{Ind-}i(F)(c) = \text{colim}(ev_c \circ i \circ \pi_F)$, where $ev_c : \text{Ind-}\mathcal{C} \rightarrow \mathbf{Set}$ is the evaluation functor at c .

Now, the functor $ev_c \circ i : \mathcal{D} \rightarrow \mathbf{Set}$ is equal to the functor $\text{Hom}_{\mathcal{C}}(c, -) : \mathcal{D} \rightarrow \mathbf{Set}$ considered above, and $\tilde{F}(c) = \text{colim}(F \circ \pi_c) : \mathcal{A}_c \rightarrow \mathbf{Set}$, where $\mathcal{A}_c$ is the category of elements of $ev_c \circ i$ and $\pi_c : \mathcal{A}_c \rightarrow \mathcal{D}^{\text{op}}$ the associated canonical projection. The commutativity of the tensor product between a presheaf and a covariant set-valued functor (cf. chapter VII of [26] or Theorem 2.5 above) thus yields a natural isomorphism between the two sets, as required. $\square$

Corollary 3.4. *Let $\mathcal{D} \hookrightarrow \mathcal{C}$ be an embedding of small categories and $F : \mathcal{D}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. With the above notation, for any object $c \in \mathcal{C}$, we have*

$$\tilde{F}(c) \cong \coprod_{(a,z) \in \mathcal{A}_c} F(c)/R_c,$$

where R_c is the equivalence relation in $\mathcal{E}$ defined by saying that, for any objects $(a, z), (a', z')$ of the category $\mathcal{A}_c$, the geometric bi-sequent

$$\begin{aligned} & (\ulcorner R \urcorner (\ulcorner \xi_{(a,z)} \urcorner (x), \ulcorner \xi_{(a',z')} \urcorner (x')) \dashv\vdash_{x^{F(a)}, x'^{F(a')}} \\ & \bigvee_{\substack{a \xrightarrow{f} b \xrightarrow{g} a' \\ f \circ z = g \circ z'}} (\exists y^{F(b)}) (\ulcorner F(f) \urcorner (y) = x \wedge \ulcorner F(g) \urcorner (y) = x')) \end{aligned}$$

is valid in the tautological $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$ (defined in section 2.7), where for any object (a, z) of the category $\mathcal{A}_c$, $\xi_{(a,z)} : F(a) \rightarrow \tilde{F}(c)$ is the canonical colimit arrow.

In particular, for any objects $(a, z), (a', z')$ of the category $\mathcal{A}_c$, the geometric bi-sequent

$$(\ulcorner R \urcorner (\ulcorner \xi_{(a,z)} \urcorner (x), \ulcorner \xi_{(a',z')} \urcorner (x)) \dashv\vdash_{x^{F(a)}} \bigvee_{\substack{a \xrightarrow{f} b \\ f \circ z = f \circ z'}} (\exists y^{F(b)}) (\ulcorner F(f) \urcorner (y) = x))$$

is valid in the $\Sigma_{\mathcal{E}}$ -structure $\mathcal{S}_{\mathcal{E}}$.

Semantically, the relation R_c can be characterized by saying that, for any objects $(a, z), (a', z')$ of the category $\mathcal{A}_c$ and any generalized elements $x : E \rightarrow F(a), x' : E \rightarrow F(a'), (\kappa_{(a,z)} \circ x, \kappa_{(a',z')} \circ x') \in R_c$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ and for each index $i \in I$ an object $b_i \in \mathcal{D}$, a generalized element $h_i : E_i \rightarrow F(b_i)$ and two arrows $f_i : a \rightarrow b_i$ and $f'_i : a' \rightarrow b_i$ in $\mathcal{D}$ such that $f'_i \circ z' = f_i \circ z$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$.

In particular, for any objects (a, z) and (a', z') of $\mathcal{A}_c$ and any generalized element $x : E \rightarrow F(a), (\kappa_{(a,z)} \circ x, \kappa_{(a',z')} \circ x) \in R_c$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ and for each index $i \in I$ an object $b_i \in \mathcal{D}$, a generalized element $h_i : E_i \rightarrow F(b_i)$ and an arrow $f_i : a \rightarrow b_i$ in $\mathcal{D}$ such that $f_i \circ z' = f_i \circ z$ and $F(f_i) \circ h_i = x \circ e_i$.

Proof The corollary can be obtained by applying Proposition 2.34 (and its categorical reformulation provided by Proposition 2.35) to the flat functor $F : \mathcal{D}^{\text{op}} \rightarrow \mathcal{E}$ and the functor $P : \mathcal{D} \rightarrow \mathbf{Set}$ given by $ev_c \circ i = \text{Hom}_{\mathcal{C}}(c, -) : \mathcal{D} \rightarrow \mathbf{Set}$, whose category of elements coincides with $\mathcal{A}_c$ and whose associated fibration π_P coincides with $\pi_c : \mathcal{A}_c \rightarrow \mathcal{D}^{\text{op}}$. $\square$

3.3 Extensions from categories of set-based models to syntactic categories

Let $\mathbb{T}$ be a geometric theory and $\mathcal{K}$ a small category of set-based $\mathbb{T}$ -models. Then the family of geometric morphisms $\mathbf{Set} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ corresponding to the $\mathbb{T}$ -models in $\mathcal{K}$ induces a geometric morphism

$$p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

whose associated $\mathbb{T}$ -model in $[\mathcal{K}, \mathbf{Set}]$ is given, at each sort, by the corresponding forgetful functor.

We thus have, for each Grothendieck topos $\mathcal{E}$, an induced functor

$$u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E}),$$

which the following theorem describes explicitly.

Before stating it, we need to introduce some notation. For any any object $\{\vec{x} . \phi\}$ of $\mathcal{C}_{\mathbb{T}}$, we write $\mathcal{A}_{\{\vec{x} . \phi\}}^{\mathcal{K}}$ or simply $\mathcal{A}_{\{\vec{x} . \phi\}}$ when the category $\mathcal{K}$ can be obviously inferred from the context, for the category whose objects are the pairs (M, w) , where $M \in \mathcal{K}$ and $w \in [[\vec{x} . \phi]]_M$ and whose arrows $(M, w) \rightarrow (N, z)$ are the $\mathbb{T}$ -model homomorphisms $g : N \rightarrow M$ in $\mathcal{K}$ such that $g(w) = z$. We denote by $\pi_{\{\vec{x} . \phi\}}^{\mathcal{K}} : \mathcal{A}_{\{\vec{x} . \phi\}}^{\mathcal{K}} \rightarrow \mathcal{K}^{\text{op}}$ the canonical projection functor.

Theorem 3.5. *Let $\mathbb{T}$ be a geometric theory. Then for any flat functor $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$, the functor $\tilde{F} := u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F) : \mathcal{C}_{\mathbb{T}} \rightarrow \mathcal{E}$ sends any object $\{\vec{x} . \phi\}$ of $\mathcal{C}_{\mathbb{T}}$ to the colimit $\text{colim}(F \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}})$ and acts on the arrows in the obvious way. In particular, for any formula $\{\vec{x} . \phi\}$ presenting a $\mathbb{T}$ -model $M_{\{\vec{x} . \phi\}}$ in $\mathcal{K}$, $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{x} . \phi\}) \cong F(M_{\{\vec{x} . \phi\}})$.*

Proof Let $g_F : \mathcal{E} \rightarrow [\mathcal{K}, \mathbf{Set}]$ be the geometric morphism corresponding, via Diaconescu's equivalence, to the flat functor F . Then the functor $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)$ is equal to the composite $g_F^* \circ p_{\mathcal{K}}^* \circ y$, where $y : \mathcal{C}_{\mathbb{T}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ is the Yoneda embedding.

Now, for any geometric formula $\{\vec{x} . \phi\}$ over Σ , $p_{\mathcal{K}}^*(y(\{\vec{x} . \phi\}))$ is the functor $F_{\{\vec{x} . \phi\}}$ sending to any model $M \in \mathcal{K}$ the set $[[\vec{x} . \phi]]_M$. This functor can be expressed as the colimit $\text{colim}(y' \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}})$, where $y' : \mathcal{K}^{\text{op}} \rightarrow [\mathcal{K}, \mathbf{Set}]$ is the Yoneda embedding, since the functor $\pi_{\{\vec{x} . \phi\}}^{\mathcal{K}}$ coincides with the canonical projection from the category of elements of the functor $F_{\{\vec{x} . \phi\}}$ to $\mathcal{K}^{\text{op}}$.

Therefore $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{x} . \phi\}) = g_F^*(\text{colim}(y' \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}})) \cong \text{colim}(g_F^* \circ y' \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}}) \cong \text{colim}(F \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}})$, as required. $\square$

Remark 3.6. From the proof of the theorem, it is clear that the isomorphism $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{x} . \phi\}) \cong F(M_{\{\vec{x} . \phi\}})$ is natural in $\{\vec{x} . \phi\}$; that is, for any geometric formulae $\{\vec{x} . \phi\}$ and $\{\vec{y} . \psi\}$ respectively presenting $\mathbb{T}$ -models $M_{\{\vec{x} . \phi\}}$ and $M_{\{\vec{y} . \psi\}}$ and any $\mathbb{T}$ -provably functional formula $\theta : \{\vec{x} . \phi\} \rightarrow \{\vec{y} . \psi\}$ inducing a $\mathbb{T}$ -model homomorphism $M_{\theta} : M_{\{\vec{y} . \psi\}} \rightarrow M_{\{\vec{x} . \phi\}}$, the arrow $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\theta)$ is given by the image of the arrow $F(M_{\theta})$ across the isomorphisms $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{x} . \phi\}) \cong F(M_{\{\vec{x} . \phi\}})$ and $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{y} . \psi\}) \cong F(M_{\{\vec{y} . \psi\}})$.

Corollary 3.7. *Let $\mathbb{T}$ be a geometric theory, $\sigma = (\phi \vdash_{\vec{x}} \psi)$ a geometric sequent over the signature of $\mathbb{T}$ and $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. If σ is valid in every $\mathbb{T}$ -model in $\mathcal{K}$ then $\tilde{F}(\{\vec{x} . \phi\}) \leq \tilde{F}(\{\vec{x} . \psi\})$ as subobjects of $\tilde{F}(\{\vec{x} . \top\})$ in $\mathcal{E}$.*

Proof By Theorem 3.5, we have that $\tilde{F}(\{\vec{x} . \phi\}) = \text{colim}(F \circ \pi_{\{\vec{x} . \phi\}}^{\mathcal{K}})$, $\tilde{F}(\{\vec{x} . \psi\}) = \text{colim}(F \circ \pi_{\{\vec{x} . \psi\}}^{\mathcal{K}})$ and $\tilde{F}(\{\vec{x} . \top\}) = \text{colim}(F \circ \pi_{\{\vec{x} . \top\}}^{\mathcal{K}})$. Now, $\mathcal{A}_{\{\vec{x} . \phi\}}$ and $\mathcal{A}_{\{\vec{x} . \psi\}}$ canonically embed as subcategories of $\mathcal{A}_{\{\vec{x} . \top\}}$, and if σ is

valid in every $\mathbb{T}$ -model in $\mathcal{K}$ then we have a canonical functor $i : \mathcal{A}_{\{\vec{x}, \phi\}} \rightarrow \mathcal{A}_{\{\vec{x}, \psi\}}$ which commutes with these embeddings. It thus follows from the functoriality of colimits that $\tilde{F}(\{\vec{x} \cdot \phi\}) \leq \tilde{F}(\{\vec{x} \cdot \psi\})$ as subobjects of $\tilde{F}(\{\vec{x} \cdot \top\})$ in $\mathcal{E}$, as required. $\square$

Let us now apply Proposition 2.35 in the context of extensions $F \rightarrow \tilde{F}$ of flat functors induced by the geometric morphism

$$p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}}).$$

The following characterization is obtained by applying it in conjunction with Proposition 4.5.

Proposition 3.8. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small category of set-based $\mathbb{T}$ -models, $\mathcal{E}$ a Grothendieck topos with a separating set S and $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. With the above notation, for any geometric formula-in-context $\phi(\vec{x})$ over Σ , we have*

$$\tilde{F}(\{\vec{x} \cdot \phi\}) \cong \left(\coprod_{(M, z) \in \mathcal{A}_{\{\vec{x}, \phi\}}} F(M) \right) / R_{\{\vec{x}, \phi\}},$$

where $R_{\{\vec{x}, \phi\}}$ is the equivalence relation in $\mathcal{E}$ defined by saying that for any objects $(M, z), (N, w)$ of the category $\mathcal{A}_{\{\vec{x}, \phi\}}$ and any generalized elements $x : E \rightarrow F(M)$, $x' : E \rightarrow F(N)$ (where $E \in S$), we have $(\xi_{(M, z)} \circ x, \xi_{(N, w)} \circ x') \in R_{\{\vec{x}, \phi\}}$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and two $\mathbb{T}$ -model homomorphisms $f_i : M \rightarrow a_i$ and $f'_i : N \rightarrow a_i$ such that $f_i(z) = f'_i(w)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$ (where for any object (M, z) of the category $\mathcal{A}_{\{\vec{x}, \phi\}}$, $\kappa_{(M, z)} : F(M) \rightarrow \tilde{F}(\{\vec{x} \cdot \phi\})$ is the canonical colimit arrow).

In particular, for any objects (M, z) and (M, w) of $\mathcal{A}_{\{\vec{x}, \phi\}}$ and any generalized element $x : E \rightarrow F(M)$ (where $E \in S$), $(\kappa_{(M, z)} \circ x, \kappa_{(M, w)} \circ x) \in R_{\{\vec{x}, \phi\}}$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and a $\mathbb{T}$ -model homomorphism $f_i : M \rightarrow a_i$ such that $f_i(z) = f_i(w)$ and $F(f_i) \circ h_i = x \circ e_i$. $\square$

Let $M_{\mathbb{T}}$ the universal model of $\mathbb{T}$ in the syntactic category $\mathcal{C}_{\mathbb{T}}$. We can represent the model $\tilde{F}(M_{\mathbb{T}})$ as a $\mathcal{E}$ -indexed filtered colimit of set-based models of $\mathbb{T}$. For simplicity, we shall first establish this representation in the case of the topos of sets, and then generalize it to an arbitrary topos.

Let $F : \mathcal{K}^{\text{op}} \rightarrow \mathbf{Set}$ be a flat functor. By Theorem 3.5, for any sort A over Σ , $\tilde{F}(M_{\mathbb{T}})A = \tilde{F}(\{x^A \cdot \top\}) = \text{colim}(F \circ \pi_{\{x^A, \top\}})$, where $\pi_{\{x^A, \top\}}$ is the canonical projection functor $\mathcal{A}_{\{x^A, \top\}} \rightarrow \mathcal{K}^{\text{op}}$ to $\mathcal{K}^{\text{op}}$ from the category

of elements $\mathcal{A}_{\{x^A, \top\}}$ of the functor $P_A : \mathcal{K} \rightarrow \mathbf{Set}$ which assigns any model M in $\mathcal{K}$ to the set MA and acts accordingly on the arrows. Now, it follows from the commutativity of the tensor product between a presheaf and a covariant set-valued functor (cf. chapter VII of [26] or section 2.5 above) that the colimit $\text{colim}(F \circ \pi_{\{x^A, \top\}})$ is isomorphic to the colimit $\text{colim}(P_A \circ \pi_F)$, where $\pi_F : \int F \rightarrow \mathcal{K}$ is the canonical projection functor from the category of elements of the functor F to $\mathcal{K}$. Since the functor F is flat, the category $\int F$ is filtered. Therefore, as filtered colimits in $\mathbb{T}\text{-mod}(\mathbf{Set})$ are computed sortwise as in $\mathbf{Set}$, $\tilde{F}(M_{\mathbb{T}}) \cong \text{colim}(i \circ \pi_F)$, where $i : \mathcal{K} \hookrightarrow \mathbb{T}\text{-mod}(\mathbf{Set})$ is the canonical inclusion. So for any object (c, x) of the category $\int F$ we have a $\mathbb{T}$ -model homomorphism $\xi_{(c,x)} : c \rightarrow \tilde{F}(M_{\mathbb{T}})$, which can be expressed in terms of the colimit arrows $\kappa_{(a,y)} : F(a) \rightarrow \tilde{F}(\{x^A, \top\}) = \tilde{F}(M_{\mathbb{T}})A$ (for $y \in cA$ and A sort over Σ) as follows: for any sort A over Σ , $\xi_{(c,x)}A(y) = \kappa_{(c,y)}(x)$. The explicit description of filtered colimits in the category $\mathbf{Set}$ thus yields, for each sort A over Σ , the following characterizing properties of the colimit $\text{colim}(P_A \circ \pi_F)$ in terms of the arrows $\xi_{(c,x)}$:

- (i) For any element x of $\tilde{F}(M_{\mathbb{T}})A$ there exists an object (c, x) of the category $\int F$ and an element y of cA such that $\xi_{(c,x)}A(y) = x$;
- (ii) For any objects (c, x) and (c', x') of the category $\int F$ and elements $y \in cA$ and $y' \in c'A$, we have $\xi_{(c,x)}A(y) = \xi_{(c',x')}A(y')$ if and only if there exists an object (c'', x'') of $\int F$ and arrows $f : c \rightarrow c''$ and $g : c' \rightarrow c''$ in $\mathcal{K}$ such that $F(f)(x'') = x$, $F(g)(x'') = x'$ and $fA(y) = gA(y')$.

Moreover, the filteredness of the category $\int F$ implies the following ‘joint embedding property’: for any objects (c, x) and (c', x') of the category $\int F$ there exists an object (c'', x'') of $\int F$ and arrows $f : c \rightarrow c''$ and $g : c' \rightarrow c''$ in $\mathcal{K}$ such that $F(f)(x'') = x$, $F(g)(x'') = x'$ and $\xi_{(c,x)} \circ f = \xi_{(c',x')} \circ g$.

Let us now proceed to establish the $\mathcal{E}$ -indexed generalization of this result.

Suppose that $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ is a flat functor, where $\mathcal{K}$ is a small subcategory of the category $\mathbb{T}\text{-mod}(\mathbf{Set})$. By Theorem 3.5, for any sort A over Σ , we have that $\tilde{F}(M_{\mathbb{T}})A = \tilde{F}(\{x^A, \top\}) = \text{colim}(F \circ \pi_{\{x^A, \top\}})$; in particular, for any object (a, y) of the category $\mathcal{A}_{\{x^A, \top\}}$ we have a colimit arrow $\kappa_{(a,y)} : F(a) \rightarrow \tilde{F}(\{x^A, \top\}) = \tilde{F}(M_{\mathbb{T}})A$ in $\mathcal{E}$. More generally, for any formula-in-context $\phi(\vec{x})$ over Σ , any model c in $\mathcal{K}$ and any element $\vec{y}$ of $[[\vec{x} \cdot \phi]]_c$, we have a colimit arrow $\kappa_{(c,\vec{y})} : F(c) \rightarrow \tilde{F}(\{\vec{x} \cdot \phi\})$.

Proposition 3.9. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small subcategory of the category $\mathbb{T}\text{-mod}(\mathbf{Set})$, $\mathcal{E}$ a Grothendieck topos and $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. With the above notation, for any pair (c, x) consisting of an object c of $\mathcal{K}$ and of a generalized element $x : E \rightarrow F(c)$, there is a Σ -structure homomorphism $\xi_{(c,x)} : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ defined as*

follows: for any sort A over Σ , the function $\xi_{(c,x)} : cA \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^A . \top\}))$ sends any element $y \in cA$ to the generalized element $\kappa_{(c,y)} \circ x : E \rightarrow \tilde{F}(\{x^A . \top\})$.

Proof We have to verify that:

- (1) For any function symbol $f : A_1, \dots, A_n \rightarrow B$ over Σ , the diagram

$$\begin{array}{ccc}
 cA_1 \times \dots \times cA_n & \xrightarrow{cf} & cB \\
 \downarrow \xi_{(c,x)} A_1 \times \dots \times \xi_{(c,x)} A_n & & \downarrow \xi_{(c,x)} B \\
 \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^{A_1} . \top\}) \times \dots \times \tilde{F}(\{x^{A_n} . \top\})) & & \\
 \downarrow i & & \\
 \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^{A_1}, \dots, x^{A_n} . \top\})) & \xrightarrow{\text{Hom}_{\mathcal{E}}(E, \tilde{F}([f]))} & \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^B . \top\})),
 \end{array}$$

where i is the canonical isomorphism $\tilde{F}(\{x^{A_1} . \top\}) \times \dots \times \tilde{F}(\{x^{A_n} . \top\}) \cong \tilde{F}(\{x^{A_1}, \dots, x^{A_n} . \top\})$ induced by the preservation of finite products by $\tilde{F}$, commutes;

- (2) For any relation symbol $R \mapsto A_1, \dots, A_n$ over Σ , we have a commutative diagram

$$\begin{array}{ccc}
 cR & \xrightarrow{\quad} & \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^{A_1}, \dots, x^{A_n} . R\})) \\
 \downarrow & & \downarrow \\
 cA_1 \times \dots \times cA_n & \xrightarrow{\quad} & \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^{A_1}, \dots, x^{A_n} . \top\})),
 \end{array}$$

where the unnamed vertical arrows are the canonical ones and the lower horizontal arrow is $\text{Hom}_{\mathcal{E}}(E, i) \circ \xi_{(c,x)} A_1 \times \dots \times \xi_{(c,x)} A_n$.

To prove (1), we first observe that for any n -tuple $\vec{y} = (y_1, \dots, y_n) \in cA_1 \times \dots \times cA_n$, $i \circ \langle \kappa_{(c,y_1)}, \dots, \kappa_{(c,y_n)} \rangle = \kappa_{(c,\vec{y})}$. Indeed, for any $i \in \{1, \dots, n\}$, the arrow $\tilde{F}(\pi_i) \circ i$, where $\pi_i : \{x^{A_1}, \dots, x^{A_n} . \top\} \rightarrow \{x^{A_i} . \top\}$ is the canonical projection arrow in the syntactic category $\mathcal{C}_{\mathbb{T}}$, is equal to the i -th product projection $\tilde{F}(\{x^{A_1} . \top\}) \times \dots \times \tilde{F}(\{x^{A_n} . \top\}) \rightarrow \tilde{F}(\{x^{A_i} . \top\})$, by Proposition 3.1(iii). Therefore, to prove the required condition it is equivalent to verify that for any n -tuple $\vec{y} = (y_1, \dots, y_n) \in cA_1 \times \dots \times cA_n$, $\xi_{(c,x)} B(cf(\vec{y})) = \tilde{F}([f]) \circ \kappa_{(c,\vec{y})} \circ x$, where $[f] : \{x^{A_1}, \dots, x^{A_n} . \top\} \rightarrow \{x^B . \top\}$ is the arrow in the syntactic category $\mathcal{C}_{\mathbb{T}}$ corresponding to the function symbol f . But $\xi_{(c,x)} B(cf(\vec{y})) = \kappa_{(c,cf(\vec{y}))} \circ x$, and we have that $\tilde{F}([f]) \circ \kappa_{(c,\vec{y})} = \kappa_{(c,cf(\vec{y}))}$, again by Proposition 3.1(iii).

To prove (2), it suffices to observe that, by Proposition 3.1(iii), for any n -tuple $\vec{y} = (y_1, \dots, y_n) \in cA_1 \times \dots \times cA_n$ in cR , $\kappa_{(c,\vec{y})}$ factors through the canonical subobject $\tilde{F}(R) \mapsto \tilde{F}(\{x^{A_1}, \dots, x^{A_n} . \top\})$.

□

The following lemma describes some basic properties of the homomorphisms $\xi_{(c,x)}$.

Lemma 3.10. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small subcategory of the category $\mathbb{T}\text{-mod}(\mathbf{Set})$ and $F : \mathcal{K}^{op} \rightarrow \mathcal{E}$ a flat functor with values in a Grothendieck topos $\mathcal{E}$. Then*

- (i) *For any generalized element $x : E \rightarrow F(c)$ and any arrow $f : d \rightarrow c$ in $\mathcal{K}$, $\xi_{(c,x)} \circ f = \xi_{(c,F(f) \circ x)}$;*
- (ii) *For any generalized element $x : E \rightarrow F(c)$ and any arrow $e : E' \rightarrow E$, $\xi_{(c,x \circ e)} = \text{Hom}(e, \tilde{F}(M_{\mathbb{T}})) \circ \xi_{(c,x)}$*

Proof These properties can be easily proved by using the definition of the arrows $\xi_{(c,x)}$ in terms of the arrows $\kappa_{(a,y)}$ and Proposition 3.1(iii). $\square$

Proposition 3.11. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small subcategory of the category $\mathbb{T}\text{-mod}(\mathbf{Set})$ and $F : \mathcal{K}^{op} \rightarrow \mathcal{E}$ a flat functor with values in a Grothendieck topos $\mathcal{E}$. Let M be the $\mathbb{T}$ -model $\tilde{F}(M_{\mathbb{T}})$ in $\mathcal{E}$. Then, for any sort A over Σ , we have that*

- (i) *For any generalized element $x : E \rightarrow MA$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$, for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $x_i : E_i \rightarrow F(a_i)$ and an element $y_i \in a_i A$ such that $\xi_{(a_i, x_i)} A(y_i) = x \circ e_i$.*
- (ii) *For any pairs (a, x) and (b, x') , where a and b are $\mathbb{T}$ -models in $\mathcal{K}$ and $x : E \rightarrow F(a)$, $x' : E \rightarrow F(b)$ are generalized elements, and any elements $y \in aA$ and $y' \in bA$, we have that $\xi_{(a,x)} A(y) = \xi_{(b,x')} A(y')$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model c_i in $\mathcal{K}$, arrows $f_i : a \rightarrow c_i$, $g_i : b \rightarrow c_i$ and a generalized element $x_i : E_i \rightarrow F(c_i)$ such that $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$ for all $i \in I$ and $f_i A(y) = g_i A(y')$.*

Moreover, the following ‘joint embedding property’ holds: for any pairs (a, x) and (b, x') , where a and b are $\mathbb{T}$ -models in $\mathcal{K}$ and $x : E \rightarrow F(a)$, $x' : E \rightarrow F(b)$ are generalized elements, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model c_i in $\mathcal{K}$, arrows $f_i : a \rightarrow c_i$, $g_i : b \rightarrow c_i$ in $\mathcal{K}$ and a generalized element $x_i : E_i \rightarrow F(c_i)$ such that $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$ for all $i \in I$ and (by Lemma 3.10) the following diagram commutes:

$$\begin{array}{ccc}
a & \xrightarrow{\xi_{(a,x)}} & \text{Hom}_{\mathcal{E}}(E, M) \\
& \searrow f & \searrow \text{Hom}_{\mathcal{E}}(e_i, M) \\
& & c_i \xrightarrow{\xi_{(c_i, x_i)}} \text{Hom}_{\mathcal{E}}(E_i, M) \\
& \nearrow g & \nearrow \text{Hom}_{\mathcal{E}}(e_i, M) \\
b & \xrightarrow{\xi_{(b, x')}} & \text{Hom}_{\mathcal{E}}(E, M)
\end{array}$$

In fact, such family can be taken to be the pullback of the family of arrows $\langle F(f), F(g) \rangle : F(c) \rightarrow F(a) \times F(b)$ (for all spans $(f : a \rightarrow c, g : b \rightarrow c)$ in the category $\mathcal{K}$) along the arrow $\langle x, x' \rangle : E \rightarrow F(a) \times F(b)$.

Proof Recall from Theorem 3.5 that for any sort A over Σ , $\tilde{F}(M_{\mathbb{T}})A = \text{colim}(F \circ \pi_{\{x^A, \top\}}^{\mathcal{K}})$; now, $\pi_{\{x^A, \top\}}^{\mathcal{K}}$ is precisely the functor $P_A : \mathcal{K} \rightarrow \mathbf{Set}$ sending a model N in $\mathcal{C}$ to the set NA .

The proposition then straightforwardly follows from Theorem 2.26, applied to the functors $P_A : \mathcal{K} \rightarrow \mathbf{Set}$ (with A varying among the sorts of Σ), in view of Theorems 2.5 and 2.14. $\square$

3.4 A general adjunction

Let $\mathcal{C}$ be a small category and $\mathcal{E}$ a Grothendieck topos. Recall from [5] that the indexed category $[\mathcal{C}, \mathcal{E}]_{\mathcal{E}}$ is locally small, that is for any two functors $F, G : \mathcal{C} \rightarrow \mathcal{E}$ there exists an object $\text{Hom}_{[\mathcal{C}, \mathcal{E}]_{\mathcal{E}}}^{\mathcal{E}}(F, G)$ of $\mathcal{E}$ satisfying the universal property that for any object E of $\mathcal{E}$ the generalized elements $E \rightarrow \text{Hom}_{[\mathcal{C}, \mathcal{E}]_{\mathcal{E}}}^{\mathcal{E}}(F, G)$ correspond bijectively, naturally in $E \in \mathcal{C}$, to the natural transformations $!_E^* \circ F \rightarrow !_E^* \circ G$.

The following theorem establishes a general adjunction between categories of $\mathcal{E}$ -valued functors induced by a functor $P : \mathcal{C} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$.

Before stating it, we need to introduce some notation. For any object c of $\mathcal{C}$, we denote by $\int P(c)$ the category of elements of the functor $P(c) : \mathcal{D}^{\text{op}} \rightarrow \mathbf{Set}$ and by $\pi_c : \int P(c) \rightarrow \mathcal{D}$ the canonical projection functor. We denote by $y_{\mathcal{D}} : \mathcal{D}^{\text{op}} \rightarrow [\mathcal{D}, \mathbf{Set}]$ the Yoneda embedding. For any functor $H : \mathcal{D} \rightarrow \mathcal{E}$ and any object (d, z) of the category $\int P(c)$, we denote by $\kappa_{(d, z)} : H(d) \rightarrow \text{colim}(H \circ \pi_c)$ the canonical colimit arrow.

Theorem 3.12. *Let $\mathcal{C}$ and $\mathcal{D}$ small categories, $\mathcal{E}$ a Grothendieck topos and $P : \mathcal{C} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ a functor. Let $(-)^{\sim} : [\mathcal{D}, \mathcal{E}] \rightarrow [\mathcal{C}, \mathcal{E}]$ be the functor sending to any functor $F : \mathcal{D} \rightarrow \mathcal{E}$ the functor $\tilde{F}$ defined by:*

for any $c \in \mathcal{C}$, $\tilde{F}(c) = \text{colim}(F \circ \pi_c)$, and

for any arrow $f : c \rightarrow c'$, $\tilde{F}(f) : \text{colim}(F \circ \pi_c) \rightarrow \text{colim}(F \circ \pi_{c'})$ is defined by the conditions $\tilde{F}(f) \circ \kappa_{(d, z)}^{\tilde{F}} = \kappa_{(d, P(f)(z))}^{\tilde{F}}$ (for any object (d, z) of the category $\int P(c)$),

and which acts on arrows in the obvious way.

Let $(-)_r : [\mathcal{C}, \mathcal{E}] \rightarrow [\mathcal{D}, \mathcal{E}]$ be the functor assigning to any functor $G : \mathcal{C} \rightarrow \mathcal{E}$ the functor $\text{Hom}_{[\mathcal{C}, \mathcal{E}]}^{\mathcal{E}}(\gamma_{\mathcal{E}}^* \circ y_{\mathcal{D}}^{\sim}, G)$, and acting on the arrows in the obvious way.

Then the functors

$$(\tilde{-}) : [\mathcal{D}, \mathcal{E}] \rightarrow [\mathcal{C}, \mathcal{E}]$$

and

$$(-)_r : [\mathcal{C}, \mathcal{E}] \rightarrow [\mathcal{D}, \mathcal{E}]$$

are adjoint to each other ($(\tilde{-})$ on the left and $(-)_r$ on the right). The unit $\eta : F \rightarrow (\tilde{F})_r$ and the counit $\epsilon^G : (\tilde{G})_r \rightarrow G$ are defined as follows:

For any $d \in \mathcal{D}$, $\eta^F(d) : F(d) \rightarrow (\tilde{F})_r(d)$ is the arrow in $\mathcal{E}$ defined by means of generalized elements by saying that it sends any generalized element $E \rightarrow F(d)$, regarded as a natural transformation $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{D}} d \rightarrow !_E^* \circ F$, to the image of this arrow $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{D}} d \rightarrow !_E^* \circ \tilde{F}$ under the functor $(\tilde{-})$, regarded as a generalized element $E \rightarrow \text{Hom}_{[\mathcal{C}, \mathcal{E}]}^{\mathcal{E}}(\gamma_{\mathcal{E}}^* \circ y_{\mathcal{D}}^{\sim} d, \tilde{F}) = (\tilde{F})_r(d)$.

For any $c \in \mathcal{C}$, $\epsilon^G(c) : (\tilde{G})_r(c) = \text{colim}(G_r \circ \pi_c) \rightarrow G(c)$ is defined by setting, for each object (d, z) of $\int P(c)$, $\epsilon^G(c) \circ \kappa_{(d, z)}^{G_r}$ equal to the arrow $G_r(d) \rightarrow G(c)$ defined by means of generalized elements by saying that a generalized element $x : E \rightarrow G_r(d)$, corresponding to a natural transformation $\bar{x} : !_E^* \circ \gamma_{\mathcal{E}}^* \circ y_{\mathcal{D}} d \cong \gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{D}} d \rightarrow !_E^* \circ \tilde{G}$, is sent to the arrow $E \rightarrow G(c)$ obtained by composing $\bar{x}(c)$ with the component of $\gamma_{\mathcal{E}/E}^*(y_{\mathcal{D}} d(c))$ corresponding to the element of $y_{\mathcal{D}} d(c)$ given by the image of the identity on d via the colimit arrow $\kappa_{(d, z)}^{y_{\mathcal{D}} d} : (y_{\mathcal{D}} d)(d) \rightarrow y_{\mathcal{D}} d(c)$.

Proof For simplicity we shall prove the result only in the case $\mathcal{E} = \mathbf{Set}$, the proof of the general case being entirely analogous (the only care that one has to take is to use generalized elements in place of standard set-theoretic elements).

We shall define a bijective correspondence between the natural transformations $\tilde{F} \rightarrow G$ and the natural transformations $F \rightarrow G_r$, natural in $F \in [\mathcal{D}, \mathbf{Set}]$ and $G \in [\mathcal{C}, \mathbf{Set}]$.

Given a natural transformation $\alpha : F \rightarrow G_r$, we define $\tau(\alpha) : \tilde{F} \rightarrow G$ by setting, for each $c \in \mathcal{C}$, $\tau_\alpha(c) : \tilde{F}(c) \rightarrow G(c)$ equal to the arrow defined as follows. As $\tilde{F}(c)$ is the colimit of the cone $\{\kappa_{(d, z)}^F : F(d) \rightarrow \tilde{F}(c) \mid (d, z) \in \int P(c)\}$, it suffices to define an arrow $u_{(d, z)} : F(d) \rightarrow G(c)$ for each pair (d, z) where $d \in \mathcal{D}$ and $z \in P(c)(d)$, checking that whenever (d, z) and (d', z') are pairs such that for an arrow $f : d \rightarrow d'$ in $\mathcal{D}$, $P(c)(f)(z') = z$ then $u_{(d', z')} \circ F(f) = u_{(d, z)}$; indeed, by the universal property of the colimit, such a family of arrows will induce a unique arrow $\tau_\alpha(c) : \tilde{F}(c) \rightarrow G(c)$ such that $\tau_\alpha(c) \circ \kappa_{(d, z)}^F = u_{(d, z)}$ (for each (d, z))

in $\int P(c)$). We set $u_{(d,z)} : F(d) \rightarrow G(c)$ equal to the function sending every element $x \in F(d)$ to the element $\alpha(d)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}^d}(1_d))$. Let us check that the compatibility condition is satisfied. Given an arrow $f : d \rightarrow d'$ such that $P(c)(f)(z') = z$, we have to verify that for every $x \in F(d)$, $\alpha(d')(Ff(x))(c)(\kappa_{(d',z')}^{y_{\mathcal{D}}^{d'}}(1_{d'})) = \alpha(d)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}^d}(1_d))$. This identity immediately follows from the commutativity of the naturality diagram

$$\begin{array}{ccc} F(d) & \xrightarrow{\alpha(d)} & G_r(d) = \text{Hom}_{[\mathcal{C}, \text{Set}]}(y_{\mathcal{D}}^{\sim} d, G) \\ \downarrow F(f) & & \downarrow G_r(f) = - \circ y_{\mathcal{D}}^{\sim} f \\ F(d') & \xrightarrow{\alpha(d')} & G_r(d') = \text{Hom}_{[\mathcal{C}, \text{Set}]}(y_{\mathcal{D}}^{\sim} d', G) \end{array}$$

for α with respect to the arrow f and that of the diagram

$$\begin{array}{ccc} y_{\mathcal{D}}^{\sim} d'(c) & \xrightarrow{f} & y_{\mathcal{D}}^{\sim} d(c) \\ \kappa_{(d',z')}^{y_{\mathcal{D}}^{d'}} \uparrow & & \kappa_{(d',z')}^{y_{\mathcal{D}}^d} \uparrow \\ y_{\mathcal{D}} d' & \xrightarrow{(y_{\mathcal{D}} f)(d')} & y_{\mathcal{D}} d, \end{array}$$

which is an instance of Proposition 3.1(i).

Indeed, $\alpha(d')(Ff(x))(c)(\kappa_{(d',z')}^{y_{\mathcal{D}}^{d'}}(1_{d'})) = \alpha(d)(x)(y_{\mathcal{D}}^{\sim} f(c)(\kappa_{(d',z')}^{y_{\mathcal{D}}^{d'}}(1_{d'}))) = \alpha(d)(x)(\kappa_{(d',z')}^{y_{\mathcal{D}}^d}(f)) = \alpha(d)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}^d}(1_d))$, where the first equality follows from the commutativity of the first diagram, the second follows from the commutativity of the second diagram, and the last follows from the fact that the family $\{\kappa_{(d,z)}^{y_{\mathcal{D}}^d} \mid (d, z) \in \int P(c)\}$ is a cocone.

To complete the definition of $\tau(\alpha)$, it remains to check that the assignment $c \rightarrow \tau(\alpha)(c)$ defines a natural transformation $\tilde{F} \rightarrow G$. We have to verify that for any arrow $f : c \rightarrow c'$ in $\mathcal{C}$, the diagram

$$\begin{array}{ccc} \tilde{F}(c) & \xrightarrow{\tau(\alpha)(c)} & G(c) \\ \downarrow \tilde{F}(f) & & \downarrow G(f) \\ \tilde{F}(c') & \xrightarrow{\tau(\alpha)(c')} & G(c') \end{array}$$

commutes.

As, by definition of $\tilde{F}(f)$, the diagram

$$\begin{array}{ccc} \tilde{F}(c) & \xrightarrow{\tilde{F}(f)} & \tilde{F}(c') \\ \kappa_{(d,z)}^F \uparrow & \nearrow \kappa_{(d,P(f)(z))}^F & \\ F(d) & & \end{array}$$

commutes and the arrows $\kappa_{(d,z)}^F$ are jointly epimorphic (as they are colimit arrows), the square above commutes if and only if the arrows $G(f) \circ \tau(\alpha)(c) \circ \kappa_{(d,z)}^F$ and $\tau(\alpha)(c') \circ \kappa_{(d,P(f)(z))}^F$ are equal, that is if and only if they take the same values at any element $x \in F(d)$. Let us set $z' = P(f)(d)(z)$. By definition of $\tau(\alpha)$, $\tau(\alpha)(c)(\kappa_{(d,z)}^F(x)) = \alpha(d)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}d}(1_d))$, while $\tau(\alpha)(c')(\kappa_{(d,z')}^F(x)) = \alpha(d)(x)(c')(\kappa_{(d,z')}^{y_{\mathcal{D}}d}(1_d))$. Now, for any $d \in \mathcal{D}$ and any $x \in F(d)$, $\alpha(d)(x)$ is a natural transformation $\tilde{y}d \rightarrow G$; in particular, the diagram

$$\begin{array}{ccc} \tilde{y}d(c) & \xrightarrow{\alpha(d)(x)(c)} & G(c) \\ \downarrow \tilde{y}d(f) & & \downarrow G(f) \\ \tilde{y}d(c') & \xrightarrow{\kappa_{(d,P(f)(z))}^F} & G(c') \end{array}$$

commutes.

The commutativity of this diagram, together with that of the diagram

$$\begin{array}{ccc} \tilde{y}d(c) & \xrightarrow{\tilde{y}d(f)} & \tilde{y}d(c') \\ \uparrow \kappa_{(d,z)}^F & \nearrow \kappa_{(d,z')}^{y_{\mathcal{D}}d} & \\ yd(d) & & \end{array}$$

(which follows by the definition of $\tilde{y}d(f)$) now immediately implies our thesis.

Let us now define a function χ assigning to any natural transformation $\beta : \tilde{F} \rightarrow G$ a natural transformation $\chi(\beta) : F \rightarrow G_r$. For any $d \in \mathcal{D}$, we define $\chi(\beta)(d) : F(d) \rightarrow G_r(d)$ by setting, for each $x \in F(d)$, $\chi(\beta)(x)$ equal to the natural transformation $\beta \circ \tilde{a}_x : yd \rightarrow G$, where $\tilde{a}_x : yd \rightarrow F$ is the natural transformation corresponding, via the Yoneda lemma, to the element $x \in F(c)$.

To verify that $\chi(\beta)$ is well-defined, we have to check that for every arrow $g : d \rightarrow d'$ in $\mathcal{D}$, the diagram

$$\begin{array}{ccc} F(d) & \xrightarrow{\chi(\beta)(d)} & G_r(d) \\ \downarrow F(g) & & \downarrow G_r(g) = - \circ yd \\ F(d') & \xrightarrow{\chi(\beta)(d')} & G_r(d') \end{array}$$

commutes, i.e. for every $x \in F(d)$, $G_r(g)(\chi(\beta)(d)(x)) = \chi(\beta)(d')(F(g)(x))$. Now, $G_r(g)(\chi(\beta)(d)(x)) = \beta \circ \tilde{a}_x \circ yd$, while $\chi(\beta)(d')(F(g)(x)) = \beta \circ \alpha_{F(g)(x)} \circ yd$; but $\alpha_{F(g)(x)} = \alpha_x \circ yd$, from which our thesis follows.

The proof of the fact that the correspondences τ and β are natural in F and G is straightforward and left to the reader.

To conclude the proof of the theorem, it thus remains to show that τ and χ are inverse to each other. The verification of the fact that the unit and counit of the adjunction coincide with those given in the statement of the theorem is straightforward and left to the reader.

Let us show that for any natural transformation $\alpha : F \rightarrow G_r$, $\chi(\tau(\alpha)) = \alpha$. Let us set $\beta = \tau(\alpha)$. We have to prove that for any $d \in \mathcal{D}$, $x \in F(d)$ and $z \in P(c)(d')$, $\alpha(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d} = \chi(\beta)(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d}$ as functions $y_{\mathcal{D}}d(d') \rightarrow G(c)$, i.e. that for any element $g \in y_{\mathcal{D}}d(d')$, $\alpha(d)(x)(c)(\kappa_{(d',z)}^{y_{\mathcal{D}}d}(g)) = (\chi(\beta)(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d})(g)$. Now, by definition of the functor $(\tilde{})$ and of the correspondence τ , the diagram

$$\begin{array}{ccccc} y_{\mathcal{D}}d(c) & \xrightarrow{\tilde{a}_x(c)} & \tilde{F}(c) & \xrightarrow{\beta(c)} & G(c) \\ \kappa_{(d',z)}^{y_{\mathcal{D}}d} \uparrow & & \kappa_{(d',z)}^F \uparrow & \nearrow u_{(d',z)} & \\ y_{\mathcal{D}}d(d') & \xrightarrow{a_x(d')} & F(d') & & \end{array}$$

commutes, and since $\chi(\beta)(x) = \beta \circ \tilde{a}_x : y_{\mathcal{D}}d \rightarrow G$, we have that $(\chi(\beta)(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d})(g)$ is equal to $\alpha(d')(F(g)(x))(c)(\kappa_{(d',z)}^{y_{\mathcal{D}}d'}(1_{d'}))$.

Now, the naturality diagram for α with respect to the arrow $g : d \rightarrow d'$ yields the equality $\alpha(d)(x) \circ y_{\mathcal{D}}g = \alpha(d')(Fg(x))$ and hence the equality $\alpha(d)(x)(c) \circ y_{\mathcal{D}}g(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d'} = \alpha(d')(Fg(x))(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d'}$.

But the commutativity of the diagram

$$\begin{array}{ccc} y_{\mathcal{D}}d'(c) & \xrightarrow{y_{\mathcal{D}}d(g)(c)} & y_{\mathcal{D}}d(c') \\ \kappa_{(d',z)}^{y_{\mathcal{D}}d'} \uparrow & & \uparrow \kappa_{(d',z)}^{y_{\mathcal{D}}d} \\ y_{\mathcal{D}}d'(d') & \xrightarrow{y_{\mathcal{D}}g(d')} & y_{\mathcal{D}}d(d') \end{array}$$

(which follows by definition of the functor $(\tilde{})$) implies that $\alpha(d)(x)(c) \circ y_{\mathcal{D}}g(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d'} = \alpha(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d} \circ y_{\mathcal{D}}g$. Therefore $\alpha(d)(x)(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d} \circ y_{\mathcal{D}}g = \alpha(d')(Fg(x))(c) \circ \kappa_{(d',z)}^{y_{\mathcal{D}}d'}$ and evaluating at $1_{d'}$ yields the desired equality.

Finally, let us show that the composite $\tau \circ \chi$ is equal to the identity. Let $\beta : \tilde{F} \rightarrow G$ be a natural transformation. We have to show that $\beta = \tau(\chi(\beta))$. Let us set $\alpha = \chi(\beta)$. To prove that $\beta = \tau(\alpha)$ it is equivalent to verify that for every object $c \in \mathcal{C}$, any pair (d, z) with $d \in \mathcal{D}$ and $z \in P(c)(d)$ and any element $x \in F(d)$, $\beta(c)(\kappa_{(d,z)}^F(x)) = \tau(\alpha)(c)(\kappa_{(d,z)}^F(x))$. But $\tau(\alpha)(c)(\kappa_{(d,z)}^F(x)) = \alpha(d)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}d}(1_d)) = \chi(\beta)(x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}d}(1_d)) = (\beta \circ \tilde{a}_x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}d}(1_d)) = \beta(c)((\tilde{a}_x)(c)(\kappa_{(d,z)}^{y_{\mathcal{D}}d}(1_d)))$. Thus our thesis follows

from the commutativity of the diagram

$$\begin{array}{ccc} y_{\mathcal{D}} d(c) & \xrightarrow{\tilde{a}_x(c)} & \tilde{F}(c) \\ \kappa_{(d,z)}^{y_{\mathcal{D}} d} \uparrow & & \uparrow \kappa_{(d,z)}^F \\ yd(d) & \xrightarrow{a_x(d)} & F(d), \end{array}$$

which is an immediate consequence of the definition of the functor $(\tilde{-})$. $\square$

Remarks 3.13. (a) Let $f : \mathcal{D} \rightarrow \mathcal{C}$ be a functor between two small categories, and let $P : \mathcal{C} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ be the functor $y_{\mathcal{C}}(-) \circ f^{\text{op}}$. Notice that P is the flat functor corresponding, via Diaconescu's equivalence, to the essential geometric morphism $[\mathcal{D}^{\text{op}}, \mathbf{Set}] \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ induced by the functor $f^{\text{op}} : \mathcal{D}^{\text{op}} \rightarrow \mathcal{C}^{\text{op}}$. The functor $(-)^{\sim} : [\mathcal{D}, \mathcal{E}] \rightarrow [\mathcal{C}, \mathcal{E}]$ coincides with the left Kan extension functor along f , while the right adjoint functor $(-)_r$ coincides with the functor $- \circ f : [\mathcal{C}, \mathcal{E}] \rightarrow [\mathcal{D}, \mathcal{E}]$.

(b) Let $u : \mathbf{Sh}(\mathcal{D}, K) \rightarrow \mathbf{Sh}(\mathcal{C}, J)$ be a geometric morphism. Then for any Grothendieck topos $\mathcal{E}$, u induces as in section 3 a functor

$$\xi_{\mathcal{E}} : \mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) \rightarrow \mathbf{Flat}_J(\mathcal{C}, \mathcal{E}).$$

By Diaconescu's equivalence, the morphism u corresponds to a flat functor $\mathcal{C} \rightarrow \mathbf{Sh}(\mathcal{D}, K)$, which composed with the canonical geometric inclusion $\mathbf{Sh}(\mathcal{D}, K) \hookrightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ yields a functor $P : \mathcal{C} \rightarrow [\mathcal{D}^{\text{op}}, \mathbf{Set}]$. Then $\xi_{\mathcal{E}}$ coincides with the restriction of the functor $(-)^{\sim} : [\mathcal{D}, \mathcal{E}] \rightarrow [\mathcal{C}, \mathcal{E}]$ induced by P as in the theorem to the full subcategories $\mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) \hookrightarrow [\mathcal{D}, \mathcal{E}]$ and $\mathbf{Flat}_J(\mathcal{C}, \mathcal{E}) \hookrightarrow [\mathcal{C}, \mathcal{E}]$. In general, the right adjoint functor $(-)_r : [\mathcal{C}, \mathcal{E}] \rightarrow [\mathcal{D}, \mathcal{E}]$ does not restrict to these subcategories, but if it does, it becomes a right adjoint to the functor $\xi_{\mathcal{E}} : \mathbf{Flat}_K(\mathcal{D}, \mathcal{E}) \rightarrow \mathbf{Flat}_J(\mathcal{C}, \mathcal{E})$.

This notably applies in the case of the geometric morphism

$$p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

considered in section 3.3. In this case, $\mathcal{C} = \mathcal{C}_{\mathbb{T}}$, $\mathcal{D} = \mathcal{K}^{\text{op}}$ and P is the functor $\mathcal{C}_{\mathbb{T}} \rightarrow [\mathcal{K}, \mathbf{Set}]$ sending any geometric formula $\phi(\vec{x})$ to the functor $M \rightarrow [[\vec{x} \cdot \phi]]_M$.

(c) By a basic property of adjoint functors, the (left adjoint) functor

$$(\tilde{-}) : [\mathcal{D}, \mathcal{E}] \rightarrow [\mathcal{C}, \mathcal{E}]$$

is full and faithful if and only if the unit $\eta^F : F \rightarrow (\tilde{F})_r$ is an isomorphism for any F in $[\mathcal{D}, \mathcal{E}]$. It easily follows (by purely formal considerations)

that for every full subcategory $\mathcal{H}$ of $[\mathcal{D}, \mathcal{E}]$ with canonical embedding $i : \mathcal{H} \hookrightarrow [\mathcal{D}, \mathcal{E}]$, the composite functor $(-) \circ i$ is full and faithful if and only if the unit $\eta^F : F \rightarrow (\tilde{F})_r$ is an isomorphism for any F in $\mathcal{H}$.

The following proposition will be useful in the sequel; see section 2.2 for the notation employed in it.

Proposition 3.14. *Let $f : \mathcal{D} \rightarrow \mathcal{C}$ be a functor between small categories, $\mathcal{E}$ a Grothendieck topos and F a functor $\mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$. Then the $\mathcal{E}$ -indexed functor $\int (F \circ f^{\text{op}})_{\mathcal{E}} \rightarrow \int F_{\mathcal{E}}$ sending any object (c, x) of $\int (F \circ f^{\text{op}})_E$ to the object $(f(c), x)$ of $(\int F)_E$ is $\mathcal{E}$ -final if and only if the $\mathcal{E}$ -indexed functor $\int \epsilon^F_{\mathcal{E}} : \int (F \circ \tilde{f}^{\text{op}})_{\mathcal{E}} \rightarrow \int F_{\mathcal{E}}$ induced by the natural transformation $\epsilon^F : F \circ \tilde{f} \rightarrow F$ of Theorem 3.12 is $\mathcal{E}$ -final.*

Proof From the general analysis of section 3.1 we know that for any $c \in \mathcal{C}$, $F \circ \tilde{f}^{\text{op}}(c) = \text{colim}(F \circ f^{\text{op}} \circ \pi_c)$, where π_c is the canonical projection to $\mathcal{D}^{\text{op}}$ from the category $\mathcal{A}_c$ whose objects are the pairs (d, h) , where d is an object of the category $\mathcal{D}$ and h is an arrow $c \rightarrow f(d)$ in $\mathcal{C}$, and whose arrows are the obvious ones. The arrow $\epsilon^F : F \circ \tilde{f} \rightarrow F$ is defined by the property that for any object (d, h) of $\mathcal{A}_c$, $\epsilon^F(c) \circ \kappa_{(d, h)}^F = F(h)$, where $\kappa_{(d, h)}^F : F(f(d)) \rightarrow \text{colim}(F \circ f^{\text{op}} \circ \pi_c)$ is the canonical colimit arrow.

Now, the thesis follows immediately from the fact that for any object of the category $\int (F \circ \tilde{f}^{\text{op}})_E$ of the form $(c, \kappa_{(d, h)}^F \circ y)$, where y is a generalized element $E \rightarrow F(f(d))$, $(\int \epsilon^F)_E((c, \kappa_{(d, h)}^F \circ y)) = (c, F(h) \circ y)$, invoking the fact that the colimit arrows are jointly epimorphic. $\square$

4 Preliminary results on theories of presheaf type

In this section we establish some results on theories of presheaf type which will be important in the sequel.

4.1 A syntactic description of finitely presentable models

For a theory of presheaf type $\mathbb{T}$, it is possible to give an explicit syntactic description of the finitely presentable $\mathbb{T}$ -models; specifically, we have the following result.

Theorem 4.1. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ and $\{\vec{x} . \phi\}$ be a formula over Σ presenting a $\mathbb{T}$ -model $U_{\{\vec{x} . \phi\}}$. Then $U_{\{\vec{x} . \phi\}}$ is isomorphic to the Σ -structure $M_{\{\vec{x} . \phi\}}$ defined as follows:*

- (i) for any sort A over Σ , $M_{\{\vec{x} . \phi\}}A$ is equal to the set $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} . \phi\}, \{x^A . \top\})$ of $\mathbb{T}$ -provably functional geometric formulae from $\{\vec{x} . \phi\}$ to $\{x^A . \top\}$;

- (ii) for any function symbol $f : A_1, \dots, A_n \rightarrow B$ over Σ , the function $M_{\{\vec{x} \cdot \phi\}} f : \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_1} \cdot \top\}) \times \dots \times \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_n} \cdot \top\}) \cong \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_1}, \dots, x^{A_n} \cdot \top\}) \rightarrow \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^B \cdot \top\})$ is equal to $[f] \circ -$ (where $[f] : \{x^{A_1}, \dots, x^{A_n} \cdot \top\} \rightarrow \{x^B \cdot \top\}$ is the morphism in $\mathcal{C}_{\mathbb{T}}$ corresponding to f);
- (iii) for any relation symbol $R \rhd A_1, \dots, A_n$, $M_{\{\vec{x} \cdot \phi\}} R$ is the subobject of $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_1} \cdot \top\}) \times \dots \times \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_n} \cdot \top\}) \cong \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, \{x^{A_1}, \dots, x^{A_n} \cdot \top\})$ given by $[R] \circ -$, (where $[R] : \{x^{A_1}, \dots, x^{A_n} \cdot \top\} \rhd \{x^{A_1}, \dots, x^{A_n} \cdot \top\}$ is the subobject in $\mathcal{C}_{\mathbb{T}}$ corresponding to R).

Proof Recall that we have a canonical equivalence of categories

$$\mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathbf{Set}) \simeq \mathbb{T}\text{-mod}(\mathbf{Set}),$$

sending any flat $J_{\mathbb{T}}$ -continuous functor $F : \mathcal{C}_{\mathbb{T}} \rightarrow \mathbf{Set}$ to the $\mathbb{T}$ -model $F(M_{\mathbb{T}})$, where $M_{\mathbb{T}}$ is the universal model of $\mathbb{T}$ in $\mathcal{C}_{\mathbb{T}}$.

We know from [8] that for any theory of presheaf type $\mathbb{T}$ over a signature Σ , any formula-in-context $\{\vec{x} \cdot \phi\}$ over Σ which presents a $\mathbb{T}$ -model is $\mathbb{T}$ -irreducible, in the sense that every $J_{\mathbb{T}}$ -covering sieve on $\{\vec{x} \cdot \phi\}$ in $\mathcal{C}_{\mathbb{T}}$ is maximal. From this it easily follows that the (flat) representable functor $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, -) : \mathcal{C}_{\mathbb{T}} \rightarrow \mathbf{Set}$ is $J_{\mathbb{T}}$ -continuous; indeed, for any $J_{\mathbb{T}}$ -covering sieve S on an object $\{\vec{y} \cdot \psi\}$ of $\mathcal{C}_{\mathbb{T}}$, any arrow $\gamma : \{\vec{x} \cdot \phi\} \rightarrow \{\vec{y} \cdot \psi\}$ in $\mathcal{C}_{\mathbb{T}}$ factors through one of the arrows belonging to S as the pullback of S along γ is $J_{\mathbb{T}}$ -covering and hence maximal. The image $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, M_{\mathbb{T}})$ under this functor of the model $M_{\mathbb{T}}$, which clearly coincides with the Σ -structure $M_{\{\vec{x} \cdot \phi\}}$ in the statement of the theorem, is therefore a $\mathbb{T}$ -model. In order to deduce our thesis, it thus remains to verify that the model $M_{\{\vec{x} \cdot \phi\}}$ satisfies the universal property of the $\mathbb{T}$ -model presented by the formula $\{\vec{x} \cdot \phi\}$, i.e. that for any $\mathbb{T}$ -model N in $\mathbf{Set}$, the $\mathbb{T}$ -model homomorphisms $M_{\{\vec{x} \cdot \phi\}} = \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, M_{\mathbb{T}}) \rightarrow N$ are in natural bijection with the elements of the set $[[\vec{x} \cdot \phi]]_N$. But the $\mathbb{T}$ -model homomorphisms $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, M_{\mathbb{T}}) \rightarrow N$, are in natural bijection, by the equivalence $\mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathbf{Set}) \simeq \mathbb{T}\text{-mod}(\mathbf{Set})$, with the natural transformations $\text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\}, -) \rightarrow F_N$, that is, by the Yoneda lemma, with the elements of the set $F_N(\{\vec{x} \cdot \phi\}) = [[\vec{x} \cdot \phi]]_N$, as required. $\square$

Remarks 4.2. (a) If $\mathbb{T}$ is a universal Horn theory (in the sense of [3]) and $\phi(\vec{x})$ is a finite conjunction of atomic formulas in a context $\vec{x} = (x^{A_1}, \dots, x^{A_n})$ then the set $M_{\{\vec{x} \cdot \phi\}} A$ can be identified with the set of equivalence classes of terms over Σ of type $A_1, \dots, A_n \rightarrow A$ modulo the equivalence relation which identifies two terms t_1 and t_2 precisely when the sequent $(\phi \vdash_{\vec{x}} t_1 = t_2)$ is provable in $\mathbb{T}$. Indeed, it is shown in [3] (cf. p. 120 therein) that any $\mathbb{T}$ -provably functional geometric formula $\theta(\vec{x}, \vec{y})$

between Horn formulae over Σ is $\mathbb{T}$ -provably equivalent to a formula of the form $\vec{y} = \vec{t}(\vec{x})$, where $\vec{t}$ is a sequence of terms of the appropriate sorts in the context $\vec{x}$.

- (b) Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small category of set-based $\mathbb{T}$ -models and $\phi(\vec{x})$ a geometric formula over Σ presenting a $\mathbb{T}$ -model in $\mathcal{K}$. If the geometric morphism

$$p : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

has the property that its inverse image p^* is full and faithful (for instance, if p is hyperconnected) then $\phi(\vec{x})$ is $\mathbb{T}$ -irreducible and the argument in the proof of Theorem 4.1 applies yielding a syntactic description of the model presented by $\phi(\vec{x})$ as specified in the statement of the theorem. Indeed, denoted by y and y' the Yoneda embeddings respectively of $\mathcal{C}_{\mathbb{T}}$ into $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ and of $\mathcal{K}^{\text{op}}$ into $[\mathcal{K}, \mathbf{Set}]$, we have that $p^*(y\{\vec{x} \cdot \phi\}) \cong y'(M_{\{\vec{x}, \phi\}})$. Now, as $y'(M_{\{\vec{x}, \phi\}})$ is an irreducible object in the topos $[\mathcal{K}, \mathbf{Set}]$ and the property of an object of a topos to be irreducible is reflected by full and faithful inverse images of geometric morphisms, $y\{\vec{x} \cdot \phi\}$ is an irreducible object of the topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$, equivalently $\phi(\vec{x})$ is $\mathbb{T}$ -irreducible, as required.

4.2 Objects of homomorphisms

For any first-order signature Σ and any Grothendieck topos $\mathcal{E}$, we have a $\mathcal{E}$ -indexed category $\Sigma\text{-}\mathbf{str}(\mathcal{E})$ whose fibre at $E \in \mathcal{E}$ is the category $\Sigma\text{-}\mathbf{str}(\mathcal{E})_E$ and whose ‘change of base’ functors are the obvious pullback functors. For any sort A over Σ , we have a $\mathcal{E}$ -indexed forgetful functor $U_A : \Sigma\text{-}\mathbf{str}(\mathcal{E}) \rightarrow \mathcal{E}_{\mathcal{E}}$ assigning to any Σ -structure M in the topos $\mathcal{E}/E$ the object $\overline{MA}$. It is easy to see, by adapting the classical proof in the case $\mathcal{E} = \mathbf{Set}$ and exploiting Theorem 2.12, that the $\mathcal{E}$ -indexed functors U_A jointly create colimits of $\mathcal{E}$ -indexed diagrams defined on $\mathcal{E}$ -final and $\mathcal{E}$ -filtered subcategories $\underline{\mathcal{A}}_{\mathcal{E}}$ of a small $\mathcal{E}$ -indexed category. Indeed, the notion of Σ -structure only involves finite set-indexed limits, and, as we have remarked above, the colimit functor $\text{colim}_{\mathcal{E}} : [\underline{\mathcal{A}}_{\mathcal{E}}, \mathcal{E}_{\mathcal{E}}] \rightarrow \mathcal{E}_{\mathcal{E}}$ preserves them. Moreover, the structure used in interpreting geometric formulae over Σ is all derived from set-indexed finite limits and arbitrary colimits, which are both preserved by $\text{colim}_{\mathcal{E}}$ (the fact that colimits commute with colimits is obvious, while the commutation with finite limits has been observed above). This implies that for any geometric theory $\mathbb{T}$ over Σ , the $\mathcal{E}$ -indexed full subcategory $\underline{\mathbb{T}\text{-mod}}(\mathcal{E})$ of $\underline{\Sigma\text{-}\mathbf{str}}(\mathcal{E})$ is closed in $\underline{\Sigma\text{-}\mathbf{str}}(\mathcal{E})$ under $\mathcal{E}$ -indexed colimits of diagrams defined on $\mathcal{E}$ -final and $\mathcal{E}$ -filtered subcategories $\underline{\mathcal{A}}_{\mathcal{E}}$ of a small $\mathcal{E}$ -indexed category.

The following result asserts that the indexed category of models of a geometric theory in a Grothendieck topos is locally small.

Theorem 4.3. *Let $\mathbb{T}$ be a geometric theory. Then for any $\mathbb{T}$ -models M and N in a Grothendieck topos $\mathcal{E}$ there exists an object $\text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N)$ of $\mathcal{E}$, called the ‘object of $\mathbb{T}$ -model homomorphisms from M to N ’, satisfying the universal property that for any object E of $\mathcal{E}$ the generalized elements $E \rightarrow \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N)$ are in bijective correspondence, naturally in $E \in \mathcal{E}$, with the $\mathbb{T}$ -model homomorphisms $!_E^*(M) \rightarrow !_E^*(N)$ in $\mathbb{T}\text{-mod}(\mathcal{E}/E)$.*

Proof We recall from [5] that for any small category $\mathcal{C}$ and any Grothendieck topos $\mathcal{E}$, for any two functors $F, G : \mathcal{C} \rightarrow \mathcal{E}$ there exists an object $\text{Hom}^{\mathcal{E}}(F, G)$ satisfying the property that for any object E of $\mathcal{E}$ the generalized elements $E \rightarrow \text{Hom}^{\mathcal{E}}(F, G)$ are in bijective correspondence, naturally in $E \in \mathcal{E}$, with the arrows $!_E^* \circ F \rightarrow !_E^* \circ G$ in $[\mathcal{C}, \mathcal{E}/E]$, that is with the natural transformations $!_E^* \circ F \rightarrow !_E^* \circ G$. This implies that for any Grothendieck topos $\mathcal{E}$ the indexed category $[\mathcal{C}, \mathcal{E}]_{\mathcal{E}}$ of functors on $\mathcal{C}$ with values in $\mathcal{E}$ and natural transformations between them is locally small. Clearly, it follows at once that any $\mathcal{E}$ -indexed full subcategory of $[\mathcal{C}, \mathcal{E}]_{\mathcal{E}}$, such as for example the indexed category of J -continuous flat functors on $\mathcal{C}$ with values in $\mathcal{E}$, is also locally small.

Now, every geometric theory $\mathbb{T}$ is Morita-equivalent to the theory of flat $J_{\mathbb{T}}$ -continuous functors $\mathcal{C}_{\mathbb{T}}$ (cf. [21]); in other words, the $\mathcal{E}$ -indexed category $\underline{\mathbb{T}\text{-mod}(\mathcal{E})}$ is equivalent to the $\mathcal{E}$ -indexed category of flat $J_{\mathbb{T}}$ -continuous functors on $\mathcal{C}_{\mathbb{T}}$ with values in $\mathcal{E}$. Hence $\underline{\mathbb{T}\text{-mod}(\mathcal{E})}$ is locally small, as required. $\square$

Remarks 4.4. (a) The assignment $(M, N) \rightarrow \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N)$ is functorial both in M and N ; that is, any homomorphism $f : M \rightarrow M'$ in $\mathbb{T}\text{-mod}(\mathcal{E})$ (resp. any homomorphism $g : N' \rightarrow N$ in $\mathbb{T}\text{-mod}(\mathcal{E})$) induces an arrow

$$\text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(f, N) : \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M', N) \rightarrow \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N)$$

(resp. an arrow

$$\text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, g) : \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N') \rightarrow \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N))$$

functorially in f (resp. functorially in g).

(b) For any Grothendieck topos $\mathcal{E}$ and $\mathbb{T}$ -models M and N in $\mathcal{E}$, we have a canonical embedding

$$\text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{F}}(M, N) \rightarrow \prod_{A \text{ sort of } \Sigma} NA^{MA}$$

induced by arrows

$$\pi_A : \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N) \rightarrow NA^{MA}$$

(for any sort A of Σ) defined in terms of generalized elements as follows: π_A sends any arrow $E \rightarrow \underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)$ in $\mathcal{E}$, corresponding to a $\mathbb{T}$ -model homomorphism $r : !_E^*(M) \rightarrow !_E^*M$ in $\mathcal{E}/E$, to the arrow $E \rightarrow NA^{MA}$ whose transpose is the ‘evaluation’ $!_E^*(MA) \rightarrow !_E^*(NA)$ at A of r .

- (c) For any $\mathbb{T}$ -models M and N in a Grothendieck topos $\mathcal{E}$ and any geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$, there is a canonical arrow

$$f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)) \rightarrow \underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{F})}^{\mathcal{F}}(f^*(M), f^*(N)).$$

Indeed, this arrow corresponds, by the universal property of the object $\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{F})}^{\mathcal{F}}(f^*(M), f^*(N))$ to the $\mathbb{T}$ -model homomorphism

$$f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)) \times f^*(M) \rightarrow f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)) \times f^*(N).$$

in the topos $\mathcal{F}/(f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)))$ whose first component, at any sort A , is the canonical projection and whose second component at A is the arrow

$$f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)) \times f^*(MA) \cong f^*(\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N) \times MA) \rightarrow f^*(NA).$$

obtained by taking the image under f^* of the arrow

$$\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N) \times MA \rightarrow NA$$

given by the transpose of the arrow $\pi_A : \underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N) \rightarrow NA^{MA}$ defined above.

- (d) Since the $\mathcal{E}$ -indexed category $\mathbb{T}\text{-mod}(\mathcal{E})$ is locally small (cf. the proof of Theorem 4.3), we have a $\mathcal{E}$ -indexed hom functor

$$\underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(-, -) : \mathbb{T}\text{-mod}(\mathcal{E}) \times \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$$

The following proposition provides an explicit description of the generalized elements of the objects of homomorphism of Theorem 4.3 in a particular case of interest. In the statement of the proposition, for a $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, we denote by $\text{Hom}_{\mathcal{E}}(E, M)$ the Σ -structure in **Set** obtained as the image of M under the product-preserving functor $\text{Hom}_{\mathcal{E}}(E, -) : \mathcal{E} \rightarrow \mathbf{Set}$.

Proposition 4.5. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , M a model of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$, c a set-based $\mathbb{T}$ -model and E an object of $\mathcal{E}$. Then the generalized elements $x : E \rightarrow \underline{\text{Hom}}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ correspond bijectively to the Σ -structure homomorphisms $\xi_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$.*

Proof By definition of $Hom_{\mathbb{T}\text{-mod}_m(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$, a generalized element $E \rightarrow Hom_{\mathbb{T}\text{-mod}_m(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ corresponds precisely to a $\mathbb{T}$ -model homomorphism $\gamma_{\mathcal{E}/E}^*(c) \rightarrow !_E^*(M)$ in the topos $\mathcal{E}/E$. Concretely, such a $\mathbb{T}$ -model homomorphism consists of a family of arrows $\tau_A : \gamma_{\mathcal{E}/E}^*(cA) \rightarrow !_E^*(MA)$ in $\mathcal{E}/E$ indexed by the sorts A over Σ which satisfies the preservation conditions defining the notion of Σ -structure homomorphism. Now, each of the arrows $\tau_A : \gamma_{\mathcal{E}/E}^*(cA) \rightarrow !_E^*(MA)$ corresponds, via the adjunction between $\gamma_{\mathcal{E}/E}^*$ and the global section functor on the topos $\mathcal{E}/E$, to a function $\xi_A : cA \rightarrow Hom_{\mathcal{E}}(E, MA)$, and it is immediate to see that the above-mentioned preservation conditions translate precisely into the requirement that the arrows $\xi_A : cA \rightarrow Hom_{\mathcal{E}}(E, M)$ should yield a Σ -structure homomorphism $c \rightarrow Hom_{\mathcal{E}}(c, M)$. $\square$

Remark 4.6. If the model c is finitely presentable also as a $\mathbb{T}_c$ -model, where $\mathbb{T}_c$ is the cartesianization of $\mathbb{T}$ as defined in section 6.4 (notice that this is always the case if Σ is finite and $\mathbb{T}$ has only a finite number of axioms, cf. Theorem 6.4 [6]) then, if $\phi(\vec{x})$ is a formula over Σ which presents it, the Σ -structure homomorphisms $c \rightarrow Hom_{\mathcal{E}}(E, M)$ can be identified with the elements of the interpretation of the formula $\phi(\vec{x})$ in the Σ -structure $Hom_{\mathcal{E}}(E, M)$.

4.3 Strong finite presentability

We shall show in this section that the finitely presentable models of a theory $\mathbb{T}$ of presheaf type enjoy a strong form of finite presentability by a geometric formula with respect to the models of $\mathbb{T}$ in arbitrary Grothendieck toposes.

Let $\mathbb{T}$ be a geometric theory over a signature Σ , c a set-based model of $\mathbb{T}$, $\phi(\vec{x})$ a geometric formula over Σ and $\vec{u}$ an element of $[[\vec{x} \cdot \phi]]_c$. Then for any model M of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$ there is a canonical arrow

$$\tau_{\phi(\vec{x}), \vec{u}}^M : Hom_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M) \rightarrow [[\vec{x} \cdot \phi]]_M$$

in $\mathcal{E}$, defined on generalized elements as follows: $\tau_{\phi(\vec{x}), \vec{u}}^M$ sends a generalized element

$$E \rightarrow Hom_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M),$$

corresponding under the bijection of Proposition 4.5 to a Σ -structure homomorphism $f : c \rightarrow Hom_{\mathcal{E}}(E, M)$, to the image of $\vec{u}$ under f ; notice that such element indeed belongs to $Hom_{\mathcal{E}}(E, [[\vec{x} \cdot \phi]]_M)$ since, as f is a Σ -structure homomorphism, the image of $[[\vec{x} \cdot \phi]]_c$ under f is contained in $[[\vec{x} \cdot \phi]]_{Hom_{\mathcal{E}}(E, M)}$, which is contained in $Hom_{\mathcal{E}}(E, [[\vec{x} \cdot \phi]]_M)$ since the functor $Hom_{\mathcal{E}}(E, -)$ is cartesian.

Definition 4.7. Let $\mathbb{T}$ be a geometric theory over a signature Σ , and c a model of $\mathbb{T}$ in **Set**. The $\mathbb{T}$ -model c is said to be *strongly finitely presented* if there exists a geometric formula $\phi(\vec{x})$ over Σ and a finite string of elements $\vec{u}$ of $[[\vec{x} \cdot \phi]]_c$, called the *strong generators* of c , such that for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ the arrow

$$\tau_{\phi(\vec{x}), \vec{u}}^M : \text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(M_{\{\vec{x}, \phi\}}), M) \rightarrow [[\vec{x} \cdot \phi]]_M$$

is an isomorphism (equivalently, the Σ -structure homomorphisms $\xi : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ are in natural bijection with the generalized elements $E \rightarrow [[\vec{x} \cdot \phi]]_M$ via the assignment $\xi \rightarrow \xi(\vec{u})$).

Remark 4.8. If the latter condition in the definition is satisfied by all models M of $\mathbb{T}$ inside Grothendieck toposes for $E = 1_{\mathcal{E}}$ then it is true in general, by the localization principle.

Theorem 4.9. Let $\mathbb{T}$ be a theory of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$, where $\mathcal{K}$ is a small subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$, M a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$ and c a $\mathbb{T}$ -model in $\mathcal{K}$. Then there is a natural bijective correspondence between the Σ -structure homomorphisms $c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and the elements of the set $\text{Hom}_{\mathcal{E}}(E, F_M c)$, where F_M is the flat functor $\mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ corresponding to the model M via the canonical Morita-equivalence for $\mathbb{T}$.

Proof We can clearly suppose without loss of generality $E = 1_{\mathcal{E}}$. The adjunction between $\gamma_{\mathcal{E}}^*$ and the global sections functor $\Gamma_{\mathcal{E}} : \mathcal{E} \rightarrow \mathbf{Set}$ provides a natural bijective correspondence between the elements of the set $\text{Hom}_{\mathcal{E}}(E, F_M c)$ and the natural transformations $\gamma_{\mathcal{E}}^* \circ y c \rightarrow F_M$. By the canonical Morita-equivalence

$$\tau_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E}),$$

for $\mathbb{T}$, such natural transformations are in natural bijective correspondence with the $\mathbb{T}$ -model homomorphisms $\gamma_{\mathcal{E}}^*(c) \cong \tau(\gamma_{\mathcal{E}}^* \circ y c) \rightarrow \tau(F_M) = M$. But these homomorphisms are, by Proposition 4.5, in natural bijective correspondence with the Σ -structure homomorphisms $c \rightarrow \text{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, M)$, as required. $\square$

Corollary 4.10. Let $\mathbb{T}$ be a theory of presheaf type and $\phi(\vec{x})$ a formula presenting a $\mathbb{T}$ -model $M_{\{\vec{x}, \phi\}}$. Then $M_{\{\vec{x}, \phi\}}$ is strongly finitely presented by $\phi(\vec{x})$, i.e. for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ and any object E of $\mathcal{E}$, the Σ -structure homomorphisms $M_{\{\vec{x}, \phi\}} \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ correspond bijectively to the elements of the set $\text{Hom}_{\mathcal{E}}(E, [[\vec{x} \cdot \phi]]_M)$, via the assignment sending a Σ -structure homomorphism $M_{\{\vec{x}, \phi\}} \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ to the image under it of the generators of $M_{\{\vec{x}, \phi\}}$.

Proof Clearly, we can suppose without loss of generality $E = 1_{\mathcal{E}}$.

By the results of [5], the canonical Morita-equivalence

$$\mathbf{Flat}(\mathbf{f.p.T-mod}(\mathbf{Set})^{\mathrm{op}}, \mathcal{E}) \simeq \mathbf{T-mod}(\mathcal{E})$$

for $\mathbf{T}$ can be described as the correspondence sending, on one hand, to any $\mathbf{T}$ -model M the flat functor $F_M := \underline{\mathrm{Hom}}_{\mathbf{T-mod}(\mathcal{E})}(\gamma_{\mathcal{E}}^*(c), M)$ and conversely to any flat functor $F : \mathbf{f.p.T-mod}(\mathbf{Set})^{\mathrm{op}} \rightarrow \mathcal{E}$ the model $\tilde{F}(M_{\mathbf{T}})$, where $\tilde{F} : \mathcal{C}_{\mathbf{T}} \rightarrow \mathcal{E}$ is the extension of F to the syntactic category $\mathcal{C}_{\mathbf{T}}$ (in the sense of section 3.3). By Theorem 3.5, for any $\mathbf{T}$ -model M in a Grothendieck topos $\mathcal{E}$, there is an isomorphism $z_{(M, \{\vec{x} \cdot \phi\})} : F_M(M_{\{\vec{x} \cdot \phi\}}) \cong \tilde{F}_M(\{\vec{x} \cdot \phi\}) = [[\vec{x} \cdot \phi]]_M$. Thus, applying Theorem 4.9 to the model $c = M_{\{\vec{x} \cdot \phi\}}$, we obtain a bijective correspondence between the Σ -structure homomorphisms $M_{\{\vec{x} \cdot \phi\}} \rightarrow \mathrm{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, M)$ and the elements of the set $\mathrm{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, [[\vec{x} \cdot \phi]]_M)$. It remains to show that this correspondence can be identified with the assignment sending a Σ -structure homomorphism $M_{\{\vec{x} \cdot \phi\}} \rightarrow \mathrm{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, M)$ to the image under it of the generators of $M_{\{\vec{x} \cdot \phi\}}$. Recall that the bijection of Theorem 4.9 can be described as follows: any Σ -structure homomorphism $f : M_{\{\vec{x} \cdot \phi\}} \rightarrow \mathrm{Hom}_{\mathcal{E}}(1_{\mathcal{E}}, M)$, corresponding to a $\mathbf{T}$ -model homomorphism $\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}}) \rightarrow M$, and hence, via the canonical Morita-equivalence for $\mathbf{T}$, to a natural transformation $\gamma_{\mathcal{E}}^* \circ y M_{\{\vec{x} \cdot \phi\}} \cong F_{\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}})} \rightarrow F_M$ (where y is the Yoneda embedding $\mathbf{f.p.T-mod}(\mathbf{Set})^{\mathrm{op}} \hookrightarrow [\mathbf{f.p.T-mod}(\mathbf{Set}), \mathbf{Set}]$), is sent to the global element of $F_M(M_{\{\vec{x} \cdot \phi\}})$ corresponding via the Yoneda lemma to this transformation. To deduce our thesis, it thus remains to verify that the canonical isomorphism of functors $\gamma_{\mathcal{E}}^* \circ y M_{\{\vec{x} \cdot \phi\}} = \gamma_{\mathcal{E}}^* \circ F_{M_{\{\vec{x} \cdot \phi\}}} \cong F_{\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}})}$, when evaluated in $M_{\{\vec{x} \cdot \phi\}}$ and composed with the isomorphism $z_{(\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}}), \{\vec{x} \cdot \phi\})} : F_{\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}})}(M_{\{\vec{x} \cdot \phi\}}) \cong [[\vec{x} \cdot \phi]]_{\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}})} \cong \gamma_{\mathcal{E}}^*([[\vec{x} \cdot \phi]]_{M_{\{\vec{x} \cdot \phi\}}})$ sends the coproduct component of $\gamma_{\mathcal{E}}^*(y M_{\{\vec{x} \cdot \phi\}}(M_{\{\vec{x} \cdot \phi\}}))$ corresponding to the identity on $M_{\{\vec{x} \cdot \phi\}}$ to the coproduct component of $\gamma_{\mathcal{E}}^*([[\vec{x} \cdot \phi]]_{M_{\{\vec{x} \cdot \phi\}}})$ corresponding to the generators of $M_{\{\vec{x} \cdot \phi\}}$. By the naturality in $\mathcal{E}$ of the Morita-equivalence for $\mathbf{T}$ we can clearly suppose, without loss of generality, $\mathcal{E}$ equal to $\mathbf{Set}$. But from the proof of Theorem 4.1, it is clear that the generators of $M_{\{\vec{x} \cdot \phi\}}$ correspond to the identity on $\{\vec{x} \cdot \phi\}$ via the Yoneda lemma; hence they correspond to the identity on $M_{\{\vec{x} \cdot \phi\}}$ via the above-mentioned bijection, as required.

Remark 4.11. We can express the bijective correspondence of Corollary 4.10 by saying that for any Grothendieck topos $\mathcal{E}$ the $\mathcal{E}$ -indexed functor $[[\vec{x} \cdot \phi]]_- : \mathbf{T-mod}(\mathcal{E}) \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ assigning to any $\mathbf{T}$ -model M the interpretation of $\phi(\vec{x})$ in M is represented as a $\mathcal{E}$ -indexed functor by the object $\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}})$, in the sense of being naturally isomorphic to the $\mathcal{E}$ -representable functor $\mathrm{Hom}_{\mathbf{T-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(M_{\{\vec{x} \cdot \phi\}}), -)$.

□

4.4 Semantic $\mathcal{E}$ -finite presentability

In this section, we introduce a semantic notion of $\mathcal{E}$ -finite presentability of a model M of a geometric theory $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$, which generalizes the classical notion in the context of finitely accessible categories, and show that all the ‘constant’ finitely presentable models of a theory of presheaf type in a Grothendieck topos $\mathcal{E}$ are $\mathcal{E}$ -finitely presentable.

For any Grothendieck topos $\mathcal{E}$, we can make any small category $\mathcal{C}$ into a $\mathcal{E}$ -indexed category $\underline{\mathcal{C}}_{\mathcal{E}}$ defined by: $\mathcal{C}_E = \mathcal{C}$ for all $E \in \mathcal{E}$ and $\mathcal{C}_{\alpha} = 1_{\mathcal{C}}$ for all arrows α in $\mathcal{E}$. To any (set-indexed) diagram $D : \mathcal{C} \rightarrow \mathcal{E}$ corresponds a $\mathcal{E}$ -indexed functor $\underline{D}_{\mathcal{E}} : \underline{\mathcal{C}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ defined by: for any object E of $\mathcal{E}$, $\underline{D}_{\mathcal{E}} E = !_E^* \circ D$, where $!_E^* : \mathcal{E} \rightarrow \mathcal{E}/E$ is the pullback functor along the unique arrow $E \rightarrow 1_{\mathcal{E}}$ in $\mathcal{E}$. Since the pullback functors preserve all small limits and colimits, giving a colimiting cocone (resp. a limiting cone) on the diagram D in the classical sense is equivalent to giving a $\mathcal{E}$ -indexed colimiting cocone (resp. limiting cone) over the $\mathcal{E}$ -indexed diagram $\underline{D}_{\mathcal{E}}$. Since the $\mathcal{E}$ -indexed category $\underline{\mathcal{E}}_{\mathcal{E}}$ is complete, for any $\mathcal{E}$ -indexed category $\underline{\mathcal{A}}_{\mathcal{E}}$, the $\mathcal{E}$ -indexed functor category $[\underline{\mathcal{A}}_{\mathcal{E}}, \underline{\mathcal{E}}_{\mathcal{E}}]$ is also complete (since limits are computed pointwise); in particular, it has limits of diagrams defined on $\mathcal{E}$ -indexed categories of the form $\underline{\mathcal{C}}_{\mathcal{E}}$. On the other hand, if $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -final $\mathcal{E}$ -filtered subcategory of a small $\mathcal{E}$ -indexed category, the cocompleteness of $\underline{\mathcal{E}}_{\mathcal{E}}$ as a $\mathcal{E}$ -indexed category ensures that there exists a well-defined $\mathcal{E}$ -indexed colimit functor $\text{colim}_{\mathcal{E}} : [\underline{\mathcal{A}}_{\mathcal{E}}, \underline{\mathcal{E}}_{\mathcal{E}}] \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$. It is easy to see, by mimicking the classical proof of the fact that finite limits commute with filtered colimits, that this functor preserves limits of diagrams defined on $\mathcal{E}$ -indexed categories of the form $\underline{\mathcal{C}}_{\mathcal{E}}$ for a finite category $\mathcal{C}$.

Definition 4.12. Let $\mathbb{T}$ be a geometric theory and M be a model of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$. Then M is said to be $\mathcal{E}$ -finitely presentable if the $\mathcal{E}$ -indexed functor $\text{Hom}_{\underline{\mathbb{T}\text{-mod}}(\mathcal{E})}^{\mathcal{E}}(M, -) : \underline{\mathbb{T}\text{-mod}}(\mathcal{E}) \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ of section 4.2 preserves $\mathcal{E}$ -filtered colimits (of $\mathcal{E}$ -final and $\mathcal{E}$ -filtered subcategories of a small $\mathcal{E}$ -indexed category).

Theorem 4.13. Let $\mathbb{T}$ be a theory of presheaf type and c a finitely presentable $\mathbb{T}$ -model. Then for any Grothendieck topos $\mathcal{E}$, the $\mathbb{T}$ -model $\gamma_{\mathcal{E}}^*(c)$ is $\mathcal{E}$ -finitely presentable.

Proof If c is presented by a geometric formula $\phi(\vec{x})$ over the signature Σ of $\mathbb{T}$, by Corollary 4.10, the functor $\text{Hom}_{\underline{\mathbb{T}\text{-mod}}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ is naturally isomorphic to the functor $[[\vec{x} . \phi]]_- : \underline{\mathbb{T}\text{-mod}}(\mathcal{E}) \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$.

This latter functor preserves $\mathcal{E}$ -filtered colimits (of $\mathcal{E}$ -final and $\mathcal{E}$ -filtered subcategories of a small $\mathcal{E}$ -indexed category) since the structure used in interpreting geometric formulae over Σ is all derived from set-indexed finite limits and arbitrary colimits, which, as remarked above, are all preserved by

$\mathcal{E}$ -indexed functors of the form $\text{colim}_{\mathcal{E}} : [\underline{\mathcal{A}}_{\mathcal{E}}, \underline{\mathcal{E}}_{\mathcal{E}}] \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ where $\underline{\mathcal{A}}_{\mathcal{E}}$ is a $\mathcal{E}$ -final $\mathcal{E}$ -filtered subcategory of a small $\mathcal{E}$ -indexed category. $\square$

Remarks 4.14. (a) The proof of the theorem shows that, more generally, for any strongly finitely presentable model c of a geometric theory $\mathbb{T}$ in the sense of Definition 4.7 (for instance, a finite model of $\mathbb{T}$ if the signature of $\mathbb{T}$ is finite - note that such a model is finitely presented with respect to the empty theory over the signature of $\mathbb{T}$ by Theorem 6.4 [6]) and any Grothendieck topos $\mathcal{E}$, the $\mathbb{T}$ -model $\gamma_{\mathcal{E}}^*(c)$ is $\mathcal{E}$ -finitely presentable.

(b) We could have more strongly required in Definition 4.12 the preservation of all existing colimits of diagrams defined on $\mathcal{E}$ -filtered $\mathcal{E}$ -indexed categories, in the sense of Definition 2.1. The theorem remains true with respect to this stronger notion, but a smallness condition for the domain category, such as the requirement for it to be a $\mathcal{E}$ -final and $\mathcal{E}$ -filtered $\mathcal{E}$ -indexed subcategory of a small $\mathcal{E}$ -indexed category, is necessary to dispose of the explicit characterization of filtered colimits provided by Corollary 2.24.

The following proposition provides an explicit characterization of the $\mathcal{E}$ -finitely presentable models of a geometric theory $\mathbb{T}$.

Proposition 4.15. *Let $\mathbb{T}$ be a geometric theory, $\mathcal{E}$ a Grothendieck topos and c a $\mathbb{T}$ -model in $\mathcal{E}$. Then c is $\mathcal{E}$ -finitely presentable if and only if for every $\mathcal{E}$ -indexed diagram $D : \underline{\mathcal{A}}_{\mathcal{E}} \rightarrow \underline{\mathbb{T}\text{-mod}}(\mathcal{E})$ defined on a $\mathcal{E}$ -filtered $\mathcal{E}$ -final subcategory $\underline{\mathcal{A}}_{\mathcal{E}}$ of a small $\mathcal{E}$ -indexed category with $\mathcal{E}$ -indexed colimiting cocone (M, μ) (we denote by $\mu_{(E,x)} : D_E(x) \rightarrow !_E^*(M)$ the colimit arrows), the following conditions are verified:*

- (a) *For any object E of $\mathcal{E}$ and $\mathbb{T}$ -model homomorphism $h : !_E^*(c) \rightarrow !_E^*(M)$ in the topos $\mathcal{E}/E$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ an object x_i of $\mathcal{A}_{E_i}$ and a $\mathbb{T}$ -model homomorphism $\alpha_i : !_{E_i}^*(c) \rightarrow D_{E_i}(x_i)$ in the topos $\mathcal{E}/E_i$ such that for all $i \in I$, $\mu_{(E,x_i)} \circ \alpha_i = e_i^*(h)$;*
- (b) *For any pairs (x, y) and (x', y') , where x and x' are objects of $\mathcal{A}_E$, f is an arrow $F \rightarrow E$ in $\mathcal{E}$, y is a $\mathbb{T}$ -model homomorphism $!_F^*(c) \rightarrow f^*(D_E(x))$ in $\mathcal{E}/F$ and y' is a $\mathbb{T}$ -model homomorphism $!_{F'}^*(c) \rightarrow f'^*(D_E(x'))$ in $\mathcal{E}/F'$, we have $f^*(\mu_E(x)) \circ y = f'^*(\mu_E(x')) \circ y'$ if and only if there exists an epimorphic family $\{f_i : F_i \rightarrow F \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ arrows $g_i : \mathcal{A}_{f \circ f_i}(x) \rightarrow z_i$ and $h_i : \mathcal{A}_{f' \circ f_i}(x') \rightarrow z_i$ in the category $\mathcal{A}_{F_i}$ such that $D_{F_i}(g_i) \circ f_i^*(y) = D_{F_i}(h_i) \circ f_i'^*(y')$.*

Proof The proposition follows as an immediate consequence of Corollary 2.24, applied to the composite $\mathcal{E}$ -indexed functor $\text{Hom}_{\underline{\mathbb{T}\text{-mod}}(\mathcal{E})}^{\mathcal{E}}(c, -) \circ D$. $\square$

Remark 4.16. By the construction of (small) $\mathcal{E}$ -indexed colimits in the $\mathcal{E}$ -indexed category $\mathbb{T}\text{-mod}(\mathcal{E})$, a $\mathcal{E}$ -indexed cocone (M, μ) over a diagram $D : \mathcal{A}_{\mathcal{E}} \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$ as in the statement of the proposition is colimiting if and only if for every sort A over the signature of $\mathbb{T}$, $UA((M, \mu))$ is a colimiting cocone over the diagram $U_A \circ D$ in $\underline{\mathcal{E}}_{\mathcal{E}}$.

5 Semantic criteria for a theory to be of presheaf type

In this section we establish our main characterization theorem providing necessary and sufficient conditions for a geometric theory to be of presheaf type. These conditions are entirely expressed in terms of the models of the theory in arbitrary Grothendieck toposes.

We shall first prove the theorem and then proceed to reformulate its conditions in more concrete terms so for them to be directly applicable in practice.

5.1 The characterization theorem

Recall from [5] that the classifying topos of a theory of presheaf type $\mathbb{T}$ can be canonically represented as the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ of set-valued functors on the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ of finitely presentable $\mathbb{T}$ -models in $\mathbf{Set}$. This is not the only possible representation of the classifying topos of $\mathbb{T}$ as a presheaf topos, but for any small category $\mathcal{K}$, $\mathbb{T}$ is classified by the topos $[\mathcal{K}, \mathbf{Set}]$ if and only if the Cauchy-completion of $\mathcal{K}$ is equivalent to $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$.

Theorem 5.1. *Let $\mathbb{T}$ be a geometric theory over a signature Σ and let $\mathcal{K}$ be a full subcategory of $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$. Then $\mathbb{T}$ is a theory of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$ if and only if all of the following conditions are satisfied:*

- (i) *For any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, the functor $H_M := \text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), M) : \mathcal{K}^{op} \rightarrow \mathcal{E}$ is flat;*
- (ii) *The extension $\tilde{H}_M : \mathcal{C}_{\mathbb{T}} \rightarrow \mathcal{E}$ of the functor $H_M : \mathcal{K}^{op} \rightarrow \mathcal{E}$ to the syntactic category $\mathcal{C}_{\mathbb{T}}$ (in the sense of section 3.3) satisfies the property that the canonical morphism $\tilde{H}_M(M_{\mathbb{T}}) \rightarrow M$ is an isomorphism;*
- (iii) *Any of the following conditions (equivalent, under assumptions (i) and (ii)) is satisfied:*

- (a) *For any model c in $\mathcal{K}$, $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ and geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$, the canonical morphism*

$$f^*(\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)) \rightarrow \text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{F})}^{\mathcal{F}}(\gamma_{\mathcal{F}}^*(c), f^*(M))$$

provided by Remark 4.4(c) using the identification $\gamma_{\mathcal{F}}^*(c) \cong f^*(\gamma_{\mathcal{E}}^*(c))$, is an isomorphism;

- (b) For any flat functor $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$, the canonical natural transformation

$$\eta^F : F \rightarrow \text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), \tilde{F}(M_{\mathbb{T}})) \cong \text{Hom}_{\mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^* \circ y_{\mathcal{K}}(-), \tilde{F})$$

of Theorem 3.12 is an isomorphism, where $y_{\mathcal{K}} : \mathcal{K}^{\text{op}} \hookrightarrow \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathbf{Set})$ is the Yoneda embedding;

- (c) The functor

$$u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

of section 3.3 is full and faithful.

Proof Let us begin by proving that each of the three listed conditions are necessary.

By the results in [5], if $\mathbb{T}$ is of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$ then we have a Morita-equivalence

$$\tau_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

which can be supposed canonical without loss of generality, i.e. which sends any finitely presentable $\mathbb{T}$ -model c in $\mathcal{K}$ to the functor $\gamma_{\mathcal{E}}^* \circ y_{\mathcal{K}} c$. It follows that for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, with corresponding flat functor F_M under this Morita-equivalence, the object $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ is isomorphic to the object $\text{Hom}_{\mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^* \circ y_{\mathcal{K}} c, F_M) \cong F_M(c)$, naturally in M and in c . Therefore the functor

$$\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), M) : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$$

is flat, it being isomorphic to F_M . This proves that condition (i) of the theorem is satisfied.

Next, we notice that the left-to-right functor forming the canonical Morita-equivalence $\tau_{\mathcal{E}}$ for $\mathbb{T}$ can be described as follows: for any flat functor $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$, the $\mathbb{T}$ -model corresponding to it is naturally isomorphic to the model $\tilde{F}(M_{\mathbb{T}})$. Indeed, as the Morita-equivalence for $\mathbb{T}$ is canonical, it is induced by the canonical geometric morphism (in fact, equivalence) $p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$, i.e. it is given by the composite of the induced equivalence $\mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \simeq \mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E})$ with the canonical equivalence $\mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$ sending any flat $J_{\mathbb{T}}$ -continuous functor G on $\mathcal{C}_{\mathbb{T}}$ to the $\mathbb{T}$ -model $G(M_{\mathbb{T}})$.

The fact that $\tau_{\mathcal{E}}$ is an equivalence thus implies that the canonical arrow $\tilde{H}_M \rightarrow M$ is an isomorphism. This shows that condition (ii) of the theorem is satisfied.

The fact that condition $(iii)(c)$ is satisfied also immediately follows from the fact that $\tau_{\mathcal{E}}$ is an equivalence.

We have thus proved that conditions (i) , (ii) and $(iii)(c)$ of the theorem are necessary.

Let us now show that conditions (i) , (ii) and $(iii)(b)$ are, all together, sufficient for the theory $\mathbb{T}$ to be classified by the topos $[\mathcal{K}, \mathbf{Set}]$.

First, we notice that condition $(iii)(b)$ implies condition $(iii)(a)$ under the assumption that condition (i) holds. Indeed, for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, condition (i) ensures that the functor $H_M : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ is flat. On the other hand, for any geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$ condition $(iii)(b)$ yields, in view of the naturality in $\mathcal{E}$ of the operation $(-)$, a natural isomorphism between the flat functor $f^* \circ H_M$ and the functor $H_{f^*(M)}$. This ensures that the requirement of condition $(iii)(a)$ is satisfied.

Under conditions (i) , (ii) and $(iii)(b)$, we shall construct, for any Grothendieck topos $\mathcal{E}$, a categorical equivalence

$$\mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

natural in $\mathcal{E}$.

We shall define two functors $G_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$ and $H_{\mathcal{E}} : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E})$ which are natural in $\mathcal{E}$ and categorical inverses to each other (up to isomorphism).

We set $H_{\mathcal{E}}(M)$ equal to the functor

$$Hom_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), M) : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}.$$

This assignment is natural in M (cf. Remark 4.4) and hence defines a functor $H_{\mathcal{E}} : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E})$, which is natural in $\mathcal{E}$ by condition $(iii)(a)$.

In the converse direction, for any flat functor $F : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$ we set $G_{\mathcal{E}}(F) = \tilde{F}(M_{\mathbb{T}})$. Clearly, this assignment is natural in F and defines a functor $G_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$.

For each Grothendieck topos $\mathcal{E}$, the functors $G_{\mathcal{E}}$ and $H_{\mathcal{E}}$ are categorical inverses to each other (up to isomorphism). Indeed, condition $(iii)(b)$ ensures that $H_{\mathcal{E}} \circ G_{\mathcal{E}}$ is naturally isomorphic to the identity, while condition (ii) ensures that $G_{\mathcal{E}} \circ H_{\mathcal{E}}$ is naturally isomorphic to the identity.

Now, the functors

$$G_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$$

and

$$H_{\mathcal{E}} : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E})$$

are natural in $\mathcal{E}$ and therefore induce geometric morphisms

$$G : [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

and

$$H : \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}}) \rightarrow [\mathcal{K}, \mathbf{Set}] .$$

The fact that G and H are two halves of a categorical equivalence between $[\mathcal{K}, \mathbf{Set}]$ and $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ follows immediately from the fact that for every Grothendieck topos $\mathcal{E}$ there are natural isomorphisms between $H_{\mathcal{E}} \circ G_{\mathcal{E}}$ and the identity functor on $\mathbf{Flat}(\mathcal{K}^{\mathrm{op}}, \mathcal{E})$ and between $G_{\mathcal{E}} \circ H_{\mathcal{E}}$ and the identity functor on $\mathbb{T}\text{-mod}(\mathcal{E})$, provided respectively by condition $(iii)(b)$ and condition (ii) .

Finally, let us show that, under the assumption of conditions (i) and (ii) of the theorem, conditions $(iii)(a)$, $(iii)(b)$, $(iii)(c)$ are all equivalent.

Conditions $(iii)(b)$ and $(iii)(c)$ are equivalent by Remark 3.13(c). The necessity of condition $(iii)(a)$ follows from Remark 4.11 and the fact that any finitely presentable model of a theory of presheaf type is finitely presented (cf. [8]).

Having already verified that conditions (i) , (ii) and $(iii)(b)$ imply all together that $\mathbb{T}$ is classified by the topos $[\mathcal{K}, \mathbf{Set}]$, and that conditions $(iii)(a)$ and $(iii)(d)$ are both necessary conditions for $\mathbb{T}$ to be of presheaf type, it remains to check that, under conditions (i) and (ii) of the theorem, condition $(iii)(a)$ implies condition $(iii)(b)$. We shall do so by verifying that conditions (i) , (ii) and $(iii)(a)$ imply all together that $\mathbb{T}$ is classified by the presheaf topos $[\mathcal{K}, \mathbf{Set}]$.

Under conditions (i) , (ii) and $(iii)(a)$, we clearly have functors

$$G_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\mathrm{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$$

and

$$H_{\mathcal{E}} : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{K}^{\mathrm{op}}, \mathcal{E})$$

which are natural in $\mathcal{E}$ and therefore induce geometric morphisms

$$G : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

and

$$H : \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}}) \rightarrow [\mathcal{K}, \mathbf{Set}] .$$

Notice that the morphism G coincides with the morphism $p_{\mathcal{K}}$ canonically induced by the universal property of the classifying topos for $\mathbb{T}$ by the $\mathbb{T}$ -models in $\mathcal{K}$.

Notice that for any $\mathbb{T}$ -model P in $\mathcal{E}$, $H_{\mathcal{E}}(P) = f_P^* \circ H^* \circ y_{\mathcal{K}}$, where $f_P : \mathcal{E} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ is the geometric morphism corresponding to P via the universal property of the classifying topos and $y_{\mathcal{K}} : \mathcal{K}^{\mathrm{op}} \rightarrow [\mathcal{K}, \mathbf{Set}]$ is the Yoneda embedding.

Under our assumptions, we have to prove that

$$H \circ G \cong 1_{[\mathcal{K}, \mathbf{Set}]}$$

or equivalently, that $G^* \circ H^* \circ y_{\mathcal{K}} \cong y_{\mathcal{K}}$.

Let us consider the geometric morphisms

$$e_N : \mathbf{Set} \rightarrow [\mathcal{K}, \mathbf{Set}]$$

corresponding to the models N of $\mathbb{T}$ in $\mathcal{K}$. As the e_N are jointly surjective, it is equivalent to prove that $e_N^* \circ G^* \circ H^* \circ y_{\mathcal{K}} \cong e_N^* \circ y_{\mathcal{K}} = \text{Hom}_{\mathcal{K}}(-, N)$ naturally in $N \in \mathcal{K}$.

Let us denote by M_G the $\mathbb{T}$ -model in $[\mathcal{K}, \mathbf{Set}]$ corresponding to the geometric morphism G . Clearly, for any $\mathbb{T}$ -model N in $\mathcal{K}$, $e_N^*(M_G) \cong N$.

We have that $G^* \circ H^* \circ y_{\mathcal{K}} = H_{[\mathcal{K}, \mathbf{Set}]}(M_G)$. But $H_{[\mathcal{K}, \mathbf{Set}]}(M_G) = \text{Hom}(\gamma_{[\mathcal{K}, \mathbf{Set}]}^*(-), M_G)$, and, by condition (iii)(a),

$$e_N^*(\text{Hom}(\gamma_{[\mathcal{K}, \mathbf{Set}]}^*(-), M_G)) \cong \text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(-, e_N^*(M_G)) \cong \text{Hom}_{\mathcal{K}}(-, N),$$

(we have omitted the subscripts and superscripts in the *Homs* above to lighten the notation), as required.

On the other hand, if condition (ii) holds then $G \circ H$ is isomorphic to the identity, so we can conclude that $\mathbb{T}$ is classified by the topos $[\mathcal{K}, \mathbf{Set}]$. In particular, $\mathbb{T}$ satisfies condition (iii)(b). This completes the proof of the theorem. $\square$

Remarks 5.2. (a) The following condition is sufficient, together with conditions (i) and (ii) of the theorem (or equivalently, together with condition (i) and the requirement that the models in $\mathcal{K}$ should be jointly conservative for $\mathbb{T}$), to ensure that $\mathbb{T}$ is classified by the topos $[\mathcal{K}, \mathbf{Set}]$ but necessary only if one assumes the axiom of choice:

(*) There is an assignment $M \rightarrow \phi_M(x_{\vec{M}})$ to a $\mathbb{T}$ -model in $\mathcal{K}$ of a formula $\phi_M(x_{\vec{M}})$ presenting it such that every $\mathbb{T}$ -model homomorphism $M \rightarrow N$ between models in $\mathcal{K}$ is induced by a $\mathbb{T}$ -provably functional formula from $\phi_N(x_{\vec{N}})$ to $\phi_M(x_{\vec{M}})$.

This condition can be easier to verify in practice than the original condition since it allows to work with distinguished presentations (rather than with *all* of them) of finitely presentable models of the theory.

The necessity of condition (*), under the axiom of choice, was established in [8]. In the converse direction, it suffices by Theorem 5.1 to prove that, under conditions (i) and (ii) (or equivalently, under condition (i) and the assertion that the models in $\mathcal{K}$ are jointly conservative for $\mathbb{T}$), condition (*) implies condition (iii)(c). First, we remark that under either of these assumptions, the canonical geometric morphism $p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$ is a surjection. From this it easily follows that for any geometric formulae $\{\vec{x} . \phi\}$ and $\{\vec{y} . \psi\}$ over Σ respectively presenting models c and d in $\mathcal{K}$, there can be at most one provably functional formula $\{\vec{x} . \phi\} \rightarrow \{\vec{y} . \psi\}$ over Σ , up to $\mathbb{T}$ -provable equivalence,

inducing a given homomorphism of $\mathbb{T}$ -models $d \rightarrow c$. Condition $(*)$ thus implies that we have a well-defined full and faithful functor $\mathcal{K} \rightarrow \mathcal{C}_{\mathbb{T}}$. From this it is immediate to see, by invoking Theorem 3.5 and Remark 3.6, that condition $(iii)(c)$ is satisfied.

- (b) If conditions (i) and (ii) in the theorem are satisfied then the canonical geometric morphism

$$p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

is a surjection; in other words, the models in $\mathcal{K}$ are jointly conservative for $\mathbb{T}$. Indeed, by condition (i) the functor H_M , where M is the universal model of $\mathbb{T}$ lying in its classifying topos, is flat. By applying Corollary 3.7 to it we thus obtain that for any geometric sequent $\sigma = (\phi \vdash_{\vec{x}} \psi)$ over the signature of $\mathbb{T}$, $\tilde{H}_M(\{\vec{x} \cdot \phi\}) \leq \tilde{H}_M(\{\vec{x} \cdot \psi\})$; condition (ii) , combined with the conservativity of M , thus allows to conclude that σ is provable in $\mathbb{T}$, as required.

- (c) Under condition (i) , if condition $(iii)(a)$ is satisfied and the models in $\mathcal{K}$ are jointly conservative for $\mathbb{T}$ (that is, every geometric sequent over Σ which is valid in every model in $\mathcal{K}$ is provable in $\mathbb{T}$) then $\mathbb{T}$ satisfies condition (ii) . Indeed, the assertion that the models in $\mathcal{K}$ should be enough for the theory $\mathbb{T}$ is precisely equivalent to the requirement that the geometric morphism $p_{\mathcal{K}}$ should be a surjection. By the naturality in $\mathcal{E}$ of the operation $(-)$ and that of the functor $H_{\mathcal{E}}$ (notice that the latter follows from condition $(iii)(a)$), it suffices to verify, since $p_{\mathcal{K}}$ is surjective, that the canonical morphism $\tilde{H}_M(M_{\mathbb{T}}) \rightarrow M$ is an isomorphism for M equal to a model in $\mathcal{K}$. But the fact that this condition holds is obvious, as required.
- (d) Conditions (ii) and $(iii)(c)$ in Theorem 5.1 admit invariant formulations, which can be profitably exploited in presence of different representations for the classifying topos of $\mathbb{T}$. Indeed, they can both be entirely reformulated in terms of the extension of flat functors operation (in the sense of section 3.1) along the canonical geometric morphism

$$p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Set}[\mathbb{T}]$$

to the classifying topos $\mathbf{Set}[\mathbb{T}]$ for $\mathbb{T}$ induced by the $\mathbb{T}$ -models in $\mathcal{K}$. Specifically, any site of definition $(\mathcal{C}, J)$ for $\mathbf{Set}[\mathbb{T}]$ gives rise to a functor

$$G_{\mathcal{E}}^{(\mathcal{C}, J)} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}_J(\mathcal{C}, \mathcal{E}).$$

Condition $(iii)(c)$ asserts that this functor, whose explicit description is given section 3.1, is full and faithful, while condition (ii) asserts that, denoting by

$$u_{\mathcal{E}} : \mathbf{Flat}_J(\mathcal{C}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

the equivalence canonically induced by the universal property of the classifying topos of $\mathbb{T}$, for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, the canonical morphism $u_{\mathcal{E}}(G_{\mathcal{E}}^{(C,J)}(H_M)) \rightarrow M$ is an isomorphism.

- (e) Condition (iii)(b) can be split into two separate conditions: η^F is pointwise monic and η^F is pointwise epic. We shall refer to the first condition as to condition (iii)(b)-(1) and to the second as to condition (iii)(b)-(2). By Theorem 3.12 and Lemma 7.11 below, condition (iii)(b)-(1) is equivalent to the requirement that the functor $u_{(\mathcal{K},\mathcal{E})}^{\mathbb{T}}$ should be faithful.

5.2 Concrete reformulations

In this section we shall give ‘concrete’ reformulations of the conditions of Theorem 5.1, in full generality as well as in some particular cases in which they admit relevant simplifications.

5.2.1 Condition (i)

In this section we shall give an explicit reformulation of condition (i) in Theorem 5.1.

Theorem 5.3. *Let $\mathbb{T}$ be a geometric theory, $\mathcal{K}$ a small category of set-based models of $\mathbb{T}$, $\mathcal{E}$ a Grothendieck topos with a separating set S and M a $\mathbb{T}$ -model in $\mathcal{E}$. Then the following conditions are equivalent:*

- (i) *The functor*

$$H_M := \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), M) : \mathcal{K}^{op} \rightarrow \mathcal{E}$$

is flat.

- (ii) (a) *There exists an epimorphic family $\{E_i \rightarrow 1_{\mathcal{E}} \mid i \in I, E_i \in S\}$ and for each $i \in I$ a $\mathbb{T}$ -model c_i in $\mathcal{K}$ and a Σ -structure homomorphism $c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$;*
- (b) *For any $\mathbb{T}$ -models c and d in $\mathcal{K}$ and Σ -structure homomorphisms $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ (where $E \in S$) and $y : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each $i \in I$ a $\mathbb{T}$ -model b_i in $\mathcal{K}$, $\mathbb{T}$ -model homomorphisms $u_i : c \rightarrow b_i$, $v_i : d \rightarrow b_i$ and a Σ -structure homomorphism $z_i : b_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $\text{Hom}_{\mathcal{E}}(e_i, M) \circ x = z_i \circ u_i$ and $\text{Hom}_{\mathcal{E}}(e_i, M) \circ y = z_i \circ v_i$.*
- (c) *For any two parallel homomorphisms $u, v : d \rightarrow c$ of $\mathbb{T}$ -models in $\mathcal{K}$ and any Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ in $\mathcal{E}$ (where $E \in S$) for which $x \circ u = x \circ v$, there is an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ in $\mathcal{E}$ and for each index i*

a homomorphism of $\mathbb{T}$ -models in $\mathcal{K}$ $w_i : c \rightarrow b_i$ and a Σ -structure homomorphism $y_i : b_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $w_i \circ u = w_i \circ v$ and $y_i \circ w_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ x$.

Proof This follows immediately from Proposition 4.5 in view of the characterization of flat functors as filtering functors established in chapter VII of [26] and reported in section 2.7. $\square$

Remarks 5.4. (a) If for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ and any object E of $\mathcal{E}$ the Σ -structure $\text{Hom}_{\mathcal{E}}(E, M)$ is a $\mathbb{T}$ -model then the three conditions (a), (b) and (c) are satisfied if they are satisfied in the case $\mathcal{E} = \mathbf{Set}$.

- (b) Condition (ii)(b) implies condition (ii)(c) if all the $\mathbb{T}$ -model homomorphisms in any Grothendieck topos are monic.
- (c) Condition (ii)(a) follows from condition (ii)(a) of Theorem 5.7 below if the signature of $\mathbb{T}$ contains at least one constant. Indeed, for any $\mathbb{T}$ -model M in a Grothendieck topos, the interpretation of such constant in M will be an arrow $1 \rightarrow MA$ in the topos, where A is the sort of the constant; applying part (a) of condition (ii) of Theorem 5.7 thus yields an epimorphic family satisfying the required property.
- (d) If all the $\mathbb{T}$ -models in $\mathcal{K}$ are finitely generated as Σ -structures and all the Σ -structure homomorphisms of the form $c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ (where c is a $\mathbb{T}$ -model in $\mathcal{K}$ and M is a $\mathbb{T}$ -model in $\mathcal{E}$) are injective if $E \not\cong 0$, a sufficient condition for property (ii)(b) to hold is that, for any context $\vec{x}$, the disjunction $(\top \vdash_{\vec{x}} \bigvee_{\phi(x) \in \mathcal{I}_{\mathcal{K}}^{\vec{x}}} \phi(\vec{x}))$ be provable in $\mathbb{T}$, where

$\mathcal{I}_{\mathcal{K}}^{\vec{x}}$ is the set of geometric formulae in the context $\vec{x}$ which strongly finitely present a $\mathbb{T}$ -model in $\mathcal{K}$ (in the sense of Definition 4.7). Indeed, for any two Σ -structure homomorphisms $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and $y : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c and d are finitely generated Σ -structures and $E \not\cong 0$, the substructure $r_e : e \hookrightarrow \text{Hom}_{\mathcal{E}}(E, M)$ of $\text{Hom}_{\mathcal{E}}(E, M)$ generated by the images of c under x and of d under y is finitely generated, say by elements $\xi_1, \dots, \xi_n$; by choosing a context $\vec{x}$ of the same length as the number of generators of e , we obtain an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S, E_i \not\cong 0\}$ with the property that for each $i \in I$ there exists a geometric formula $\phi_i(\vec{x})$ strongly presenting a $\mathbb{T}$ -model u_i in $\mathcal{K}$ such that $(\xi_1 \circ e_i, \dots, \xi_n \circ e_i)$ factors through $[[\vec{x} \cdot \phi_i]]_M$. Therefore, by the universal property of u_i as $\mathbb{T}$ -model strongly presented by the formula $\phi_i(\vec{x})$, for each $i \in I$ we have a Σ -structure homomorphism $z_i : u_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ which sends the generators of u_i to the element $(\xi_1 \circ e_i, \dots, \xi_n \circ e_i)$. Since z_i is injective by our hypothesis and its image contains a set of generators for e , we have a factorization of

$\text{Hom}_{\mathcal{E}}(e_i, M) \circ r_e$ through r_i . Therefore both $\text{Hom}_{\mathcal{E}}(e_i, M) \circ x$ and $\text{Hom}_{\mathcal{E}}(e_i, M) \circ y$ factor through z_i . This shows that condition (ii)(b) is satisfied.

Let us suppose that every $\mathbb{T}$ -model in $\mathcal{K}$ is strongly finitely presented by a formula over Σ (in the sense of Definition 4.7) and that every $\mathbb{T}$ -model homomorphism between two models in $\mathcal{K}$ is induced by a $\mathbb{T}$ -provably functional formula between formulas which present them. Then, in light of the discussion preceding Definition 4.7, we can express the conditions for the functor H_M to be flat (equivalently, filtering) in terms of the satisfaction by M of certain geometric sequents involving these formulas. Specifically, we have the following result.

Theorem 5.5. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small full category of the category of set-based $\mathbb{T}$ -models and $\mathcal{P}$ a family of geometric formulae over Σ such that every $\mathbb{T}$ -model in $\mathcal{K}$ is strongly presented by a formula in $\mathcal{P}$ and for any two formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$ presenting respectively models c and d in $\mathcal{K}$, any $\mathbb{T}$ -model homomorphism $d \rightarrow c$ is induced by a $\mathbb{T}$ -provably functional formula from $\phi(\vec{x})$ to $\psi(\vec{y})$. Then $\mathbb{T}$ satisfies condition (i) of Theorem 5.1 with respect to the category $\mathcal{K}$ if and only if the following conditions are satisfied:*

(i) *The sequent*

$$(\top \vdash_{\square} \bigvee_{\phi(\vec{x}) \in \mathcal{P}} (\exists \vec{x}) \phi(\vec{x}))$$

is valid in M ;

(ii) *For any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, the sequent*

$$\phi(\vec{x}) \wedge \psi(\vec{y}) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, \\ \{\vec{x}, \phi\} \xrightarrow{\theta_1} \{\vec{z}, \chi\} \xrightarrow{\theta_2} \{\vec{y}, \psi\} \text{ in } \mathcal{C}_{\mathbb{T}}}} (\exists \vec{z}) (\theta_1(\vec{z}, \vec{x}) \wedge \theta_2(\vec{z}, \vec{y})),$$

is valid in M ;

(iii) *For any $\mathbb{T}$ -provably functional formulae $\theta_1, \theta_2 : \phi(\vec{x}) \rightarrow \psi(\vec{y})$ between two formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, the sequent*

$$\theta_1(\vec{x}, \vec{y}) \wedge \theta_2(\vec{x}, \vec{y}) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, \\ \{\vec{z}, \chi\} \xrightarrow{\tau} \{\vec{x}, \phi\} \text{ in } \mathcal{C}_{\mathbb{T}}, \\ \tau \wedge \theta_1 \dashv \vdash_{\mathbb{T}} \tau \wedge \theta_2}} (\exists \vec{z}) \tau(\vec{z}, \vec{x})$$

is valid in M .

Proof The fact that every model c in $\mathcal{K}$ is strongly finitely presented by a formula $\phi(\vec{x})$ in $\mathcal{P}$ ensures that for any object E of $\mathcal{E}$ and any $\mathbb{T}$ -model M in $\mathcal{E}$

the Σ -homomorphism $c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ are in bijective correspondence with the generalized elements $E \rightarrow [[\vec{x} \cdot \phi]]_M$ in $\mathcal{E}$. The thesis then follows from the Kripke-Joyal semantics for the topos $\mathcal{E}$, noticing that by our hypotheses for any two formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$ presenting respectively models c and d in $\mathcal{K}$, any $\mathbb{T}$ -model homomorphism $d \rightarrow c$ is induced by a $\mathbb{T}$ -provably functional formula from $\phi(\vec{x})$ to $\psi(\vec{y})$. $\square$

Remarks 5.6. (a) For any geometric theory $\mathbb{T}$ and category $\mathcal{K}$ satisfying the hypotheses of the theorem, all the sequents in the statement of the theorem are satisfied by every model in $\mathcal{K}$. Therefore, adding them to $\mathbb{T}$ yields a quotient of $\mathbb{T}$ satisfying condition (i) of Theorem 5.1 with respect to the category $\mathcal{K}$.

(b) Under the alternative hypothesis that every model of $\mathcal{K}$ is both strongly finitely presentable and finitely generated (with respect to the same generators), for any two formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$ presenting respectively models c and d in $\mathcal{K}$, the $\mathbb{T}$ -model homomorphisms $c \rightarrow d$ are in bijection (via the evaluation of such homomorphisms at the generators of c) with the tuples of elements of d which satisfy ϕ , each of which has the form $(t_1(\vec{z}), \dots, t_n(\vec{z}))$ (where n is the length of the context $\vec{x}$) for some terms $t_1, \dots, t_n$ in the context $\vec{z}$ over the signature Σ . Conditions (ii) and (iii) of Theorem 5.5 can thus be reformulated more explicitly as follows:

(ii') For any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, the sequent

$$(\phi(\vec{x}) \wedge \psi(\vec{y})) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, t_1^{A_1}(\vec{z}), \dots, t_n^{A_n}(\vec{z}) \\ s_1^{B_1}(\vec{z}), \dots, s_m^{B_m}(\vec{z})}} (\exists \vec{z}) (\chi(\vec{z}) \wedge \bigwedge_{\substack{i \in \{1, \dots, n\}, \\ j \in \{1, \dots, m\}}} (x_i = t_i(\vec{z}) \wedge y_j = s_j(\vec{z}))),$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $t_1^{A_1}(\vec{z}), \dots, t_n^{A_n}(\vec{z})$ and $s_1^{B_1}(\vec{z}), \dots, s_m^{B_m}(\vec{z})$ whose output sorts are respectively $A_1, \dots, A_n, B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(t_1^{A_1}(\vec{\xi}), \dots, t_n^{A_n}(\vec{\xi})) \in [[\vec{x} \cdot \phi]]_{M_{\{\vec{z}, \chi\}}}$ and $(s_1^{B_1}(\vec{\xi}), \dots, s_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$, is valid in M ;

(iii') For any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, and any terms $t_1^{A_1}(\vec{y}), s_1^{A_1}(\vec{y}), \dots, t_n^{A_n}(\vec{y}), s_n^{A_n}(\vec{y})$ whose output sorts are respectively $A_1, \dots, A_n$, the sequent

$$\begin{aligned} & \left(\bigwedge_{i \in \{1, \dots, n\}} (t_i(\vec{y}) = s_i(\vec{y})) \right) \wedge \phi(t_1/x_1, \dots, t_n/x_n) \wedge \phi(s_1/x_1, \dots, s_n/x_n) \wedge \psi(\vec{y}) \\ & \vdash_{\vec{y}} \bigvee_{\chi(\vec{z}) \in \mathcal{P}, u_1^{B_1}(\vec{z}), \dots, u_m^{B_m}(\vec{z})} ((\exists \vec{z}) (\chi(\vec{z}) \wedge \bigwedge_{j \in \{1, \dots, m\}} (y_j = u_j(\vec{z}))), \end{aligned}$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $u_1^{B_1}(\vec{z}), \dots, u_m^{B_m}(\vec{z})$ whose output sorts are respectively $B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(u_1^{B_1}(\vec{\xi}), \dots, u_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$ and $t_i(u_1(\vec{\xi}), \dots, u_m(\vec{\xi})) = s_i(u_1(\vec{\xi}), \dots, u_m(\vec{\xi}))$ in $M_{\{\vec{z}, \chi\}}$ for all $i \in \{1, \dots, n\}$, is valid in M .

5.2.2 Condition (ii)

In this section we shall give concrete reformulations of condition (ii) of Theorem 5.1, under the assumption that condition (i) of Theorem 5.1 is satisfied.

First, notice that the canonical morphism $\tilde{H}_M(M_{\mathbb{T}}) \rightarrow M$ considered in condition (ii) is an isomorphism if and only if for every sort A over Σ , the induced arrow $\tilde{H}_M(\{x^A \cdot \top\}) \rightarrow MA$ is an isomorphism in $\mathcal{E}$. To understand this condition more concretely, we apply Proposition 3.8 to the geometric formula $\{x^A \cdot \top\}$ and the flat functor $F = H_M : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$. Recall that the category $\mathcal{A}_{\{x^A, \top\}}^{\mathcal{K}}$ defined in that context has as objects the pairs (c, z) where c is a $\mathbb{T}$ -model in $\mathcal{K}$ and z is an element of the set cA and as arrows $(c, z) \rightarrow (d, w)$ the $\mathbb{T}$ -model homomorphisms $g : d \rightarrow c$ in $\mathcal{K}$ such that $gA(w) = z$, and that the equivalence relation $R_{\{x^A, \top\}}$ is defined by the condition that for any generalized elements $x : E \rightarrow F(c)$ and $x' : E \rightarrow F(d)$, $(x, x') \in R_{\{x^A, \top\}}$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(b_i)$ and two $\mathbb{T}$ -model homomorphisms $f_i : c \rightarrow b_i$ and $f'_i : d \rightarrow b_i$ in $\mathcal{K}$ such that $f_i A(z) = f'_i A(w)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$.

This yields that the canonical arrow $\tilde{H}_M(\{x^A \cdot \top\}) \rightarrow MA$ is an isomorphism if and only if (using the notation of Proposition 3.8):

- (1) the canonical arrows $\kappa_{(c, z)} : H_M(c) \rightarrow MA$ for $(c, z) \in \mathcal{A}_{\{x^A, \top\}}$ are jointly epimorphic and
- (2) for any two objects (c, z) and (d, w) of the category $\mathcal{A}_{\{x^A, \top\}}$ and any generalized elements $x : E \rightarrow H_M(c)$ and $x' : E \rightarrow H_M(d)$ (where $E \in \mathcal{S}$), $\kappa_{(c, z)} \circ x = \kappa_{(d, w)} \circ x'$ if and only if $(x, x') \in R_{\{x^A, \top\}}$.

Thanks to the identification between the generalized elements $x : E \rightarrow H_M(c)$ and the Σ -structure homomorphisms $f_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ provided by Proposition 4.5, we can rewrite conditions (1) and (2) more explicitly.

To this end, we preliminarily notice that for any object (c, z) of the category $\mathcal{A}_{\{x^A, \top\}}$, the canonical arrow $\kappa_{(c, z)} : H_M(c) = \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}}(\gamma_{\mathcal{E}}^*(c), M) \rightarrow MA$ can be described in terms of generalized elements as the arrow sending any generalized element $x : E \rightarrow H_M(c)$, corresponding via the identification of Proposition 4.5 to a Σ -structure homomorphism $f_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, to the generalized element $E \rightarrow MA$ of MA given by $f_x A(z)$. Therefore, for

any generalized elements $x : E \rightarrow H_M(c)$ and $x' : E \rightarrow H_M(d)$, corresponding respectively to Σ -structure homomorphisms $f_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and $f_{x'} : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, we have that $\kappa_{(c,z)} \circ x = \kappa_{(d,w)} \circ x'$ if and only if $f_x A(z) = f_{x'} A(w)$.

Now, condition (1) can be formulated by saying that for any generalized element $x : E \rightarrow MA$ (where $E \in S$) there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$, a family $\{(c_i, z_i) \mid i \in I\}$ of objects of the category $\mathcal{A}_{\{x^A, \top\}}$ and generalized elements $\{y_i : E_i \rightarrow H_M(c_i) \mid i \in I\}$ such that $x \circ e_i = \xi_{(c_i, z_i)} \circ y_i$ for every $i \in I$.

Under the identification of Proposition 4.5, condition (1) thus rewrites as follows: for any generalized element $x : E \rightarrow MA$ (where $E \in S$) there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$, a family $\{(c_i, z_i) \mid i \in I\}$ of objects of the category $\mathcal{A}_{\{x^A, \top\}}$ and Σ -structure homomorphisms $f_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $f_i A(z_i) = x \circ e_i$.

Concerning condition (2), we observe that for any objects (c, z) and (d, w) of the category $\mathcal{A}_{\{x^A, \top\}}$ and any Σ -structure homomorphisms $f_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and $f_{x'} : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ (where $E \in S$) corresponding respectively to generalized elements $x : E \rightarrow H_M(c)$ and $x' : E \rightarrow H_M(d)$ via the identification of Proposition 4.5, we have that $(x, x') \in R_{\{x^A, \top\}}$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a Σ -structure homomorphism $h_i : b_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ and two $\mathbb{T}$ -model homomorphisms $f_i : c \rightarrow b_i$ and $f'_i : d \rightarrow b_i$ in $\mathcal{K}$ such that $f_i A(z) = f'_i A(w)$ and $h_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ f_x$ and $h_i \circ f'_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ f_{x'}$.

Summarizing, we have the following result.

Theorem 5.7. *Let $\mathbb{T}$ be a geometric theory, $\mathcal{K}$ a small full subcategory of the category of set-based models of $\mathbb{T}$, $\mathcal{E}$ a Grothendieck topos with a separating set S and M a $\mathbb{T}$ -model in $\mathcal{E}$. Then the following conditions are equivalent:*

- (i) *The extension $\tilde{H}_M : \mathcal{C}_{\mathbb{T}} \rightarrow \mathcal{E}$ of the functor $H_M : \mathcal{K}^{op} \rightarrow \mathcal{E}$ to the syntactic category $\mathcal{C}_{\mathbb{T}}$ (in the sense of section 3.3) satisfies the property that the canonical morphism $\tilde{H}_M(M_{\mathbb{T}}) \rightarrow M$ is an isomorphism;*
- (ii) *For any sort A over Σ , the following conditions are satisfied:*
 - (a) *For any generalized element $x : E \rightarrow MA$ (where $E \in S$) there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a $\mathbb{T}$ -model c_i in $\mathcal{K}$, an element z_i of $c_i A$ and a Σ -structure homomorphism $f_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $(f_i A)(z_i) = x \circ e_i$;*
 - (b) *For any two pairs (c, z) and (d, w) consisting of $\mathbb{T}$ -models c and d in $\mathcal{K}$ and elements $z \in cA, w \in dA$, and any Σ -structure homomorphisms $f : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and $f' : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ (where E is an object of S), we have that $f A(z) = f' A(w)$ if and only if there*

exists an epimorphic family $\{e_j : E_j \rightarrow E \mid j \in J, E_j \in S\}$ and for each index $j \in J$ a $\mathbb{T}$ -model b_j in $\mathcal{K}$, a Σ -structure homomorphism $h_j : b_j \rightarrow \text{Hom}_{\mathcal{E}}(E_j, M)$ and two $\mathbb{T}$ -model homomorphisms $f_j : c \rightarrow b_j$ and $f'_j : d \rightarrow b_j$ in $\mathcal{K}$ such that $f_j A(z) = f_j A'(w)$, $h_j \circ f_j = \text{Hom}_{\mathcal{E}}(e_j, M) \circ f$ and $h_j \circ f'_j = \text{Hom}_{\mathcal{E}}(e_j, M) \circ f'$.

Remarks 5.8. (a) If all the $\mathbb{T}$ -model homomorphisms in any Grothendieck topos are monic and $\mathbb{T}$ satisfies condition (ii)(b) of Theorem 5.3 then condition (ii)(b) of Theorem 5.7 is automatically satisfied.

- (b) A sufficient condition for condition (ii)(a) to hold is that the disjunction $(\top \vdash_x \bigvee_{\phi(x) \in \mathcal{I}_{\mathcal{K}}^x} \phi(x))$ be provable in $\mathbb{T}$, where $\mathcal{I}_{\mathcal{K}}^x$ is the set of geometric formulae in one variable which strongly finitely present a $\mathbb{T}$ -model in $\mathcal{K}$ (in the sense of Definition 4.7).
- (c) If for every sort A over Σ the formula $\{x^A \cdot \top\}$ strongly presents a $\mathbb{T}$ -model F_A in $\mathcal{K}$ then condition (i) of the theorem is automatically satisfied; indeed, under this hypothesis for any sort A over Σ the canonical arrow $\tilde{H}_M(M_{\mathbb{T}})A = \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}}(\gamma_{\mathcal{E}}^*(F_A), M) \rightarrow MA$ is an isomorphism (cf. section 4.3).

The following result provides an explicit formulation of condition (ii) of Theorem 5.1 holding for theories $\mathbb{T}$ with respect to small categories $\mathcal{K}$ such that every model of $\mathcal{K}$ is both strongly finitely presentable and finitely generated (with respect to the same generators).

Theorem 5.9. *Let $\mathbb{T}$ be a geometric theory over a signature Σ and $\mathcal{K}$ a small full subcategory of the category of set-based $\mathbb{T}$ -models such that every model in $\mathcal{K}$ is both strongly finitely presentable and finitely generated (with respect to the same generators). Then $\mathbb{T}$ satisfies condition (ii) of Theorem 5.1 with respect to the category $\mathcal{K}$ if and only if for every model M of $\mathbb{T}$ in a Grothendieck topos, the following conditions are satisfied (where $\mathcal{P}$ denotes the set of formulae over Σ which present a model in $\mathcal{K}$):*

- (i) *For any sort A over Σ , the sequent*

$$(\top \vdash_{x_A} \bigvee_{\chi(\vec{z}) \in \mathcal{P}, t^A(\vec{z})} (\exists \vec{z})(\chi(\vec{z}) \wedge x = t(\vec{z}))),$$

where the the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the terms $t^A(\vec{z})$ whose output sort is A ;

- (ii) *For any sort A over Σ , any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, and any terms $t^A(\vec{x})$ and $s^A(\vec{y})$, the sequent*

$$\begin{aligned}
& (\phi(\vec{x}) \wedge \psi(\vec{y}) \wedge t(\vec{x}) = s(\vec{y})) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, p_1^{A_1}(\vec{z}), \dots, p_n^{A_n}(\vec{z}) \\ q_1^{B_1}(\vec{z}), \dots, q_m^{B_m}(\vec{z})}} (\exists \vec{z})(\chi(\vec{z}) \wedge \\
& \wedge \bigwedge_{\substack{i \in \{1, \dots, n\}, \\ j \in \{1, \dots, m\}}} (x_i = p_i(\vec{z}) \wedge y_j = q_j(\vec{z}))),
\end{aligned}$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $p_1^{A_1}(\vec{z}), \dots, p_n^{A_n}(\vec{z})$ and $q_1^{B_1}(\vec{z}), \dots, q_m^{B_m}(\vec{z})$ whose output sorts are respectively $A_1, \dots, A_n, B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(p_1^{A_1}(\vec{\xi}), \dots, p_n^{A_n}(\vec{\xi})) \in [[\vec{x} \cdot \phi]]_{M_{\{\vec{z}, \chi\}}}$ and $(q_1^{B_1}(\vec{\xi}), \dots, q_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$ and $t(p_1(\vec{\xi}), \dots, p_n(\vec{\xi})) = s(q_1(\vec{\xi}), \dots, q_m(\vec{\xi}))$ in $M_{\{\vec{z}, \chi\}}$.

Proof The proof is analogous to that of Theorem 5.3 and left to the reader. $\square$

Remark 5.10. All the sequents in the statement of the theorem are satisfied by every model in $\mathcal{K}$. Therefore, adding them to any theory $\mathbb{T}$ satisfying the hypotheses of the theorem yields a quotient of $\mathbb{T}$ satisfying condition (ii) of Theorem 5.1 with respect to the category $\mathcal{K}$.

5.2.3 Condition (iii)

In this section, we shall give concrete reformulations of conditions (iii)(a)-(b)-(c) of Theorem 5.1.

Let us begin our analysis with a proposition which shows that, under some natural assumptions which are often verified in practice, condition (iii)(a) of Theorem 5.1 is satisfied.

Proposition 5.11. *Let $\mathbb{T}$ be a geometric theory over a signature Σ and let $\mathcal{K}$ be a small category of $\mathbb{T}$ -mod(**Set**). Then*

- (i) *If $\mathbb{T}$ is a quotient of a theory $\mathbb{S}$ satisfying condition (iii)(a) of Theorem 5.1 with respect to a category $\mathcal{H}$ of set-based $\mathbb{S}$ -models and $\mathcal{K}$ is a subcategory of $\mathcal{H}$ then $\mathbb{T}$ satisfies property (iii)(a) of Theorem 5.1 with respect to $\mathcal{K}$;*
- (ii) *If every $\mathbb{T}$ -model in $\mathcal{K}$ is strongly finitely presented (in the sense of section 4.3) then $\mathbb{T}$ satisfies condition (iii)(a) of Theorem 5.1 with respect to the category $\mathcal{K}$;*

(iii) If for every $\mathbb{T}$ -model c in $\mathcal{K}$ the object $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ can be built from c and M by only using geometric constructions (i.e. constructions only involving finite limits and arbitrary small colimits) then $\mathbb{T}$ satisfies condition (iii)(a) of Theorem 5.1 with respect to the category $\mathcal{K}$.

Proof If $\mathbb{T}$ is a quotient of $\mathbb{S}$ then for every Grothendieck topos $\mathcal{E}$, the category $\mathbb{T}\text{-mod}(\mathcal{E})$ is a full subcategory of the category $\mathbb{S}\text{-mod}(\mathcal{E})$. This clearly implies that for any $\mathbb{T}$ -models M and N in a Grothendieck topos $\mathcal{E}$, $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N) \cong \text{Hom}_{\mathbb{S}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)$; in particular, for any $\mathbb{T}$ -model c in $\mathcal{K}$ and any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M) \cong \text{Hom}_{\mathbb{S}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$. The fact that $\mathbb{S}$ satisfies condition (iii)(a) of Theorem 5.1 with respect to the category $\mathcal{H}$ thus implies that $\mathbb{T}$ does with respect to the category $\mathcal{K}$, as required.

If every $\mathbb{T}$ -model c in $\mathcal{K}$ is strongly finitely presented (in the sense of section 4.3) by a formula $\phi(\vec{x})$ over the signature of $\mathbb{T}$ then for any $\mathbb{T}$ -model M in a Grothendieck topos, $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M) \cong [[\vec{x} \cdot \phi]]_M$ (cf. section 4.3). Therefore, as the interpretation of geometric formulae is always preserved by inverse image functors of geometric morphisms, condition (iii)(a) of Theorem 5.1 is satisfied by the theory $\mathbb{T}$ with respect to the category $\mathcal{K}$.

The fact that condition (iii) implies condition (iii)(a) of Theorem 5.1 follows immediately from the fact that geometric constructions are preserved by inverse image functors of geometric morphisms. $\square$

We shall now proceed to giving concrete reformulations of the conditions (iii)(b)-(1) and (iii)(b)-(2) of Theorem 5.1 introduced in Remark 5.2(e), in order to make them more easily verifiable in practice.

First, let us explicitly describe, for any object c of $\mathcal{K}$, the arrow

$$\eta^F(c) : F(c) \rightarrow \text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), \tilde{F}(M_{\mathbb{T}}))$$

of condition (iii)(b) of Theorem 5.1 in terms of generalized elements.

For any generalized element $x : E \rightarrow F(c)$, $\eta^F(c)(x)$ corresponds under the identification of Proposition 4.5 to the Σ -structure homomorphism $z_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ defined at each sort A over Σ as the function $cA \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(\{x^A \cdot \top\}))$ sending any element $y \in cA$ to the generalized element $E \rightarrow \tilde{F}(\{x^A \cdot \top\})$ obtained by composing the canonical colimit arrow $\kappa_{(c,y)}^F : F(c) \rightarrow \tilde{F}(\{x^A \cdot \top\})$ with the generalized element $x : E \rightarrow F(c)$.

It follows that $\eta^F(c)(x)$ is a monomorphism if and only if for any generalized elements $x, x' : E \rightarrow F(c)$, $\kappa_{(c,y)}^F \circ x = \kappa_{(c,y)}^F \circ x'$ for every sort A over Σ and element $y \in cA$ implies $x = x'$. By Proposition 3.8, the condition ' $\kappa_{(c,y)}^F \circ x = \kappa_{(c,y)}^F \circ x'$ ' is satisfied if and only if there exists an

epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and two $\mathbb{T}$ -model homomorphisms $f_i, f'_i : c \rightarrow a_i$ in $\mathcal{K}$ such that $f_i A(y) = f'_i A(y)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$.

To obtain an explicit characterization of the condition for $\eta^F(c)(x)$ to be an epimorphism, we notice that an arrow $f : A \rightarrow B$ in a Grothendieck topos $\mathcal{E}$ is an epimorphism if and only if for every generalized element $x : E \rightarrow B$ of B there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a generalized element $y_i : E_i \rightarrow A$ such that $f \circ y_i = x \circ e_i$. Applying this characterization to the arrow $\eta^F(c)(x)$ modulo the identification of Proposition 4.5, we obtain the following criterion: $\eta^F(c)(x)$ is an epimorphism if and only if for every object E of $\mathcal{E}$ and any Σ -structure homomorphism $v : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a generalized element $x_i : E_i \rightarrow F(c)$ such that $z_{x_i} = \text{Hom}_{\mathcal{E}}(e_i, \tilde{F}(M_{\mathbb{T}})) \circ v$ for all i . The condition ' $z_{x_i} = \text{Hom}_{\mathcal{E}}(e_i, \tilde{F}(M_{\mathbb{T}})) \circ v$ ' can be explicitly reformulated as the requirement that for every sort A over Σ and every element $y \in cA$, $\kappa_{(c,y)}^F \circ x_i = vA(y) \circ e_i$.

Summarizing, we have the following

Theorem 5.12. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ be a small subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$, $\mathcal{E}$ a Grothendieck topos with a separating set S and $F : \mathcal{K}^{op} \rightarrow \mathcal{E}$ a flat functor. Then*

- (i) *F satisfies condition (iii)(b)-(1) of Theorem 5.1 if and only if for any $\mathbb{T}$ -model c in $\mathcal{K}$ and any generalized elements $x, x' : E \rightarrow F(c)$ (where $E \in S$), if for every sort A over Σ and any element $y \in cA$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and two $\mathbb{T}$ -model homomorphisms $f_i, f'_i : c \rightarrow a_i$ in $\mathcal{K}$ such that $f_i A(y) = f'_i A(y)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$ then $x = x'$.*
- (ii) *F satisfies condition (iii)(b)-(2) of Theorem 5.1 if and only if for any $\mathbb{T}$ -model c in $\mathcal{K}$, any object E of S and any Σ -structure homomorphism $v : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I, E_i \in S\}$ and for each index $i \in I$ a generalized element $x_i : E_i \rightarrow F(c)$ such that for every sort A over Σ and any element $y \in cA$, $\kappa_{(c,y)}^F \circ x_i = vA(y) \circ e_i$.*

□

Under the hypothesis that for any sort A over Σ the formula $\{x^A \cdot \top\}$ presents a $\mathbb{T}$ -model F_A , the model $\tilde{F}(M_{\mathbb{T}})$ is isomorphic to the model interpreting each sort A with the object $F(F_A)$, and the homomorphism $z_x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ corresponding to a generalized element $x : E \rightarrow F(c)$ can be described as follows: for any sort A over Σ , $z_x A : cA \rightarrow \text{Hom}_{\mathcal{E}}(E, F(P_A))$

assigns to any element $y \in cA$, corresponding to a $\mathbb{T}$ -model homomorphism $s_y : F_A \rightarrow c$ via the universal property of F_A , the generalized element $F(s_y) \circ x$. Therefore, given two generalized elements $x, x' : E \rightarrow F(c)$ and an element $y \in cA$, $z_x A(y) = z_{x'} A(y)$ if and only if $F(s_y) \circ x = F(s_y) \circ x'$.

Summarizing, we have the following

Theorem 5.13. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , such that for any sort A over Σ the formula $\{x^A . \top\}$ presents a $\mathbb{T}$ -model F_A , $\mathcal{K}$ be a small subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ containing the models F_A and $F : \mathcal{K}^{op} \rightarrow \mathcal{E}$ a flat functor with values in a Grothendieck topos $\mathcal{E}$. Then F satisfies condition (iii)(b)-(1) of Theorem 5.1 with respect to the category $\mathcal{K}$ if and only if for every $\mathbb{T}$ -model c in $\mathcal{K}$ the family of arrows $\{F(s_y) : F(c) \rightarrow F(F_A) \mid A \in \Sigma, y \in cA\}$ is jointly monic.*

□

The representation of $\tilde{F}(M_{\mathbb{T}})$ as a filtered $\mathcal{E}$ -indexed colimit established in section 3.3 allows us to obtain a different reformulation, in terms of the Σ -structure homomorphisms $\xi_{(c,x)}$ defined in that context, of Theorem 5.12:

Theorem 5.14. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ and $F : \mathcal{K}^{op} \rightarrow \mathcal{E}$ be a flat functor with values in a Grothendieck topos $\mathcal{E}$. Then*

- (i) *F satisfies condition (iii)(b)-(1) of Theorem 5.1 if and only if for any $\mathbb{T}$ -model c in $\mathcal{K}$ and any generalized elements $x, x' : E \rightarrow F(c)$, the Σ -structure homomorphisms $\xi_{(c,x)}$ and $\xi_{(c,x')}$ are equal if and only if $x = x'$.*
- (ii) *F satisfies condition (iii)(b)-(2) of Theorem 5.1 if and only if for any $\mathbb{T}$ -model c in $\mathcal{K}$, any object E of $\mathcal{E}$ and any Σ -structure homomorphism $z : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a generalized element $x_i : E_i \rightarrow F(c)$ such that $\text{Hom}_{\mathcal{E}}(e_i, \tilde{F}(M_{\mathbb{T}})) \circ z = \xi_{(c,x_i)}$ for all $i \in I$.*

Proof (i) The equality $\xi_{(c,x)} = \xi_{(c,x')}$ holds if and only if for every sort A over Σ and any $y \in cA$, $\xi_{(c,x)} A(y) = \xi_{(c,x')} A(y)$, i.e. $\kappa_{(c,y)} \circ x = \kappa_{(c,y)} \circ x'$. Now, by Proposition 3.8, this latter condition is satisfied if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and two $\mathbb{T}$ -model homomorphisms $f_i, f'_i : c \rightarrow a_i$ such that $f_i A(y) = f'_i A(y)$ and $\langle F(f_i), F(f'_i) \rangle \circ h_i = \langle x, x' \rangle \circ e_i$, as required.

(ii) It suffices to notice that the condition that for every sort A over Σ and every element $y \in cA$, $\kappa_{(c,y)}^F \circ x_i = z A(y) \circ e_i$ can be reformulated as the requirement that $\text{Hom}_{\mathcal{E}}(e_i, \tilde{F}(M_{\mathbb{T}})) \circ z = \xi_{(c,x_i)}$ (for any $i \in I$). □

Thanks to this different reformulation of condition (iii)(b)-(1) of Theorem 5.1, we can prove that if all the arrows in $\mathcal{C}$ are sortwise monic then $\mathbb{T}$ always satisfies the condition.

First, we need a lemma.

Lemma 5.15. *Under the hypotheses specified above, if all the homomorphisms in the category $\mathcal{K}$ are sortwise monic then for any pair (c, x) consisting of an object c of $\mathcal{K}$ and of a generalized element $x : E \rightarrow F(c)$ such that $E \not\cong 0_{\mathcal{E}}$, the Σ -structure homomorphism $\xi_{(c,x)} : a \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ is sortwise injective.*

Proof For any sort A over Σ , the function $\xi_{(c,x)}A : cA \rightarrow \text{Hom}_{\mathcal{E}}(E, MA)$ sends any element $y \in cA$ to the generalized element $\kappa_{(c,y)} \circ x$. Now, for any $y_1, y_2 \in cA$, we have, by Proposition 3.8, that $\kappa_{(c,y_1)} \circ x = \kappa_{(c,y_2)} \circ x$ if and only if there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model a_i in $\mathcal{K}$, a generalized element $h_i : E_i \rightarrow F(a_i)$ and a $\mathbb{T}$ -model homomorphism $f_i : c \rightarrow a_i$ in $\mathcal{K}$ such that $f_iA(y_1) = f_iA(y_2)$ and $F(f_i) \circ h_i = x \circ e_i$. If $E \not\cong 0_{\mathcal{E}}$ then the set I is non-empty, that is there exists $i \in I$; thus we have that $f_iA(y_1) = f_iA(y_2)$, which entails $y_1 = y_2$ as f_i is sortwise monic by our hypothesis. $\square$

Corollary 5.16. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{K}$ a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ whose arrows are all sortwise monic homomorphisms. Then $\mathbb{T}$ satisfies condition (iii)(b)-(1) of Theorem 5.1.*

Proof By Proposition 3.11, for any $x : E \rightarrow F(c)$ and $x' : E \rightarrow F(c)$, the following ‘joint embedding property’ holds: there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a $\mathbb{T}$ -model c_i in $\mathcal{K}$, homomorphisms $f_i : c \rightarrow c_i$, $g_i : c \rightarrow c_i$ in $\mathcal{K}$ and a generalized element $x_i : E_i \rightarrow F(c_i)$ such that $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$, $\text{Hom}_{\mathcal{E}}(E_i, M) \circ \xi_{(c,x)} = \xi_{(c_i,x_i)} \circ f_i$ and $\text{Hom}_{\mathcal{E}}(E_i, M) \circ \xi_{(c,x')} = \xi_{(c_i,x_i)} \circ g_i$ (for all $i \in I$). Clearly, we can suppose without loss of generality all the objects E_i to be non-zero.

Now, since all the arrows in $\mathcal{K}$ are sortwise monic homomorphisms, by Lemma 5.15 for each $i \in I$ the arrow $\xi_{(c_i,x_i)}$ is sortwise monic and hence $\xi_{(c,x)} = \xi_{(c,x')}$ implies $f_i = g_i$. Therefore $x \circ e_i = x' \circ e_i$ for all $i \in I$ (since $\langle x, x' \rangle \circ e_i = \langle F(f_i), F(g_i) \rangle \circ x_i$), and hence $x = x'$, as required. $\square$

We shall now proceed to identifying some natural sufficient conditions for a theory $\mathbb{T}$ to satisfy condition (iii)(b)-(2) of Theorem 5.1. Before stating the relevant theorem, we need a number of preliminary results.

Lemma 5.17. *Let Σ be a signature without relation symbols and $\mathbb{T}$ a geometric theory over a signature Σ' obtained from Σ by solely adding relation symbols whose interpretation in any $\mathbb{T}$ -model in a Grothendieck topos is the*

complement of the interpretation of a geometric formula over Σ (for instance, the injectivization of a geometric theory over Σ in the sense of Definition 6.33). Let $f : M \rightarrow N$ and $g : P \rightarrow N$ be homomorphisms of $\mathbb{T}$ -models in a Grothendieck topos $\mathcal{E}$, and let k be an assignment to any sort A over Σ of an arrow $kA : MA \rightarrow PA$ such that $gA \circ kA = fA$. Then, if g is sortwise monic, k defines a $\mathbb{T}$ -model homomorphism $M \rightarrow P$ such that $g \circ k = f$.

Proof The fact that k preserves the interpretation of function symbols over Σ' follows from the fact that f and g do, as g is sortwise monic. It remains to prove that for any relation symbol $R \mapsto A_1, \dots, A_n$ over Σ' and any generalized element $x : E \rightarrow MA_1 \times \dots \times MA_n$ in $\mathcal{E}$, if x factors through $R_M \mapsto MA_1 \times \dots \times MA_n$ then $(kA_1 \times \dots \times kA_n) \circ x$ factors through $R_P \mapsto PA_1 \times \dots \times PA_n$. Now, if R_P is the complement of the interpretation $i_\phi : [[\phi(\vec{x})]]_P \mapsto PA_1 \times \dots \times PA_n$ of a geometric formula $\phi(\vec{x}) = \phi(x^{A_1}, \dots, x^{A_n})$ in the model P , this latter condition is equivalent to the requirement that the equalizer e of $(kA_1 \times \dots \times kA_n) \circ x$ and i_ϕ be zero; but, since g is a Σ' -structure homomorphism, the subobject e is contained in the equalizer of the arrows $(gA_1 \times \dots \times gA_n) \circ (kA_1 \times \dots \times kA_n) \circ x = (fA_1 \times \dots \times fA_n) \circ x$ and $[[\phi(\vec{x})]]_N \mapsto NA_1 \times \dots \times NA_n$, which is zero since f is a Σ' -structure homomorphism and R_N is the complement of $[[\phi(\vec{x})]]_N$. Therefore $e \cong 0_{\mathcal{E}}$, as required. $\square$

Lemma 5.18. *Let $\mathcal{E}$ be a Grothendieck topos. Then the inverse image functor $\gamma_{\mathcal{E}}^* : \mathbf{Set} \rightarrow \mathcal{E}$ of the unique geometric morphism $\gamma_{\mathcal{E}} : \mathcal{E} \rightarrow \mathbf{Set}$ is faithful if and only if $\mathcal{E}$ is non-trivial (i.e., $1_{\mathcal{E}} \not\cong 0_{\mathcal{E}}$).*

Proof It is clear that if $\mathcal{E}$ is trivial then $\gamma_{\mathcal{E}}^* : \mathbf{Set} \rightarrow \mathcal{E}$ is not faithful, it being the constant functor with value $0_{\mathcal{E}}$.

In the converse direction, suppose that $\mathcal{E}$ is non-trivial. Given two functions $f, g : A \rightarrow B$ in $\mathbf{Set}$, the arrows $\gamma_{\mathcal{E}}^*(f), \gamma_{\mathcal{E}}^*(g) : \coprod_{a \in A} 1_{\mathcal{E}} \rightarrow \coprod_{b \in B} 1_{\mathcal{E}}$ in $\mathcal{E}$ are characterized by the following identities: for any $a \in A$ $\gamma_{\mathcal{E}}^*(f) \circ s_a = t_{f(a)}$ and $\gamma_{\mathcal{E}}^*(g) \circ s_a = t_{g(a)}$, where $s_a : 1 \rightarrow \coprod_{a \in A} 1_{\mathcal{E}}$ and $t_b : 1 \rightarrow \coprod_{b \in B} 1_{\mathcal{E}}$ are respectively the a -th and b -th coproduct arrows (for any $a \in A$ and $b \in B$); so $\gamma_{\mathcal{E}}^*(f) = \gamma_{\mathcal{E}}^*(g)$ if and only if for every $a \in A$, $t_{f(a)} = t_{g(a)}$. Now, since a subobject with domain $1_{\mathcal{E}}$ in a non-trivial topos $\mathcal{E}$ cannot be disjoint from itself, it follows that $f(a) = g(a)$ for every $a \in A$. Therefore $f = g$, as required. $\square$

Corollary 5.19. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , $\mathcal{E}$ a non-trivial Grothendieck topos (i.e., $1_{\mathcal{E}} \not\cong 0_{\mathcal{E}}$), M and N two $\mathbb{T}$ -models in $\mathbf{Set}$ and f a function sending each sort A over Σ to a map $fA : MA \rightarrow NA$ in such a way that the assignment $A \mapsto \gamma_{\mathcal{E}}^*(fA) : \gamma_{\mathcal{E}}^*(MA) \rightarrow \gamma_{\mathcal{E}}^*(NA)$ is a $\mathbb{T}$ -model homomorphism in $\mathcal{E}$. Then the assignment $A \mapsto fA$ is a $\mathbb{T}$ -model homomorphism $M \rightarrow N$ in $\mathbf{Set}$.*

Proof We have to prove that f preserves the interpretation of function symbols over Σ and the satisfaction of atomic relations over Σ . Since the preservation of the interpretation of function symbols over Σ can be expressed as the commutativity of a certain square involving finite products of sets of the form MA and NA , its satisfaction follows from that of $\gamma_{\mathcal{E}}^*(f)$ by virtue of Lemma 5.18. It remains to prove that f preserves the satisfaction of atomic relations over Σ . Let $R \mapsto A_1, \dots, A_n$ be a relation symbol over Σ . From the fact that $\gamma_{\mathcal{E}}^*(f) : \gamma_{\mathcal{E}}^*(M) \rightarrow \gamma_{\mathcal{E}}^*(N)$ preserves the satisfaction by R it follows that for any n -tuple $\vec{a} = (a_1, \dots, a_n) \in R_M$, the coproduct arrow $1_{\mathcal{E}} \rightarrow \coprod_{(b_1, \dots, b_n) \in NA_1 \times \dots \times NA_n} 1_{\mathcal{E}}$ corresponding to the n -tuple $(fA_1(a_1), \dots, fA_n(a_n))$ factors through the subobject $\coprod_{(b_1, \dots, b_n) \in R_N} 1_{\mathcal{E}} \mapsto \coprod_{(b_1, \dots, b_n) \in NA_1 \times \dots \times NA_n} 1_{\mathcal{E}}$. But this immediately implies, since distinct coproduct arrows are disjoint from each other and the topos $\mathcal{E}$ is non-trivial, that $f(\vec{a}) \in R_N$, as required. $\square$

Corollary 5.20. *Let Σ be a signature without relation symbols and $\mathbb{T}$ a geometric theory over a signature Σ' obtained from Σ by solely adding relation symbols whose interpretation in any $\mathbb{T}$ -model in a Grothendieck topos is the complement of the interpretation of a geometric formula over Σ (for instance, the injectivization of a geometric theory over Σ in the sense of Definition 6.33). Let $\mathcal{K}$ be a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$, a and b $\mathbb{T}$ -models in $\mathcal{K}$, M a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$ and $f : a \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, $g : b \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ Σ' -structure homomorphisms, where E is an object of $\mathcal{E}$. Let k be an assignment sending any sort A over Σ to a function $kA : aA \rightarrow bA$ such that $gA \circ kA = fA$. Suppose that either*

- (i) *there are no relation symbols in Σ' except possibly for a binary relation symbol which is $\mathbb{T}$ -provably complemented to the equality relation on a sort of Σ and the Σ' -structure homomorphisms f and g are sortwise monic; or*
- (ii) *$E \not\cong 0_{\mathcal{E}}$ and the $\mathbb{T}$ -model homomorphism $\tilde{g} : \gamma_{\mathcal{E}/E}^*(b) \rightarrow M$ corresponding to g under the identification of Proposition 4.5 is sortwise monic.*

Then the assignment $A \mapsto kA$ defines a $\mathbb{T}$ -model homomorphism $a \rightarrow b$ in $\mathbf{Set}$.

Proof We have to verify that k preserves the interpretation of function symbols over Σ' as well as the satisfaction by any binary relation symbol which is $\mathbb{T}$ -provably complemented to the equality relation on some sort over Σ .

By Lemma 6.34 below, k preserves the interpretation of any binary relation symbol which is $\mathbb{T}$ -provably complemented to the equality relation on some sort over Σ if and only if it is sortwise monic. This holds under hypothesis (i) since f is sortwise monic and for every sort A over Σ , $gA \circ kA = fA$.

On the other hand, under hypothesis (i), the fact that k preserves the interpretation of function symbols over Σ' follows from the fact that g and f do, since g is sortwise monic and $gA \circ kA = fA$ for every sort A over Σ . This shows that our thesis is satisfied under hypothesis (i).

Let us now suppose that hypothesis (ii) holds. Consider the $\mathbb{T}$ -model homomorphisms $\tilde{f} : \gamma_{\mathcal{E}/E}^*(a) \rightarrow M$ and $\tilde{g} : \gamma_{\mathcal{E}/E}^*(b) \rightarrow M$ corresponding to f and g under the identification of Proposition 4.5. Clearly, the assignment $A \rightarrow \tilde{k}A = \gamma_{\mathcal{E}/E}^*(kA)$ satisfies $\tilde{g}A \circ \tilde{k}A = \tilde{f}A$ for all sorts A over Σ . It thus follows from Lemma 5.17 that $A \rightarrow \tilde{k}A$ is a $\mathbb{T}$ -model homomorphism $\gamma_{\mathcal{E}/E}^*(a) \rightarrow \gamma_{\mathcal{E}/E}^*(b)$; but this in turn implies, by Corollary 5.19 (notice that $E \not\cong 0_{\mathcal{E}}$ if and only if the topos $\mathcal{E}/E$ is non-trivial), that the assignment $A \rightarrow kA$ is a $\mathbb{T}$ -model homomorphism $a \rightarrow b$, as required. $\square$

Theorem 5.21. *Let Σ be a signature without relation symbols and $\mathbb{T}$ a geometric theory over a signature Σ' obtained from Σ by solely adding relation symbols whose interpretation in any $\mathbb{T}$ -model in a Grothendieck topos is the complement of the interpretation of a geometric formula over Σ (for instance, the injectivization of a geometric theory over Σ in the sense of Definition 6.33). Let $\mathcal{K}$ be a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ whose objects are all finitely generated $\mathbb{T}$ -models. Suppose that*

- (i) *either all the arrows in $\mathcal{K}$ are sortwise monic homomorphisms and there are no relation symbols except possibly for a binary relation symbol which is $\mathbb{T}$ -provably complemented to the equality relation on a given sort of Σ or*
- (ii) *all the $\mathbb{T}$ -model homomorphisms in any Grothendieck topos are sortwise monic.*

Then $\mathbb{T}$ satisfies condition (iii)(b)-(2) of Theorem 5.1 with respect to the category $\mathcal{K}$.

Proof By Theorem 5.14, we have to verify that for any $\mathbb{T}$ -model c in $\mathcal{K}$, any object E of $\mathcal{E}$ and any Σ -structure homomorphism $z : c \rightarrow \text{Hom}_{\mathcal{E}}(E, \tilde{F}(M_{\mathbb{T}}))$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each index $i \in I$ a generalized element $x_i : E_i \rightarrow F(c)$ such that $\text{Hom}_{\mathcal{E}}(e_i, M) \circ z = \xi_{(c, x_i)}$ for all $i \in I$.

If $E \cong 0_{\mathcal{E}}$ then the condition is trivially satisfied; indeed, one can take $I = \emptyset$. We shall therefore suppose $E \not\cong 0_{\mathcal{E}}$ or, equivalently, that the topos $\mathcal{E}/E$ is non-trivial.

From now on, we shall suppose for simplicity that Σ is one-sorted, but all our arguments can be straightforwardly extended to the general case.

Let $\{r_1, \dots, r_n\}$ be a set of generators for the model c . Consider their images $z(r_1), \dots, z(r_n) : E \rightarrow M$ under the homomorphism z . By Proposition 3.11, for any $i \in \{1, \dots, n\}$ there exists an epimorphic family $\{e_j^i :$

$E_j^i \rightarrow E \mid j \in I_i\}$ in $\mathcal{E}$ and for each index $j \in J_i$ a $\mathbb{T}$ -model a_j^i in $\mathcal{K}$, a generalized element $x_j^i : E_j^i \rightarrow F(a_j^i)$ and an element $y_j^i \in a_j^i$ such that $\xi_{(a_j^i, x_j^i)}(y_j^i) = z(r_i) \circ e_j^i$.

For any tuple $\vec{k} = (k_1, \dots, k_n) \in J_1 \times \dots \times J_n$, consider the iterated pullback $e_{\vec{k}} : E_{\vec{k}} =: E_{k_1}^1 \times_E \dots \times_E E_{k_n}^n \rightarrow E$. The family of arrows $\{e_{\vec{k}} : E_{\vec{k}} \rightarrow E \mid \vec{k} \in J_1 \times \dots \times J_n\}$ is clearly epimorphic. For any $i \in \{1, \dots, n\}$ and $k_i \in J_i$, set $x_{\vec{k}}^i : E_{\vec{k}} \rightarrow F(a_{k_i}^i)$ equal to the composite of the generalized element $x_{k_i}^i : E_{k_i}^i \rightarrow F(a_{k_i}^i)$ with the canonical pullback arrow $p_{k_i}^i : E_{\vec{k}} \rightarrow E_{k_i}^i$. For any fixed $\vec{k} \in J_1 \times \dots \times J_n$, by inductively applying the joint embedding property of Proposition 3.11, we can find an epimorphic family $\{u_l^{\vec{k}} : U_l^{\vec{k}} \rightarrow E_{\vec{k}} \mid l \in L_{\vec{k}}\}$ and for each index $l \in L_{\vec{k}}$ a $\mathbb{T}$ -model $d_l^{\vec{k}}$ in $\mathcal{K}$, a generalized element $x_l^{\vec{k}} : U_l^{\vec{k}} \rightarrow F(d_l^{\vec{k}})$, and arrows $f_l^{\vec{k}} : a_{k_i}^i \rightarrow d_l^{\vec{k}}$ in $\mathcal{K}$ such that $x_{\vec{k}}^i \circ u_l^{\vec{k}} = F(f_l^{\vec{k}}) \circ x_l^{\vec{k}}$ (for all $i \in \{1, \dots, n\}$ and $l \in L_{\vec{k}}$).

Let us prove that $z(r_i) \circ e_{\vec{k}} \circ u_l^{\vec{k}} = \xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}(f_l^{\vec{k}}(y_{k_i}^i))$ (for any $i, \vec{k}$ and l).

We have already observed that for any $i \in \{1, \dots, n\}$, we have $z(r_i) \circ e_{k_i}^i = \xi_{(a_{k_i}^i, x_{k_i}^i)}(y_{k_i}^i)$. Composing both sides of the equation with $p_{k_i}^i$ and applying Lemma 3.10(ii) yields the equality $z(r_i) \circ e_{\vec{k}} = \xi_{(a_{k_i}^i, x_{\vec{k}}^i)}(y_{k_i}^i)$. On the other hand, by Lemma 3.10(i) we have that $\xi_{(a_{k_i}^i, x_{\vec{k}}^i)}(y_{k_i}^i) \circ u_l^{\vec{k}} = \xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}(f_l^{\vec{k}}(y_{k_i}^i))$. Our thesis thus follows by combining the former identity with the one obtained by composing both sides of the latter identity with $u_l^{\vec{k}}$.

Let us now consider the Σ' -structure homomorphisms

$$\text{Hom}_{\mathcal{E}}(e_{\vec{k}} \circ u_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) \circ z : c \rightarrow \text{Hom}_{\mathcal{E}}(U_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}}))$$

and

$$\xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})} : d_l^{\vec{k}} \rightarrow \text{Hom}_{\mathcal{E}}(U_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) .$$

Since, under either assumption (i) or (ii), the arrows of the category $\mathcal{K}$ are sortwise monic homomorphisms, by Proposition 5.15 the homomorphism $\xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}$ is injective. Therefore, since the image of all the generators $r_1, \dots, r_n$ of c under the homomorphism $\text{Hom}_{\mathcal{E}}(e_{\vec{k}} \circ u_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) \circ z$ is contained in the image of the homomorphism $\xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}$, an easy induction on the structure $t(r_1, \dots, r_n)$ of the elements of c shows that the image of all elements of c belongs to the image of $\xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}$; in other words, there exists a factorization $k_l^{\vec{k}} : c \rightarrow d_l^{\vec{k}}$ of $\text{Hom}_{\mathcal{E}}(e_{\vec{k}} \circ u_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) \circ z$ across $\xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})}$. By Corollary 5.20, such factorization is a $\mathbb{T}$ -model homomorphism. We have thus found an epimorphic family on E , namely $\{e_{\vec{k}} \circ u_l^{\vec{k}} \mid \vec{k} \in J_1 \times \dots \times J_n, l \in L_{\vec{k}}\}$, and for every $\vec{k}$ and l a $\mathbb{T}$ -model homomorphism $k_l^{\vec{k}} : c \rightarrow d_l^{\vec{k}}$ in $\mathcal{K}$ such that

$$\text{Hom}_{\mathcal{E}}(e_{\vec{k}} \circ u_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) \circ z = \xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})} \circ k_l^{\vec{k}} .$$

The composites $x_l^{\vec{k}'} : U_l^{\vec{k}} \rightarrow F(c)$ of the generalized elements $x_l^{\vec{k}} : U_l^{\vec{k}} \rightarrow F(d_l^{\vec{k}})$ with the arrows $F(k_l^{\vec{k}})$ thus yield generalized elements such that $\text{Hom}_{\mathcal{E}}(e_{\vec{k}} \circ u_l^{\vec{k}}, \tilde{F}(M_{\mathbb{T}})) \circ z = \xi_{(d_l^{\vec{k}}, x_l^{\vec{k}})} \circ k_l^{\vec{k}} = \xi_{(c, x_l^{\vec{k}'})}$, where the last equality follows by Lemma 3.10(i). This completes our proof. $\square$

Remark 5.22. The assumption that all the arrows in $\mathcal{K}$ be sortwise monic in condition (i) is weaker in general than the requirement of condition (ii) that all the $\mathbb{T}$ -model homomorphisms in any Grothendieck topos be sortwise monic. Anyway, condition (ii) is a necessary condition for $\mathbb{T}$ to be classified by the topos $[\mathcal{K}, \mathbf{Set}]$ if all the arrows in $\mathcal{K}$ are sortwise monic (cf. Corollary 6.9).

Corollary 5.23. *Let $\mathbb{T}$ be a geometric theory satisfying the hypotheses of Theorem 5.21 with respect to a small full subcategory $\mathcal{K}$ of $\mathbb{T}\text{-mod}(\mathbf{Set})$. Then $\mathbb{T}$ is of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$ if and only if it satisfies condition (i) of Theorem 5.1 and condition (ii)(a) of Theorem 5.7.*

Proof Condition (iii) of Theorem 5.1 is satisfied by Theorem 5.21, while the first part of condition (ii) holds by Remark 5.8(a). $\square$

5.3 Abstract reformulation

Thanks to the general theory of indexed filtered colimits developed in section 2, we can reformulate the conditions of Theorem 5.1 in more abstract, though less explicit, terms as follows.

Let $\mathbb{T}$ be a geometric theory over a signature Σ and $\mathcal{K}$ a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$.

Condition (i) of Theorem 5.1 for $\mathbb{T}$ with respect to $\mathcal{K}$ can be reformulated as the requirement that for every model M of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$, the $\mathcal{E}$ -indexed category of elements $\int^J \overline{H_M}$ of the functor H_M (or equivalently its $\mathcal{E}$ -final subcategory $\int \overline{H_{M_{\mathcal{E}}}}$, cf. Theorem 2.13) be $\mathcal{E}$ -filtered.

For any sort A over Σ , we have a functor $P_A : \mathcal{K} \rightarrow \mathbf{Set}$ of evaluation of models in $\mathcal{K}$ at the sort A . Using the notation of Theorem 2.5, we have an associated internal diagram $(P_A)_{\mathcal{E}}$ in $[\mathbb{K}, \mathcal{E}]$ and hence $\mathcal{E}$ -indexed functor $\underline{\mathbb{K}}_{\mathcal{E}} \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$. These functors, with A varying among the sorts over Σ , clearly lift to a $\mathcal{E}$ -indexed functor $P : \underline{\mathbb{K}}_{\mathcal{E}} \rightarrow \mathbb{T}\text{-mod}(\underline{\mathcal{E}})$.

In these terms, condition (ii) of Theorem 5.1 for the model M can be reformulated as the requirement that the canonical $\mathcal{E}$ -indexed cone over the $\mathcal{E}$ -indexed functor $P \circ \pi_{\overline{H_M}}^f$ be colimiting (cf. the proof of Proposition 3.11).

Conditions (i) and (ii) can thus be interpreted by saying that every $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ is canonically a $\mathcal{E}$ -filtered colimit of constant finitely presentable $\mathbb{T}$ -models in $\mathbf{Set}$ which belong to $\mathcal{K}$.

Under the assumption that conditions (i) and (ii) are satisfied, condition (iii) is equivalent to the requirement that for any $\mathbb{T}$ -model c in $\mathcal{K}$ and

any Grothendieck topos $\mathcal{E}$, the $\mathbb{T}$ -model $\gamma_{\mathcal{E}}^*(c)$ be $\mathcal{E}$ -finitely presentable (in the sense of section 4.4) or, more weakly, that the internal $\mathcal{E}$ -indexed hom functor $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), -) : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \underline{\mathcal{E}}_{\mathcal{E}}$ preserve $\mathcal{E}$ -filtered colimits of diagrams of the form $\gamma_{\mathcal{E}} \circ D$, where D is a diagram defined on a ($\mathcal{E}$ -final subcategory of a) small internal filtered category. Indeed, the fact that this condition is necessary for $\mathbb{T}$ to be of presheaf type follows from Theorem 4.13, while the fact that it is sufficient, together with conditions (i) and (ii) of Theorem 5.1, for $\mathbb{T}$ to be classified by the topos $[\mathcal{K}, \mathbf{Set}]$ can be proved by showing that it implies condition (iii)(a) of Theorem 5.1. In fact, this immediately follows from the fact that inverse image functors of geometric morphisms preserve internal (filtered) colimits in view of the fact that, by condition (ii), every $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ is canonically a $\mathcal{E}$ -filtered colimit of ‘constant’ finitely presentable $\mathbb{T}$ -models in $\mathcal{K}$.

Overall, we can conclude that a geometric theory $\mathbb{T}$ is of presheaf type if and only if every $\mathbb{T}$ -model in any Grothendieck topos $\mathcal{E}$ is a $\mathcal{E}$ -filtered colimit of its canonical diagram made of ‘constant’ finitely presentable models which are $\mathcal{E}$ -finitely presentable.

6 Faithful interpretations of theories of presheaf type

In this section we introduce the notion of faithful interpretation of geometric theories and establish sufficient criteria for faithful interpretations of theories of presheaf type to be again of presheaf type. We shall treat conditions (i), (ii) and (iii) of Theorem 5.1 separately.

6.1 General results

Definition 6.1. A geometric morphism $a : \mathbf{Set}[\mathbb{T}] \rightarrow \mathbf{Set}[\mathbb{T}']$ between the classifying toposes of two geometric theories is said to be a *faithful interpretation* of $\mathbb{T}'$ into $\mathbb{T}$ if the induced morphism

$$a_{\mathcal{E}} : \mathbb{T}\text{-mod}(\mathcal{E}) \rightarrow \mathbb{T}'\text{-mod}(\mathcal{E})$$

of categories of models is faithful, reflects isomorphisms and the equality relation between objects.

For any faithful interpretation of $\mathbb{T}'$ into $\mathbb{T}$ and any Grothendieck topos $\mathcal{E}$, we have a subcategory $\text{Im}(a_{\mathcal{E}})$ of $\mathbb{T}'\text{-mod}(\mathcal{E})$ whose objects (resp. arrows) are exactly the models (resp. the model homomorphisms) in the image of the functor $a_{\mathcal{E}}$. For any functor $F : \mathcal{A} \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$, F is full and faithful if and only if its composite $a_{\mathcal{E}} \circ F$ with $a_{\mathcal{E}}$ is full and faithful as a functor $\mathcal{A} \rightarrow \text{Im}(a_{\mathcal{E}})$.

Remarks 6.2. (a) Any *quotient* $\mathbb{S}$ of a geometric theory $\mathbb{T}$ defines a faithful interpretation of $\mathbb{T}$ into $\mathbb{S}$;

(b) Any *expansion* (in the sense of section 7.3) $\mathbb{T}'$ of a geometric theory $\mathbb{T}$ over a signature which does not contain new sorts with respect to the signature of $\mathbb{T}$ defines a faithful interpretation of $\mathbb{T}'$ into $\mathbb{T}$; in particular, the *injectivization* $\mathbb{T}_m$ (in the sense of Definition 6.33) of a geometric theory $\mathbb{T}$ defines a faithful interpretation of $\mathbb{T}$ into $\mathbb{T}_m$.

Given a faithful interpretation of $\mathbb{T}'$ into $\mathbb{T}$ as above, we can reformulate the conditions of Theorem 5.1 for the theory $\mathbb{T}$ with respect to a small full subcategory $\mathcal{K}$ of $\mathbb{T}\text{-mod}(\mathbf{Set})$ in alternative ways, as follows.

Condition (iii)(c) of Theorem 5.1 for $\mathbb{T}$ with respect to $\mathcal{K}$ can be formulated as follows: the composite functor

$$q_{\mathcal{E}} = a_{\mathcal{E}} \circ u_{\mathcal{K}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}'\text{-mod}(\mathcal{E})$$

is full and faithful into the image $Im(a_{\mathcal{E}})$ of $a_{\mathcal{E}}$, where $u_{\mathcal{K}}$ is the functor $\mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$ induced by the canonical geometric morphism $p_{\mathcal{K}} : [\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Set}[\mathbb{T}]$.

Condition (ii) of Theorem 5.1 for $\mathbb{T}$ with respect to $\mathcal{K}$ can be reformulated by saying that for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, the canonical morphism

$$q_{\mathcal{E}}(H_M) \rightarrow a_{\mathcal{E}}(M)$$

is an isomorphism.

Suppose moreover that the functor

$$a_{\mathbf{Set}} : \mathbb{T}\text{-mod}(\mathbf{Set}) \rightarrow \mathbb{T}'\text{-mod}(\mathbf{Set})$$

restricts to a functor

$$f : \mathcal{K} \rightarrow \mathcal{H},$$

where $\mathcal{H}$ is a small full subcategory of $\mathbb{T}'\text{-mod}(\mathbf{Set})$.

We have a commutative diagram

$$\begin{array}{ccc} [\mathcal{K}, \mathbf{Set}] & \xrightarrow{[f, \mathbf{Set}]} & [\mathcal{H}, \mathbf{Set}] \\ \downarrow p_{\mathcal{K}} & & \downarrow p_{\mathcal{H}} \\ \mathbf{Set}[\mathbb{T}] & \xrightarrow{a} & \mathbf{Set}[\mathbb{T}'], \end{array}$$

where $p_{\mathcal{K}}$ (resp. $p_{\mathcal{H}}$) is the canonical geometric morphism determined by the universal property of the classifying topos of $\mathbb{T}$ (resp. of $\mathbb{T}'$).

Also, for any $\mathbb{T}'$ -model M in a Grothendieck topos $\mathcal{E}$, the functors $a_{\mathcal{E}}$ induce a $\mathcal{E}$ -indexed functor

$$\int a : \int H_M^{\mathbb{T}} \rightarrow \int H_{a_{\mathcal{E}}(M)}^{\mathbb{T}'}$$

between the $\mathcal{E}$ -indexed categories of elements $\int H_M^{\mathbb{T}}$ and $\int H_{a_{\mathcal{E}}(M)}^{\mathbb{T}'}$ of the functors

$$H_M^{\mathbb{T}} := \text{Hom}_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), M) : \mathcal{K}^{\text{op}} \rightarrow \mathcal{E}$$

and

$$H_{a_{\mathcal{E}}(M)}^{\mathbb{T}'} := \text{Hom}_{\underline{\mathbb{T}'\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(-), a_{\mathcal{E}}(M)) : \mathcal{H}^{\text{op}} \rightarrow \mathcal{E}.$$

Since a is by our hypothesis a faithful interpretation of theories, $\int a$ is faithful and reflects equalities and isomorphisms. Suppose now that $\int a$ is moreover $\mathcal{E}$ -full (in the sense of Definition 2.9). Since for any functor F with values in a Grothendieck topos, F is flat if and only if its $\mathcal{E}$ -indexed category of elements is $\mathcal{E}$ -filtered, we conclude by Proposition 2.11 that if $\int a$ is $\mathcal{E}$ -final then the theory $\mathbb{T}$ satisfies condition (i) of Theorem 5.1 with respect to the category $\mathcal{K}$ if $\mathbb{T}'$ does with respect to the category $\mathcal{H}$. Moreover, Proposition 3.14 and Theorem 2.12 ensure that if $\mathbb{T}'$ satisfies condition (ii) of Theorem 5.1 with respect to the category $\mathcal{H}$ then $\mathbb{T}$ does with respect to the category $\mathcal{K}$.

Summarizing, we have the following result.

Theorem 6.3. *Let $a : \mathbf{Set}[\mathbb{T}] \rightarrow \mathbf{Set}[\mathbb{T}']$ be a faithful interpretation of geometric theories and let $\mathcal{K}$ and $\mathcal{H}$ be small subcategories respectively of $\mathbb{T}\text{-mod}(\mathbf{Set})$ and of $\mathbb{T}'\text{-mod}(\mathbf{Set})$ such that the functor*

$$a_{\mathbf{Set}} : \mathbb{T}\text{-mod}(\mathbf{Set}) \rightarrow \mathbb{T}'\text{-mod}(\mathbf{Set})$$

restricts to a functor $\mathcal{K} \rightarrow \mathcal{H}$ and the $\mathcal{E}$ -indexed functor

$$\int a : \int H_M^{\mathbb{T}} \rightarrow \int H_{a_{\mathcal{E}}(M)}^{\mathbb{T}'}$$

is $\mathcal{E}$ -full and satisfies the equivalent conditions of Proposition 6.5.

Then the theory $\mathbb{T}$ satisfies condition (i) (resp. condition (ii)) of Theorem 5.1 with respect to the category $\mathcal{K}$ if $\mathbb{T}'$ satisfies condition (i) (resp. condition (ii)) of Theorem 5.1 with respect to the category $\mathcal{H}$.

□

Remark 6.4. If $\mathbb{T}$ is the injectivization of $\mathbb{T}'$ (in the sense of Definition 6.33) or if $\mathbb{T}$ is a quotient of $\mathbb{T}'$ then the functor $\int a$ is $\mathcal{E}$ -full. Indeed, in the first case this follows from the fact that if the composite of two arrows is monic then the first one is monic, while in the second it follows from the fact that the category of models of $\mathbb{T}$ is a full subcategory of the category of models of $\mathbb{T}'$.

The following proposition provides an explicit rephrasing of the finality condition for the functor $\int a$.

Proposition 6.5. *Let $a : \mathbf{Set}[\mathbb{T}] \rightarrow \mathbf{Set}[\mathbb{T}']$ be a faithful interpretation of geometric theories let M be a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$ such that the functor $H_{a_{\mathcal{E}}(M)}^{\mathbb{T}'}$ is flat. Then the following conditions are equivalent:*

- (a) *the functor $\int a$ is $\mathcal{E}$ -final;*
- (b) *For any object E of $\mathcal{E}$, any $\mathbb{T}'$ -model c in $\mathcal{H}$, any $\mathbb{T}$ -model M in $\mathcal{E}$ and any $\mathbb{T}'$ -model homomorphism $x : \gamma_{\mathcal{E}/E}^*(c) \rightarrow !_E^*(a_{\mathcal{E}}(M))$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{T}'$ -model homomorphism $f_i : c \rightarrow a_{\mathbf{Set}}(c_i)$, where c_i is a $\mathbb{T}$ -model in $\mathcal{K}$, and a $\mathbb{T}'$ -model homomorphism $x_i : \gamma_{\mathcal{E}/E_i}^*(a_{\mathbf{Set}}(c_i)) \rightarrow !_{E_i}^*(a_{\mathcal{E}}(M))$ such that $x_i \circ \gamma_{\mathcal{E}/E_i}^*(f_i) = e_i^*(x)$ for all $i \in I$;*
- (c) *For any object E of $\mathcal{E}$ and any Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, a_{\mathcal{E}}(M))$, where c is a $\mathbb{T}'$ -model in $\mathcal{H}$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{T}'$ -model homomorphism $f_i : c \rightarrow a_{\mathbf{Set}}(c_i)$, where c_i is a $\mathbb{T}$ -model in $\mathcal{K}$, and a Σ -structure homomorphism $x_i : a_{\mathbf{Set}}(c_i) \rightarrow \text{Hom}_{\mathcal{E}}(E_i, a_{\mathcal{E}}(M))$ such that $x_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, a_{\mathcal{E}}(M)) \circ x$ for all $i \in I$.*

Proof The equivalence between the first two conditions immediately follows from the fact that, H_M being flat, the $\mathcal{E}$ -indexed category $\int H_M$ is $\mathcal{E}$ -filtered and hence Remark 2.10(c) applies, while the equivalence between the second and the third condition follows from Proposition 4.5. $\square$

Remark 6.6. If $\mathcal{E} = \mathbf{Set}$ and $\mathbb{T}$ is a quotient of $\mathbb{T}'$ the condition rewrites as follows: for any $\mathbb{T}$ -model M in $\mathbf{Set}$, any model c in $\mathcal{H}$ and any $\mathbb{T}'$ -model homomorphism $f : c \rightarrow M$, there exists a $\mathbb{T}$ -model d in $\mathcal{K}$ and homomorphisms $g : c \rightarrow d$ and $h : d \rightarrow M$ such that $h \circ g = f$.

In the particular case where $\mathbb{T}'$ is the empty theory over a signature Σ , $\mathbb{T}$ is a $\mathcal{F}$ -finitary geometric theory over Σ (in the sense of [32]), $\mathcal{H}$ is the category of finite Σ -structures and $\mathcal{K}$ is the category of finite $\mathbb{T}$ -models, the condition, required for every model M of $\mathbb{T}$ in $\mathbf{Set}$, specializes precisely to the ‘finite structure condition’ of [32].

The following result shows a relation between the action on points of a morphism of classifying toposes and a related induced action on flat functors.

Theorem 6.7. *Let $a : \mathbf{Set}[\mathbb{T}] \rightarrow \mathbf{Set}[\mathbb{T}']$ be an interpretation of geometric theories and $\mathcal{K}$ and $\mathcal{H}$ small subcategories respectively of $\mathbb{T}\text{-mod}(\mathbf{Set})$ and of $\mathbb{T}'\text{-mod}(\mathbf{Set})$ such that the functor*

$$a_{\mathbf{Set}} : \mathbb{T}\text{-mod}(\mathbf{Set}) \rightarrow \mathbb{T}'\text{-mod}(\mathbf{Set})$$

restricts to a functor $f : \mathcal{K} \rightarrow \mathcal{H}$. Then for any Grothendieck topos $\mathcal{E}$ the extension functor

$$\mathbf{Flat}(\mathcal{H}^{op}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{K}^{op}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$$

along the geometric morphism $[f, \mathbf{Set}] : [\mathcal{K}, \mathbf{Set}] \rightarrow [\mathcal{H}, \mathbf{Set}]$ takes values in the full subcategory $\text{Im}(a_{\mathcal{E}})$ of $\mathbb{T}\text{-mod}(\mathcal{E})$.

Proof The diagram

$$\begin{array}{ccc} [\mathcal{H}, \mathbf{Set}] & \xrightarrow{p_{\mathcal{H}}} & \mathbf{Set}[\mathbb{T}'] \\ \uparrow [f, \mathbf{Set}] & & \uparrow a \\ [\mathcal{K}, \mathbf{Set}] & \xrightarrow{p_{\mathcal{K}}} & \mathbf{Set}[\mathbb{T}] \end{array}$$

clearly commutes by definition of the functor f . Therefore, in view of Diaconescu's theorem, for any Grothendieck topos $\mathcal{E}$ we have a commutative diagram

$$\begin{array}{ccc} \mathbf{Flat}(\mathcal{H}^{\text{op}}, \mathcal{E}) & \xrightarrow{u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'}} & \mathbb{T}'\text{-mod}(\mathcal{E}) \\ \uparrow \xi_{\mathcal{E}} & & \uparrow a_{\mathcal{E}} \\ \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) & \xrightarrow{u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}} & \mathbb{T}\text{-mod}(\mathcal{E}), \end{array}$$

where $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}$ and $u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'}$ are the functors of section 3.3 and $\xi_{\mathcal{E}}$ is the extension functor induced by the geometric morphism $[f, \mathbf{Set}]$, from which our thesis immediately follows. $\square$

The following corollary can be obtained as a particular case of the theorem by taking $\mathbb{T}'$ to be the geometric theory of flat functors on $\mathcal{C}^{\text{op}}$, $\mathbb{T}$ to be its injectivization (in the sense of Definition 6.33), $\mathcal{H}$ equal to $\mathcal{C}$ and $\mathcal{K}$ equal to $\mathcal{D}$.

Corollary 6.8. *Let $\mathcal{D}$ be a small subcategory of a small category $\mathcal{C}$ whose arrows are all monic (in $\mathcal{C}$), and $\mathcal{E}$ a Grothendieck topos. Then*

- (a) *For any object $d \in \mathcal{D}$, $\tilde{F}(d)$ is a decidable object of $\mathcal{E}$;*
- (b) *For any natural transformation $\alpha : F \rightarrow G$ between two flat functors $F, G : \mathcal{D}^{\text{op}} \rightarrow \mathcal{E}$, $\tilde{\alpha}(c) : \tilde{F}(c) \rightarrow \tilde{G}(c)$ is monic in $\mathcal{E}$ for every $c \in \mathcal{C}$. In particular, $\alpha(d) : F(d) \rightarrow G(d)$ is monic in $\mathcal{E}$ for every $d \in \mathcal{D}$.*

$\square$

The following corollary is obtained by applying the theorem in the case $\mathbb{T}$ is equal to the injectivization of $\mathbb{T}'$ (in the sense of Definition 6.33) and $\mathcal{H} = \mathcal{K} = \text{f.p.}\mathbb{T}'\text{-mod}(\mathbf{Set})$.

Corollary 6.9. *Let $\mathbb{T}$ be a theory of presheaf type such that all the $\mathbb{T}$ -model homomorphisms between its finitely presentable (set-based) models are sortwise injective. Then every $\mathbb{T}$ -model homomorphism in every Grothendieck topos is sortwise monic.*

$\square$

6.2 Finitely presentable and finitely generated models

In this section, we discuss, for the purpose of establishing our main criteria for the injectivization of a theory of presheaf type to be again of presheaf type, the relationship between finitely presentable and finitely generated models of a given geometric theory.

Throughout this section, we shall assume for simplicity all our first-order signatures to be one-sorted, if not otherwise stated. It is certainly possible to extend our definitions and results to the general situation, but we shall not embark in the straightforward task of making this explicit.

Let Σ be a first-order signature. Recall from [20] that, given a Σ -structure M and a subset $A \subseteq M$, the Σ -structure $\langle A \rangle$ generated by A is the smallest Σ -substructure of M containing A ; by the proof of Theorem 1.2.3 [20], $\langle A \rangle$ can be concretely represented as the union $\bigcup_{n \in \mathbb{N}} A_n$, where the subsets A_n are defined inductively as follows: $A_0 = A \cup \{c^M\}$, where c^M are the interpretations in M of the constants over Σ , and $A_{n+1} = A_n \cup \{f^M(\vec{a}) : \text{for some } n > 0, f \text{ is a } n\text{-ary function symbol of } \Sigma \text{ and } \vec{a} \text{ is a } n\text{-tuple of elements of } A_n\}$. A Σ -structure M is said to be *finitely generated* if there exists a finite subset $A \subseteq M$ such that $M = \langle A \rangle$.

In the category $\mathbb{T}\text{-mod}(\mathbf{Set})$ of set-based models of a geometric theory $\mathbb{T}$, we say that an object B is a *quotient* of an object A if there exists a surjective $\mathbb{T}$ -model homomorphism $A \rightarrow B$.

Proposition 6.10. *Let $\mathbb{T}$ be a geometric theory over a signature Σ . Then any quotient in $\mathbb{T}\text{-mod}(\mathbf{Set})$ of a finitely generated $\mathbb{T}$ -model is finitely generated.*

Proof Let $q : M \rightarrow N$ be a surjective homomorphism of $\mathbb{T}$ -models. Suppose that M is finitely generated, by a finite subset $A \subseteq M$. Set $B = q(A)$. This set is clearly finite, and N is equal to $\langle B \rangle$. To prove this we show, by induction on n , that for any $n \in \mathbb{N}$, $q(A_n) = B_n$. As q is a Σ -structure homomorphism, we have that $q(A_0) = B_0$ and if $q(A_n) = B_n$ then clearly $q(A_{n+1}) = B_{n+1}$. Now, as q is surjective, $N = q(A) = q(\bigcup_{n \in \mathbb{N}} A_n) = \bigcup_{n \in \mathbb{N}} q(A_n) = \bigcup_{n \in \mathbb{N}} B_n$, as required. $\square$

Proposition 6.11. *Let $\mathbb{T}$ be a geometric theory over a signature Σ whose axioms are all of the form $(\phi \vdash_{\vec{x}} \psi)$, where ψ is a quantifier-free geometric formula. Then any finitely presentable $\mathbb{T}$ -model is finitely generated.*

Proof First, we notice that any geometric theory over a signature Σ such that all its axioms are of the form $(\phi \vdash_{\vec{x}} \psi)$, where ψ is a quantifier-free geometric formula, satisfies the property that any substructure of a model of $\mathbb{T}$ is a model of $\mathbb{T}$.

Let M be a finitely presentable $\mathbb{T}$ -model. We can clearly represent M as a filtered colimit (actually, directed union) of its finitely generated substructures (equivalently, submodels, cf. the above remark). By definition of finite presentability, the identity on M factors through one of the embeddings of a finitely generated $\mathbb{T}$ -submodel of M into M ; such embedding is thus necessarily an isomorphism, and M coincides with its domain. In particular, M is finitely generated, as required. $\square$

Proposition 6.12. *Let $\mathbb{T}$ be a geometric theory over a signature Σ whose category of set-based models is finitely accessible (in particular, a theory of presheaf type). Then any finitely generated $\mathbb{T}$ -model is a quotient of a finitely presentable models of $\mathbb{T}$ in the category $\mathbb{T}\text{-mod}(\mathbf{Set})$.*

If moreover all the axioms of $\mathbb{T}$ are of the form $(\phi \vdash_{\vec{x}} \psi)$, where ψ is a quantifier-free geometric formula, then the finitely generated $\mathbb{T}$ -models are precisely the quotients of the finitely presentable models of $\mathbb{T}$ in the category $\mathbb{T}\text{-mod}(\mathbf{Set})$.

Proof As the category $\mathbb{T}\text{-mod}(\mathbf{Set})$ is finitely accessible, M can be expressed as a filtered colimit ($M = \text{colim}(N_i), \{J_i : N_i \rightarrow M \mid i \in \mathcal{I}\}$) of finitely presentable models N_i . Let $a_1, \dots, a_n$ a finite set of generators for M . As the family of arrows $\{J_i : N_i \rightarrow M \mid i \in \mathcal{I}\}$ is jointly surjective (since filtered colimits in $\mathbb{T}\text{-mod}(\mathbf{Set})$ are computed pointwise as in $\mathbf{Set}$), for any generator a_j there exists an index $k_j \in \mathcal{I}$ and an element d_j of N_{k_j} such that $J_{k_j}(d_j) = a_j$. Now, the set of generators a_j being finite and the index category $\mathcal{I}$ being filtered, we can suppose without loss of generality that all the k_j are equal. We thus have an index $k \in \mathcal{I}$ and a string of elements of N_k which are sent by J_k to the generators of M ; hence the arrow $J_k : N_k \rightarrow M$ is surjective.

The second part of the proposition follows from Propositions 6.11 and 6.10, which ensure that every quotient of a finitely presentable model is finitely generated. $\square$

Proposition 6.13. *Let Σ be a signature without relation symbols and $\mathbb{T}$ a geometric theory over a larger signature Σ' obtained from Σ by solely adding relation symbols whose interpretation in any set-based $\mathbb{T}$ -model coincides with the complement of a geometric formula over Σ (for instance, the injectivization of a geometric theory over Σ). Suppose that all the $\mathbb{T}$ -model homomorphisms between set-based $\mathbb{T}$ -models are injective. Then any finitely generated $\mathbb{T}$ -model is finitely presentable.*

Proof From Proposition 6.11, we know that every finitely presentable $\mathbb{T}$ -model is finitely generated. It thus remains to prove the converse.

We observed in [6] (Lemma 6.2) that, given a category $\mathcal{D}$ with filtered colimits, an object M of $\mathcal{D}$ is finitely presentable in $\mathcal{D}$ if and only if for any

filtered diagram $D : \mathcal{I} \rightarrow \mathcal{D}$, any arrow $M \rightarrow \text{colim}(D)$ factors through one of the canonical colimit arrows $J_i : D(i) \rightarrow \text{colim}(D)$ and for any arrows $f : M \rightarrow D(i)$ and $g : M \rightarrow D(j)$ in $\mathcal{D}$ such that $J_i \circ f = J_j \circ g$, there exists $k \in \mathcal{I}$ and two arrows $s : i \rightarrow k$ and $t : j \rightarrow k$ such that $D(s) \circ f = D(t) \circ g$. Now, the latter condition is automatically satisfied if all the arrows of $\mathcal{D}$ are monic (since the category $\mathcal{I}$ is filtered); in particular, applying this criterion to the case of our theory $\mathbb{T}$ yields the following characterization of the finitely presentable $\mathbb{T}$ -models: a set-based $\mathbb{T}$ -model M is finitely presentable if and only if for any filtered colimit $\text{colim}(N_i)$ of set-based $\mathbb{T}$ -models, with colimit arrows $J_i : N_i \rightarrow \text{colim}(N_i)$, any $\mathbb{T}$ -model homomorphism $f : M \rightarrow \text{colim}(N_i)$ factors through at least one arrow J_i in the category $\mathbb{T}\text{-mod}(\mathbf{Set})$.

Now, if M is finitely generated by elements $a_1, \dots, a_n$, for any of generator a_i of M there exists an index k_i of $\mathcal{I}$ and an element $b_i \in N_{k_i}$ such that $f(a_i) = J_i(b_i)$. As the category $\mathcal{I}$ is filtered, we can suppose without loss of generality that all the k_i are equal to each other. Therefore there exists an index $k \in \mathcal{I}$ and elements $b_1, \dots, b_n$ of N_k such that $f(a_i) = J_k(b_i)$ for all i . The image of M under f is thus entirely contained in N_k and hence we have a function $g : M \rightarrow N_k$ such that $J_k \circ g = f$. Let us prove that g is a Σ' -structure homomorphism. The fact that g is a Σ -structure homomorphism, i.e. that it commutes with the interpretation of the function symbols over Σ , follows from the fact that f is by the injectivity of J_k . Concerning the preservation of atomic relations over Σ' , this is guaranteed by the fact that, were they not preserved, f would preserve the relations of satisfaction by elements of the geometric formulae over Σ defining the complements of such relations, contradicting the fact that f is a Σ' -structure homomorphism. $\square$

A useful sufficient condition for condition (iii)(b)-(1) of Theorem 5.1 to hold, which is applicable to categories $\mathcal{K}$ whose arrows are not necessarily monomorphisms is the following.

Theorem 6.14. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , and $\mathcal{K}$ a small full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ such that every model in $\mathcal{K}$ is a quotient of a $\mathbb{T}$ -model which is finitely presented by a geometric formula over Σ . Then $\mathbb{T}$ satisfies condition (iii)(b)-(1) of Theorem 5.1 with respect to the category $\mathcal{K}$.*

Proof We have to show that the extension functor

$$u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}_{J_{\mathbb{T}}}(\mathcal{C}_{\mathbb{T}}, \mathcal{E})$$

of section 3.3 is faithful.

By Theorem 3.5, for any formula $\{\vec{x} \cdot \phi\}$ presenting a $\mathbb{T}$ -model $M_{\{\vec{x} \cdot \phi\}}$, $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}(F)(\{\vec{x} \cdot \phi\}) \cong F(M_{\{\vec{x} \cdot \phi\}})$. The thesis thus follows immediately from Proposition 6.17. $\square$

6.3 Reformulations of condition (iii) of Theorem 5.1

Let us work in the context of a faithful interpretation of theories as in the last section.

Let

$$u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$$

and

$$u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'} : \mathbf{Flat}(\mathcal{H}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}'\text{-mod}(\mathcal{E})$$

be the functors of section 3.3.

Suppose that $\mathbb{T}'$ is of presheaf type and that $\mathcal{H}$ contains a category $\mathcal{P}$ whose Cauchy-completion coincides with the category of finitely presentable $\mathbb{T}'$ -models (so that $\mathbb{T}'$ is classified by the topos $[\mathcal{P}, \mathbf{Set}]$). Notice that, since the functor $f : \mathcal{K} \rightarrow \mathcal{H}$ is a restriction of the functor $a_{\mathbf{Set}}$, $\mathcal{K}$ gets identified under it with a subcategory of $\mathcal{H}$ and hence, by Remark 3.2(c), the extension functor

$$\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{K}^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{H}^{\text{op}}, \mathcal{E})$$

along the geometric morphism $[f, \mathbf{Set}]$ is faithful.

From the fact that $\mathbb{T}'$ faithfully interprets in $\mathbb{T}$ (in the sense of Definition 6.1) it follows that the functor $u_{(\mathcal{K}, \mathcal{E})}^{\mathbb{T}}$ is faithful (resp. full and faithful) if and only if the composite functor $u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'} \circ \xi_{\mathcal{E}}$ is, when regarded as a functor with values in the subcategory $Im(a_{\mathcal{E}})$ of $\mathbb{T}'\text{-mod}(\mathcal{E})$. The interest of this reformulation lies in the alternative description of the functor

$$u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'} : \mathbf{Flat}(\mathcal{H}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}'\text{-mod}(\mathcal{E})$$

which is available under our hypotheses. Indeed, the following proposition holds.

Proposition 6.15. *Under the hypotheses specified above, the functor*

$$u_{(\mathcal{H}, \mathcal{E})}^{\mathbb{T}'} : \mathbf{Flat}(\mathcal{H}^{\text{op}}, \mathcal{E}) \rightarrow \mathbb{T}'\text{-mod}(\mathcal{E}) \simeq \mathbf{Flat}(\mathcal{P}^{\text{op}}, \mathcal{E})$$

sends any flat functor $F : \mathcal{H}^{\text{op}} \rightarrow \mathcal{E}$ to its restriction $F|_{\mathcal{P}} : \mathcal{P}^{\text{op}} \rightarrow \mathcal{E}$, and acts accordingly on the natural transformations.

Proof This immediately follows from Theorem 3.5 in view of the fact that $\mathcal{H}$ contains $\mathcal{P}$ and all the objects of $\mathcal{P}$ are finitely presented $\mathbb{T}'$ -models (since $\mathbb{T}'$ is by our hypothesis of presheaf type classified by the topos $[\mathcal{P}, \mathbf{Set}]$). $\square$

As a corollary of the proposition, we immediately obtain the following result.

Theorem 6.16. *Under the hypotheses specified above, we have that*

- (i) $\mathbb{T}$ satisfies condition (iii)(b)-(1) of Theorem 5.1 with respect to the category $\mathcal{K}$ if and only if for any Grothendieck topos $\mathcal{E}$, every natural transformation between flat functors in the image of the functor $\xi_{\mathcal{E}}$ is determined by its restriction to the category $\mathcal{P}$;
- (ii) $\mathbb{T}$ satisfies condition (iii)(b)-(2) of Theorem 5.1 with respect to the category $\mathcal{K}$ if and only if for any Grothendieck topos $\mathcal{E}$, every natural transformation between restrictions to $\mathcal{P}$ of flat functors in the image of the functor $\xi_{\mathcal{E}}$ can be extended to a natural transformation between the functors themselves.

Let us now consider the satisfaction of condition (iii)(b) of Theorem 5.1 by the injectivization of a geometric theory $\mathbb{T}$ satisfying the hypotheses of Propositions 6.11 and 6.13. Since by the proposition every finitely generated $\mathbb{T}$ -model is finitely presentable in $\mathbb{T}\text{-mod}(\mathbf{Set})$ and every model of $\mathbb{T}$ is a directed union of its finitely generated submodels, the category $\mathbb{T}\text{-mod}(\mathbf{Set})$ is equivalent to $\mathbf{Flat}(\text{f.g.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, \mathbf{Set})$, where $\text{f.g.}\mathbb{T}\text{-mod}(\mathbf{Set})$ is the full subcategory of $\mathbb{T}\text{-mod}(\mathbf{Set})$ on the finitely generated $\mathbb{T}$ -models. The following theorem shows that condition (iii)(b)-(1) is satisfied. Before stating it, we need a lemma.

Lemma 6.17. *Let $\mathcal{C}$ be a small category, $\mathcal{E}$ a topos and $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. Then for any epimorphism $f : a \rightarrow b$ in $\mathcal{C}$, the arrow $F(f) : F(b) \rightarrow F(a)$ is monic in $\mathcal{E}$.*

Proof The thesis follows at once from the fact that flat functors preserve all the finite limits which exist in the domain category using the well-known characterization of monomorphisms in terms of pullbacks. $\square$

Theorem 6.18. *Let $\mathbb{T}$ be the injectivization of a theory of presheaf type $\mathbb{T}'$ (in the sense of Definition 6.33) such that the every finitely presentable $\mathbb{T}'$ -model is finitely generated (cf. for instance Proposition 6.11) and the finitely generated $\mathbb{T}'$ -models are all quotients (in the sense of section 6.2) of finitely presentable $\mathbb{T}'$ -models (cf. for instance Proposition 6.12). Then $\mathbb{T}$ satisfies condition (iii)(b)-(1) of Theorem 5.1 with respect to the category of finitely generated $\mathbb{T}'$ -models.*

Proof It suffices to apply Proposition 6.15 and Lemma 6.17 in conjunction with Theorem 6.16 by taking $\mathcal{K}$ equal to the category of finitely generated $\mathbb{T}'$ -models and sortwise monic homomorphisms between them, $\mathcal{H}$ equal to the category of finitely generated $\mathbb{T}'$ -models and homomorphisms between them and $\mathcal{P}$ equal to the category of finitely presentable $\mathbb{T}'$ -models. $\square$

The following proposition can be useful in verifying that condition (ii) of Theorem 6.16 holds.

Proposition 6.19. *Let $\mathcal{D}$ be a full subcategory of a category $\mathcal{C}$ with filtered colimits such that*

- *for any object c of $\mathcal{C}$ there exists an epimorphism $a \rightarrow c$ in $\mathcal{C}$ from an object a of $\mathcal{D}$ to c ,*
- *every object of $\mathcal{D}$ is finitely presentable in $\mathcal{C}$ and*
- *every object c of $\mathcal{C}$ can be canonically expressed as a filtered colimit $\text{colim}(d_i)$ of objects of $\mathcal{D}$, with colimit arrows $J_i : d_i \rightarrow c$ (for $i \in \mathcal{I}$).*

Let F and G be flat functors $\mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$. Then a natural transformation $\beta : F|_{\mathcal{D}} \rightarrow G|_{\mathcal{D}}$ lifts (uniquely) to a natural transformation $F \rightarrow G$ if and only if for every epimorphism $q : a \rightarrow b$ of the form J_i , the arrow $\beta(a) : F(a) \rightarrow G(a)$ restricts to an arrow $F(b) \rightarrow G(b)$ along the monomorphisms $F(q) : F(b) \rightarrow F(a)$ and $G(q) : G(b) \rightarrow G(a)$.

Proof Clearly, the ‘only if’ direction is trivially satisfied, so we only have to care about the ‘if’ direction. The uniqueness of the extended natural transformation follows from Proposition 6.17, so we just have to prove the existence.

Notice that for every object c of $\mathcal{C}$ there exists an $i \in \mathcal{I}$ such that the arrow $J_i : d_i \rightarrow c$ is an epimorphism. Indeed, by our hypotheses there exists an epimorphism $q : a \rightarrow c$ to c from an object a of $\mathcal{D}$ to c and, since a is finitely presentable in $\mathcal{C}$, q necessarily factors through an arrow of the form J_i , which is necessarily epic as q is.

Let $\beta : F|_{\mathcal{D}} \rightarrow G|_{\mathcal{D}}$ be a natural transformation. Choose an $i \in \mathcal{I}$ such that the arrow $J_i : d_i \rightarrow c$ is an epimorphism (such arrow exists by the above remark), and set $\alpha(c)$ equal to the restriction of $\beta(d_i)$ along the arrows $F(J_i)$ and $G(J_i)$.

First, let us verify that this definition is well-posed, i.e. that it does not depend on the choice of $i \in \mathcal{I}$. For any other index j such that J_j is an epimorphism with codomain c , the filteredness of $\mathcal{I}$ ensures the existence of an index k and two arrows $\xi : i \rightarrow k$ and $\chi : j \rightarrow k$ in $\mathcal{I}$, whence $J_k \circ \xi = J_i$ and $J_k \circ \chi = J_j$. Now, if we denote by u the restriction of the arrow $\beta(d_k)$ along the arrows $F(J_k)$ and $G(J_k)$ we obtain, by invoking the naturality of β with respect to the arrows ξ and χ in $\mathcal{I}$, that u is equal to the unique restriction of $\beta(d_i)$ along the arrows $F(J_i)$ and $G(J_i)$ and to the unique restriction of $\beta(d_j)$ along the arrows $F(J_j)$ and $G(J_j)$; in particular, these two restrictions are equal, as required.

It remains to show that the assignment $c \rightarrow \alpha(c)$ is natural in c , i.e. that for any arrow $f : c \rightarrow c'$ in $\mathcal{C}$, the following naturality square commutes:

$$\begin{array}{ccc} F(c) & \xrightarrow{\alpha(c)} & G(c) \\ F(f) \uparrow & & \uparrow G(f) \\ F(c') & \xrightarrow{\alpha(c')} & G(c') \end{array}.$$

To prove this we observe that, using the fact that there exists an epimorphism $J_i : d_i \rightarrow c$ from an object of $\mathcal{D}$ to c , we can suppose without loss of generality c to lie in $\mathcal{D}$; indeed, since G sends epimorphisms to monomorphisms, the commutativity of the naturality square above is equivalent, by definition of $\alpha(c)$, to the commutativity of the naturality square relative to the arrow $f \circ J_i$. Now, consider the canonical representation of c' as a filtered colimit $\text{colim}(e_j)$ (for $j \in \mathcal{J}$) of objects in $\mathcal{D}$, with colimit arrows $K_j : e_j \rightarrow c'$ ($j \in \mathcal{J}$). Since c lies in $\mathcal{D}$, c is finitely presentable in $\mathcal{C}$ by our hypotheses and hence there exists an index $j \in \mathcal{J}$ and an arrow $r : c \rightarrow e_j$ such that $K_j \circ r = f$. The commutativity of our diagram thus follows from the commutativity of the naturality square of β relative to the arrow r and the definition of $\alpha(c')$ in terms of $\beta(e_j)$. $\square$

The following result, obtained by combining Theorem 6.45 and Proposition 6.19, gives a criterion for condition (iii)(b)-(2) to hold.

Theorem 6.20. *Let $\mathbb{T}$ be the injectivization of a theory of presheaf type $\mathbb{T}'$ such that every finitely presentable $\mathbb{T}'$ -model is finitely generated (cf. for instance Proposition 6.11), the finitely generated $\mathbb{T}'$ -models are all quotients (in the sense of section 6.2) of finitely presentable $\mathbb{T}'$ -models (cf. for instance Proposition 6.12) and every monic $\mathbb{T}'$ -model homomorphism between finitely generated models of $\mathbb{T}$ is sortwise monic. Then $\mathbb{T}$ satisfies condition (iii)(b)-(2) of Theorem 5.1 with respect to the category $\mathcal{C}$ of finitely generated models of $\mathbb{T}'$ and sortwise monic homomorphisms between them if and only if for any flat functors $F, G : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$, denoting by $\tilde{F}$ and $\tilde{G}$ their extension to the category of finitely generated $\mathbb{T}'$ -models and homomorphisms between them and by $\mathcal{P}$ the full subcategory of $\mathcal{C}$ on the finitely presentable $\mathbb{T}'$ -models, any natural transformation $\alpha : F|_{\mathcal{P}} \rightarrow G|_{\mathcal{P}}$ satisfies the following property: for any quotient $\mathbb{T}$ -model homomorphism $q : a \rightarrow b$, where a is finitely presentable and b is finitely generated, the arrow $\alpha(b)$ restricts along the monic arrows $F(q)$ and $G(q)$.*

$\square$

Remark 6.21. Notice that if F and G are representable functors then the condition of the theorem is trivially satisfied, the category $\mathbb{T}'\text{-mod}(\mathbf{Set})$ being finitely accessible.

6.4 Quotient theories

Recall from [9] that a *quotient* of a geometric theory $\mathbb{T}$ over a signature Σ is a geometric theory $\mathbb{T}'$ over Σ such that every axiom of $\mathbb{T}$ is provable in $\mathbb{T}'$.

It is often useful, in investigating whether a given geometric theory $\mathbb{S}$ is of presheaf type, to consider it in relation to a theory $\mathbb{T}$ of presheaf type of which $\mathbb{S}$ is a quotient.

The following result is a corollary of Theorem 6.3 and Proposition 6.5.

Corollary 6.22. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ and $\mathbb{S}$ be a quotient of $\mathbb{T}$ such that all the finitely presentable $\mathbb{S}$ -models are finitely presentable as $\mathbb{T}$ -models (for instance, $\mathbb{T}$ can be the empty theory over a finite signature and $\mathbb{S}$ can be any geometric theory over Σ whose finitely presentable models are all finite, cf. Theorem 6.4 [6]). Suppose moreover that for any object E of a Grothendieck topos $\mathcal{E}$ and any Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c is a finitely presentable $\mathbb{T}$ -model and M is a $\mathbb{S}$ -model in $\mathcal{E}$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{T}$ -model homomorphism $f_i : c \rightarrow c_i$, where c_i is a finitely presentable $\mathbb{S}$ -model, and a Σ -structure homomorphism $x_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $x_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ x$ for all $i \in I$. Then $\mathbb{S}$ is of presheaf type.*

Proof Condition (iii) of Theorem 5.1 satisfied by Proposition 5.11, while the fact that conditions (i) and (ii) of Theorem 5.1 are satisfied follows immediately from Theorem 6.3 in view of Proposition 6.5 (take $\mathbb{T}$ equal to $\mathbb{S}$, $\mathbb{T}'$ equal to $\mathbb{T}$, $\mathcal{K}$ equal to the category of finitely presentable $\mathbb{S}$ -models and $\mathcal{H}$ equal to the category of finitely presentable $\mathbb{T}$ -models - the fact that $\int a$ is $\mathcal{E}$ -full is clear). $\square$

We shall say that the family of homomorphisms x_i in the statement of the corollary defines a ‘localization’ of the homomorphism x .

Remarks 6.23. (a) Corollary 6.22 generalizes Theorem 3.6 [32], whose hypotheses, which are expressed syntactically in terms of geometric logic, once interpreted in topos-theoretic semantics, are stronger than those of Corollary 6.22 in the particular case where $\mathbb{T}$ is the empty theory over a finite signature and $\mathbb{S}$ is any geometric theory over Σ whose finitely presentable models are all finite.

(b) Theorem 6.28 follows as an immediate consequence of the corollary, observing that if $\mathbb{T}$ and $\mathbb{T}'$ are theories as in the hypotheses of the theorem then any $\mathbb{T}$ -model homomorphism having as codomain a $\mathbb{T}'$ -model also has as domain a $\mathbb{T}'$ -model whence the hypotheses of the corollary are trivially satisfied.

- (c) Assuming the first hypothesis of the corollary, the second is actually necessary for $\mathbb{S}$ to be of presheaf type. Indeed, if $\mathbb{S}$ is of presheaf type then every model M of $\mathbb{S}$ is a $\mathcal{E}$ -filtered colimit of the associated canonical diagram of ‘constant’ finitely presentable $\mathbb{S}$ -models (cf. sections 4.4 and 5.3), which by the first hypothesis are also finitely presentable $\mathbb{T}$ -models; the $\mathcal{E}$ -finite presentability of c thus implies the second hypothesis of the corollary (cf. Proposition 4.15).
- (d) Corollary 6.22 is often applied to pairs of the form $(\mathbb{S}, \mathbb{T} = \mathbb{S}_c)$, where $\mathbb{S}$ is a geometric theory over a signature Σ and $\mathbb{T}$ equal to the *cartesianization* (or finite-limit part) of $\mathbb{S}_c$ of $\mathbb{S}$, namely the cartesian theory over Σ consisting of all the cartesian sequents over Σ which are provable in $\mathbb{S}$ (recall that a $\mathbb{S}$ -cartesian sequent is a sequent over Σ involving $\mathbb{S}$ -cartesian formulae over Σ , that is formulae built from atomic formulae by only using finitary conjunctions and $\mathbb{S}$ -provably unique existential quantifications). Notice that the cartesianization of $\mathbb{S}$ is the biggest cartesian theory over the signature of $\mathbb{S}$ of which $\mathbb{S}$ is a quotient.

The question of whether for any theory of presheaf type $\mathbb{T}$, every finitely presentable $\mathbb{T}$ -model is finitely presentable as a $\mathbb{T}_c$ -model is still open and will be addressed in full generality in another paper. For the moment, we limit ourselves to remarking that this property is satisfied by all the examples of theories of presheaf type considered in this paper (cf. section 8).

Corollary 6.22 can also be applied to pairs consisting of a geometric theory over a signature Σ and the empty theory over the same signature, provided that the former satisfies appropriate hypotheses. In order to apply the corollary in this context, we need the following lemma. Below, when we say that a Σ -structure c is finite we mean that for every sort A over Σ , cA is finite.

Lemma 6.24. *Let $\mathbb{T}$ be a geometric theory over a finite signature Σ with a finite number of axioms each of which is of the form $(\top \vdash_{\vec{x}} \bigvee_{i \in I} \phi_i)$, where the ϕ_i are finite conjunctions of atomic formulae. Then for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$, any object E of $\mathcal{E}$ and any Σ -structure homomorphism $f : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ from a finite Σ -structure c there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a finite Σ -substructure c_i of $\text{Hom}_{\mathcal{E}}(E_i, M)$ which is a model of $\mathbb{T}$ and a Σ -structure homomorphism $f_i : c \rightarrow c_i$ such that $\text{Hom}_{\mathcal{E}}(e_i, M) \circ f = j_i \circ f_i$ (where j_i is the canonical inclusion of c_i into $\text{Hom}_{\mathcal{E}}(E_i, M)$).*

Proof For each axiom σ of the form $(\top \vdash_{\vec{x}_\sigma} \bigvee_{i \in I_\sigma} \phi_i^\sigma)$ and any finite tuple $\vec{\xi}$ of elements of c of the same type as $\vec{x}_\sigma$, since M is by our hypotheses a

$\mathbb{T}$ -model, there exists an epimorphic family $\{e_i^{\vec{\xi}} : E_i^{\vec{\xi}} \rightarrow E \mid i \in I_\sigma\}$ in $\mathcal{E}$ such that for any $i \in I_\sigma$, $f(\vec{\xi}) \circ e_i^{\vec{\xi}}$ factors through $[[\vec{x}_\sigma \cdot \phi_i]]_M$. Since there is only a finite number of axioms of $\mathbb{T}$ and of elements of c , and the fibered product of a finite number of epimorphic families is again an epimorphic family, there exists an epimorphic family $\{e_k : E_k \rightarrow E \mid k \in K\}$ such that for any $k \in K$, any axiom σ of $\mathbb{T}$ and any tuple $\vec{\xi}$ of elements of c of the appropriate sorts, the element $f(\vec{\xi}) \circ e_k \in \text{Hom}_{\mathcal{E}}(E_k, M)$ satisfies the formula on the right-hand side of the sequent σ . Hence if we consider, for each $k \in K$, the surjection-inclusion factorization $i_k \circ f_k$ of the homomorphism $\text{Hom}_{\mathcal{E}}(e_k, M) \circ f : c \rightarrow \text{Hom}_{\mathcal{E}}(E_k, M)$ (in the sense of Lemma 6.51), we obtain that c_k is finite, since f_k is surjective, and that all the axioms of $\mathbb{T}$ are satisfied in c_k (since the formulae ϕ_i are finite conjunctions of atomic formulae and c_k is a substructure of $\text{Hom}_{\mathcal{E}}(E_k, M)$), that is all the c_k are models of $\mathbb{T}$. This proves our thesis. $\square$

Corollary 6.25. *Let $\mathbb{T}$ be a geometric theory over a finite signature Σ with a finite number of axioms each of which is of the form $(\top \vdash_{\vec{x}} \bigvee_{i \in I} \phi_i)$, where the ϕ_i are finite conjunctions of atomic formulae. Suppose that for any $\mathbb{T}$ -model M in a Grothendieck topos $\mathcal{E}$ and object E of $\mathcal{E}$, every finitely generated Σ -substructure of $\text{Hom}_{\mathcal{E}}(E, M)$ has sortwise only a finite number of elements besides the constants (for instance, when the signature Σ does not contain function symbols except for a finite number of constants). Then $\mathbb{T}$ is of presheaf type, classified by the category of covariant set-valued functors on the category of finite models of $\mathbb{T}$.*

Proof As the signature of $\mathbb{T}$ is finite, every model which sortwise contains only finitely many elements besides the constants is finitely presentable as a model of the empty theory over the signature of $\mathbb{T}$ (cf. Theorem 6.4 [6]). This remark ensures, by Lemma 6.24, that the theory $\mathbb{T}$ satisfies the hypotheses of Corollary 6.22 with respect to the empty theory over its signature. Since the latter theory is clearly of presheaf type, it follows that $\mathbb{T}$ is of presheaf type as well, as required. $\square$

6.4.1 Presheaf-type quotients and rigid topologies

In this section we shall analyze the presheaf-type quotients of a presheaf-type theory $\mathbb{T}$ in terms of the associated subtoposes of the classifying topos for $\mathbb{T}$.

We know from [9] that every quotient $\mathbb{T}'$ a theory of presheaf type $\mathbb{T}$ corresponds to a unique Grothendieck topology J on the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ (under the duality theorem of [9]).

Recall from [21] (Definition C2.2.18) that a Grothendieck topology J on a category $\mathcal{C}$ is said to be *rigid* if for every object c of $\mathcal{C}$, the family of all the arrows to c in $\mathcal{C}$ from J -irreducible objects of $\mathcal{C}$ (i.e., objects d with the

property that the only J -covering sieve on d is the maximal one), generates a J -covering sieve.

We can characterize the topologies J such that the corresponding subtopos $\mathbf{Sh}(\mathbf{f.p.T-mod}(\mathbf{Set})^{\text{op}}, J) \hookrightarrow [\mathbf{f.p.T-mod}(\mathbf{Set}), \mathbf{Set}]$ is an essential geometric inclusion (that is, a geometric inclusion whose inverse image which admits a left adjoint) by using the following site characterizations:

(1) A canonical geometric inclusion $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ from a topos $\mathbf{Sh}(\mathcal{C}, J)$ which is equivalent to a presheaf topos and such that $\mathcal{C}$ is a Cauchy-complete category, is essential if and only if the topology J is rigid;

(2) A geometric morphism (resp. geometric inclusion) $[\mathcal{C}, \mathbf{Set}] \rightarrow [\mathcal{D}, \mathbf{Set}]$, where $\mathcal{D}$ is a Cauchy-complete category, is essential if and only if it is induced by a functor (resp. by a full and faithful functor) $\mathcal{C} \rightarrow \mathcal{D}$.

The latter characterization is well-known (cf. Lemma A4.1.5 [21] and Example A4.2.12(b)), while the former can be proved as follows. If J is rigid then, denoting by $\mathcal{D}$ the full subcategory of $\mathcal{C}$ on the J -irreducible objects, the Comparison Lemma yields an equivalence $\mathbf{Sh}(\mathcal{C}, J) \simeq [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ which makes the canonical inclusion $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ isomorphic to the canonical geometric inclusion $[\mathcal{D}^{\text{op}}, \mathbf{Set}] \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ induced by the full inclusion of categories $\mathcal{D}^{\text{op}} \hookrightarrow \mathcal{C}^{\text{op}}$; in particular, the morphism $\mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is essential. Conversely, if the canonical geometric inclusion $i_J : \mathbf{Sh}(\mathcal{C}, J) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ is essential then if $\mathbf{Sh}(\mathcal{C}, J)$ is equivalent to $[\mathcal{D}^{\text{op}}, \mathbf{Set}]$, by property (2) i_J is isomorphic to a geometric inclusion $[\mathcal{D}^{\text{op}}, \mathbf{Set}] \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ induced by a full embedding $\mathcal{D}^{\text{op}} \hookrightarrow \mathcal{C}^{\text{op}}$. Now, the topology $J_{\mathcal{D}}$ on $\mathcal{C}$ defined by saying that a sieve R on c is J -covering if and only if it contains all the morphisms from objects of $\mathcal{D}$ to c is clearly rigid, and the Comparison Lemma yields an equivalence $\mathbf{Sh}(\mathcal{C}, J_{\mathcal{D}}) \simeq [\mathcal{D}^{\text{op}}, \mathbf{Set}]$ which makes the geometric morphism $[\mathcal{D}^{\text{op}}, \mathbf{Set}] \rightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ isomorphic to the canonical inclusion $\mathbf{Sh}(\mathcal{C}, J_{\mathcal{D}}) \hookrightarrow [\mathcal{C}^{\text{op}}, \mathbf{Set}]$ (cf. Example A2.2.4(d) [21]). It thus follows that $J = J_{\mathcal{D}}$; in particular, J is rigid.

These site characterizations define the arches of a ‘bridge’ (in the sense of [13]) leading to the following result (cf. also Theorem 6.8 [6] for a related result).

Theorem 6.26. *Let $\mathbb{T}'$ be a quotient of a theory of presheaf type $\mathbb{T}$, corresponding to a Grothendieck topology J on the category $\mathbf{f.p.T-mod}(\mathbf{Set})^{\text{op}}$ under the duality theorem of [9]. Suppose that $\mathbb{T}'$ is itself of presheaf type. Then every finitely presentable $\mathbb{T}'$ -model is finitely presentable also as a $\mathbb{T}$ -model if and only if the topology J is rigid.*

Proof If every finitely presentable $\mathbb{T}'$ -model is finitely presentable also as a $\mathbb{T}$ -model then the geometric inclusion corresponding via the duality theorem to the quotient $\mathbb{T}'$ of $\mathbb{T}$ is induced by the inclusion of the respective full subcategory of $\mathbf{f.p.T-mod}(\mathbf{Set})$ on the $\mathbb{T}'$ -models. In view of the above characterizations, the topology J is rigid. Conversely, if J is rigid then

$\mathbf{Sh}(\mathbf{f.p.T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$ is equivalent to the presheaf topos $[\mathcal{D}, \mathbf{Set}]$, where $\mathcal{D}^{\text{op}}$ is the full subcategory of $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})^{\text{op}}$ on the J -irreducible objects. Clearly, this subcategory is Cauchy-complete (as $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})$ is Cauchy-complete and the condition of J -irreducibility is stable under retracts) and hence, since $\mathbb{T}'$ is of presheaf type, it is equivalent to the category of finitely presentable $\mathbb{T}'$ -models; in particular, any such model is finitely presentable as a $\mathbb{T}$ -model. $\square$

Remark 6.27. (i) Under the hypothesis that every finitely presentable $\mathbb{T}'$ -model is finitely presentable as a $\mathbb{T}$ -model, the full subcategory $\mathbf{f.p.T}'\text{-mod}(\mathbf{Set})$ of $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})$ and the topology J can be defined in terms of each other as follows (cf. the proof of the theorem): the objects of $\mathbf{f.p.T}'\text{-mod}(\mathbf{Set})$ are precisely the J -irreducible objects of $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})$, and a sieve S in $\mathbf{f.p.T}'\text{-mod}(\mathbf{Set})$ on an object c is J -covering if and only if it contains all the arrows in $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})$ from finitely presentable $\mathbb{T}'$ -models to c .

(ii) If $\mathbb{T}'$ is a presheaf-type quotient of a theory of presheaf type $\mathbb{T}$ then for any finitely presentable $\mathbb{T}$ -model c , any $\mathbb{T}'$ -model M and any $\mathbb{T}$ -model homomorphism $f : c \rightarrow M$, there exists a $\mathbb{T}$ -model homomorphism $g : c \rightarrow c'$ to a finitely presentable $\mathbb{T}'$ -model c' and a $\mathbb{T}$ -model homomorphism $h : c' \rightarrow M$ such that $h \circ g = f$. Indeed, M can be expressed as a filtered colimit of finitely presentable $\mathbb{T}'$ -models c' ; therefore, as c is finitely presentable as a $\mathbb{T}$ -model, f necessarily factors through a colimit arrow $c' \rightarrow M$.

(iii) Under the hypothesis that every finitely presentable $\mathbb{T}'$ -model is finitely presentable as a $\mathbb{T}$ -model, by the duality theorem of [9], the syntactic description of $\mathbb{T}'$ in terms of J given therein and the description of J in terms of $\mathbf{f.p.T}'\text{-mod}(\mathbf{Set})$ given in Remark 6.27(a), $\mathbb{T}'$ can be characterized as the quotient of $\mathbb{T}$ obtained by adding all the sequents of the form $(\phi(\vec{x}) \vdash_{\vec{x}} \bigvee_{i \in I} \exists \vec{x}_i \theta_i(\vec{x}_i, \vec{x}))$, where $\{\theta_i(\vec{x}_i, \vec{x}) : \phi_i(\vec{x}_i) \rightarrow \phi(\vec{x})\}$ is the family of $\mathbb{T}$ -provably functional formulae from the formulae $\phi_i(\vec{x}_i)$ presenting a $\mathbb{T}'$ -model to a $\mathbb{T}$ -irreducible formula (equivalently, a formula presenting a $\mathbb{T}$ -model) $\phi(\vec{x})$.

The following result provides a natural class of presheaf-type quotients of presheaf-type theories whose associated topologies are rigid.

Theorem 6.28. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ . Then any quotient $\mathbb{T}'$ of $\mathbb{T}$ obtained from $\mathbb{T}$ by adding sequents of the form $\phi \vdash_{\vec{x}} \perp$, where $\phi(\vec{x})$ is a geometric formula over Σ , is classified by the topos $[\mathcal{T}, \mathbf{Set}]$, where $\mathcal{T}$ is the full subcategory of $\mathbf{f.p.T}\text{-mod}(\mathbf{Set})$ on the $\mathbb{T}'$ -models.*

Proof By covering each $\phi(\vec{x})$ by $\mathbb{T}$ -irreducible formulae in the syntactic category $\mathcal{C}_{\mathbb{T}}$ of $\mathbb{T}$, we can suppose without loss of generality all of the $\phi(\vec{x})$ to

be $\mathbb{T}$ -irreducible, that is to present a $\mathbb{T}$ -model M_ϕ . Then, by the results of [9], the quotient $\mathbb{T}'$ is classified by the topos $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J)$, where J is the smallest Grothendieck topology on $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}$ which contains all the empty sieves on the models M_ϕ presented by the formulae ϕ involved in the axioms $\phi \vdash_{\vec{x}} \perp$ added to $\mathbb{T}$ to form $\mathbb{T}'$. Let $\mathcal{T}$ be the full subcategory of $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ on the $\mathbb{T}'$ -models. Then $\mathcal{T}$ is J -dense; indeed, any object not in $\mathcal{T}$ admits an arrow from a model of the form M_ϕ and hence is covered by the empty sieve. Further, for any object of $\mathcal{T}$, the J -covering sieves on it are exactly the maximal ones; therefore, the Comparison Lemma yields an equivalence $\mathbf{Sh}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, J) \simeq [\mathcal{T}, \mathbf{Set}]$, as required. $\square$

6.4.2 Finding theories classified by a given presheaf topos

The following theorem provides a method for constructing theories of presheaf type whose categories of finitely presentable models are equivalent, up to Cauchy-completion, to a given small category of structures.

Theorem 6.29. *Let $\mathbb{T}$ be a theory of presheaf type and $\mathcal{A}$ a full subcategory of $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$. Then the $\mathcal{A}$ -completion $\mathbb{T}'$ of $\mathbb{T}$ (i.e., the set of all geometric sequents over the signature of $\mathbb{T}$ which are valid in all models in $\mathcal{A}$) is of presheaf type classified by the topos $[\mathcal{A}, \mathbf{Set}]$; in particular, every finitely presentable $\mathbb{T}'$ -model is a retract of a model in $\mathcal{A}$.*

Proof Since every model in $\mathcal{A}$ is finitely presentable as a $\mathbb{T}$ -model, we have a geometric inclusion $i : [\mathcal{A}, \mathbf{Set}] \hookrightarrow \mathbf{Set}[\mathbb{T}] \simeq [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ induced by the canonical inclusion $\mathcal{A} \hookrightarrow \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$. This subtopos corresponds, by the duality theorem of [9], to a quotient $\mathbb{T}'$ of $\mathbb{T}$ classified by the topos $[\mathcal{A}, \mathbf{Set}]$, which can be characterized as the collection of all geometric sequents which hold in every model in $\mathcal{A}$, that is, as the $\mathcal{A}$ -completion $\mathbb{T}'$ of $\mathbb{T}$ (recall that theories of presheaf type have enough set-based models). Therefore $\mathbb{T}'$ is of presheaf type classified by the topos $[\mathcal{A}, \mathbf{Set}]$; but the finitely presentable $\mathbb{T}'$ -models are the finitely presentable objects of $\text{Ind-}\mathcal{A}$, that is the retracts of objects of $\mathcal{A}$ in $\text{Ind-}\mathcal{A} \simeq \mathbb{T}'\text{-mod}(\mathbf{Set})$. This completes the proof of the theorem. $\square$

Remarks 6.30. (a) Theorem 6.29 is a generalization of Joyal and Wraith's recognition theorem (Theorem 1.1 [2]), as well as of Proposition 5.3 [22]. Indeed, the former theorem can be obtained as the particular case of Theorem 6.29 when $\mathbb{T}'$ has enough set-based models and every set-based model of $\mathbb{T}'$ is a filtered colimit of finitely presentable models of $\mathbb{T}'$ which are also finitely presentable as $\mathbb{T}$ -models, while Proposition 5.3 [22] can be obtained as the specialization of this latter theorem to the case when $\mathbb{T}'$ is a disjunctive theory and $\mathbb{T}$ is a cartesian theory (in view of the fact, proved in [22], that the category of set-based models of a disjunctive

theory is multiply finitely presentable and hence that every set-based model of a disjunctive theory can be expressed as a filtered colimit of finitely presentable models of it).

- (b) It is worth to compare Theorem 6.29 with Theorem 5.1. If $\mathbb{T}'$ is a quotient of a theory of presheaf type $\mathbb{T}$ and $\mathcal{K}$ is a small category of set-based $\mathbb{T}'$ -models such that every model in $\mathcal{K}$ is finitely presentable as a $\mathbb{T}$ -model then, in order to conclude that $\mathbb{T}'$ is of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$, we can either verify conditions (i) and (ii) of Theorem 5.1 or to verify that the models in $\mathcal{K}$ are jointly conservative for $\mathbb{T}'$; in fact, as observed in Remark 5.2(c), conditions (i) and (ii) of Theorem 5.1 together imply that the models in $\mathcal{K}$ are jointly conservative for $\mathbb{T}'$.

Corollary 6.31. *Assuming the axiom of choice, every coherent theory (or, more generally, any theory in a countable fragment of geometric logic as in the hypotheses of Theorem 5.1.7 [27]) $\mathbb{T}$ over a finite relational signature whose axioms do not contain existential quantifications is of presheaf type.*

Proof Since the axioms of $\mathbb{T}$ do not contain quantifications, every substructure of a model of $\mathbb{T}$ is a model of $\mathbb{T}$. Moreover, since the signature of $\mathbb{T}$ is relational, every finitely generated substructure over the signature of $\mathbb{T}$ contains only a finite number of elements besides the constants. Therefore every finitely presentable model of $\mathbb{T}$ contains only a finite number of elements besides the constants (as any model of $\mathbb{T}$ is a filtered union of its finite submodels); on the other hand, since the signature of $\mathbb{T}$ is finite, every such model is finitely presentable as a model of the empty theory over the signature of $\mathbb{T}$ (cf. Theorem 6.4 [6]). Therefore condition (iii) of Theorem 5.1 is satisfied (cf. Proposition 5.11(i)). Now, under the axiom of choice, every theory satisfying the hypotheses of Theorem 5.1.7 [27], in particular any coherent theory, has enough models, whence our thesis follows from Theorem 6.29. $\square$

It is nonetheless important, from a constructive viewpoint, to be able to prove that a theory is of presheaf type without invoking the axiom of choice. Theorem 5.1 allows us to do so in a great variety of cases.

Theorem 6.29 shows that a good first step in constructing a geometric theory classified by a presheaf topos $[\mathcal{K}, \mathbf{Set}]$ consists in finding a theory of presheaf type $\mathbb{T}$ such that the category $\mathcal{K}$ can be identified as a full subcategory of the category of finitely presentable models of $\mathbb{T}$. Indeed, under these hypotheses, Theorem 6.29 ensures the existence of a quotient $\mathbb{T}_{\mathcal{K}}$ of $\mathbb{T}$ classified by the topos $[\mathcal{K}, \mathbf{Set}]$, which can be characterized as the theory consisting of all the geometric sequents over the signature of $\mathbb{T}$ which are valid in every model in $\mathcal{K}$. In most cases, if one has a natural candidate $\mathbb{S}$ for a theory classified by the topos $[\mathcal{K}, \mathbf{Set}]$, the theory $\mathbb{T}$ can be chosen to be the Horn part of $\mathbb{S}$ or the cartesianization $\mathbb{S}_c$ of $\mathbb{S}$.

Of course, the abstract characterization of $\mathbb{T}_{\mathcal{K}}$ as the theory consisting of all the geometric sequents over the signature of $\mathbb{T}$ which are valid in every model in $\mathcal{K}$ is not very useful in specific contexts, where one looks for an axiomatization of $\mathbb{T}_{\mathcal{K}}$ as simple and ‘economical’ as possible. To this end, we observe that if every M in $\mathcal{K}$ is strongly finitely presented as a model of $\mathbb{T}$ as well as finitely generated (in the sense that for any sort A over the signature of $\mathbb{T}$ the elements of the set MA are precisely given by the interpretations in M of terms $t^A(\vec{x})$ (or more generally of $\mathbb{T}$ -provably functional predicates) over the signature of $\mathbb{T}$, where $\vec{x}$ are the generators of M as strongly finitely presented model of $\mathbb{T}$) then we dispose of an explicit axiomatization of the theory $\mathbb{T}_{\mathcal{K}}$, as given by the following

Theorem 6.32. *Let $\mathbb{T}$ be a geometric theory over a signature Σ and $\mathcal{K}$ a full subcategory of the category set-based $\mathbb{T}$ -models such that every $\mathbb{T}$ -model in $\mathcal{K}$ is both strongly finitely presentable and finitely generated (with respect to the same generators). Then the following sequents (where we denote by $\mathcal{P}$ the set of geometric formulae over Σ which strongly present a $\mathbb{T}$ -model in $\mathcal{K}$), added to the axioms of $\mathbb{T}$, yield an axiomatization of a quotient of $\mathbb{T}$ classified by the topos $[\mathcal{K}, \mathbf{Set}]$ via a Morita-equivalence induced by the canonical geometric morphism $[\mathcal{K}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$.*

In particular, if the theory $\mathbb{T}$ is of presheaf type (whence every finitely presentable $\mathbb{T}$ -model is strongly finitely presentable) and all the models in the full subcategory $\mathcal{K}$ of $\mathbb{T}\text{-mod}(\mathbf{Set})$ are finitely presented and finitely generated (with respect to the same generators), the following sequents yield an axiomatization of the theory $\mathbb{T}_{\mathcal{K}}$ defined above:

(i) *The sequent*

$$(\top \vdash_{\square} \bigvee_{\phi(\vec{x}) \in \mathcal{P}} (\exists \vec{x}) \phi(\vec{x}));$$

(ii) *For any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, the sequent*

$$(\phi(\vec{x}) \wedge \psi(\vec{y}) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, t_1^{A_1}(\vec{z}), \dots, t_n^{A_n}(\vec{z}) \\ s_1^{B_1}(\vec{z}), \dots, s_m^{B_m}(\vec{z})}} (\exists \vec{z}) (\chi(\vec{z}) \wedge \bigwedge_{\substack{i \in \{1, \dots, n\}, \\ j \in \{1, \dots, m\}}} (x_i = t_i(\vec{z}) \wedge y_j = s_j(\vec{z}))))),$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $t_1^{A_1}(\vec{z}), \dots, t_n^{A_n}(\vec{z})$ and $s_1^{B_1}(\vec{z}), \dots, s_m^{B_m}(\vec{z})$ whose output sorts are respectively $A_1, \dots, A_n, B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(t_1^{A_1}(\vec{\xi}), \dots, t_n^{A_n}(\vec{\xi})) \in [[\vec{x} \cdot \phi]]_{M_{\{\vec{z}, \chi\}}}$ and $(s_1^{B_1}(\vec{\xi}), \dots, s_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$;

- (iii) For any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, and any terms $t_1^{A_1}(\vec{y}), s_1^{A_1}(\vec{y}), \dots, t_n^{A_n}(\vec{y}), s_n^{A_n}(\vec{y})$ whose output sorts are respectively $A_1, \dots, A_n$, the sequent

$$\left(\bigwedge_{i \in \{1, \dots, n\}} (t_i(\vec{y}) = s_i(\vec{y})) \wedge \phi(t_1/x_1, \dots, t_n/x_n) \wedge \phi(s_1/x_1, \dots, s_n/x_n) \wedge \psi(\vec{y}) \right) \\ \vdash_{\vec{y}} \bigvee_{\chi(\vec{z}) \in \mathcal{P}, u_1^{B_1}(\vec{z}), \dots, u_m^{B_m}(\vec{z})} ((\exists \vec{z})(\chi(\vec{z}) \wedge \bigwedge_{j \in \{1, \dots, m\}} (y_j = u_j(\vec{z}))),$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $u_1^{B_1}(\vec{z}), \dots, u_m^{B_m}(\vec{z})$ whose output sorts are respectively $B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(u_1^{B_1}(\vec{\xi}), \dots, u_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$ and $t_i(u_1(\vec{\xi}), \dots, u_m(\vec{\xi})) = s_i(u_1(\vec{\xi}), \dots, u_m(\vec{\xi}))$ in $M_{\{\vec{z}, \chi\}}$ for all $i \in \{1, \dots, n\}$;

- (iv) For any sort A over Σ , the sequent

$$(\top \vdash_{x_A} \bigvee_{\chi(\vec{z}) \in \mathcal{P}, t^A(\vec{z})} (\exists \vec{z})(\chi(\vec{z}) \wedge x = t(\vec{z}))),$$

where the the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the terms $t^A(\vec{z})$ whose output sort is A ;

- (v) For any sort A over Σ , any formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ in $\mathcal{P}$, where $\vec{x} = (x_1^{A_1}, \dots, x_n^{A_n})$ and $\vec{y} = (y_1^{B_1}, \dots, y_m^{B_m})$, and any terms $t^A(\vec{x})$ and $s^A(\vec{y})$, the sequent

$$(\phi(\vec{x}) \wedge \psi(\vec{y}) \wedge t(\vec{x}) = s(\vec{y})) \vdash_{\vec{x}, \vec{y}} \bigvee_{\substack{\chi(\vec{z}) \in \mathcal{P}, p_1^{A_1}(\vec{z}), \dots, p_n^{A_n}(\vec{z}) \\ q_1^{B_1}(\vec{z}), \dots, q_m^{B_m}(\vec{z})}} (\exists \vec{z})(\chi(\vec{z}) \wedge \\ \wedge \bigwedge_{\substack{i \in \{1, \dots, n\}, \\ j \in \{1, \dots, m\}}} (x_i = p_i(\vec{z}) \wedge y_j = q_j(\vec{z}))),$$

where the disjunction is taken over all the formulae $\chi(\vec{z})$ in $\mathcal{P}$ and all the sequences of terms $p_1^{A_1}(\vec{z}), \dots, p_n^{A_n}(\vec{z})$ and $q_1^{B_1}(\vec{z}), \dots, q_m^{B_m}(\vec{z})$ whose output sorts are respectively $A_1, \dots, A_n, B_1, \dots, B_m$ and such that, denoting by $\vec{\xi}$ the set of generators of the model $M_{\{\vec{z}, \chi\}}$ (strongly) finitely presented by the formula $\chi(\vec{z})$, $(p_1^{A_1}(\vec{\xi}), \dots, p_n^{A_n}(\vec{\xi})) \in [[\vec{x} \cdot \phi]]_{M_{\{\vec{z}, \chi\}}}$ and $(q_1^{B_1}(\vec{\xi}), \dots, q_m^{B_m}(\vec{\xi})) \in [[\vec{y} \cdot \psi]]_{M_{\{\vec{z}, \chi\}}}$ and $t(p_1(\vec{\xi}), \dots, p_n(\vec{\xi})) = s(q_1(\vec{\xi}), \dots, q_m(\vec{\xi}))$ in $M_{\{\vec{z}, \chi\}}$.

Proof Let $\mathbb{R}$ be the geometric theory obtained from $\mathbb{T}$ by adding all the sequents specified above. The objects in $\mathcal{K}$ are clearly models of $\mathbb{R}$ (cf. Remarks 5.6(a) and 5.10). From the fact that they are strongly finitely presentable it follows by Proposition 5.11(ii) that condition (iii) of Theorem 5.1 is satisfied by the theory $\mathbb{R}$ with respect to the category $\mathcal{K}$. By Theorems 5.3 and 5.9 and Remarks 5.6(a) and 5.10, $\mathbb{R}$ also satisfies conditions (i) and (ii) of Theorem 5.1 with respect to $\mathcal{K}$. Theorem 5.1 thus implies that $\mathbb{R}$ is of presheaf type classified by the topos $[\mathcal{K}, \mathbf{Set}]$. Moreover, it can be readily seen that the resulting Morita-equivalence is induced by the canonical geometric morphism from $[\mathcal{K}, \mathbf{Set}]$ to the classifying topos for $\mathbb{T}$; this ensures in particular that if $\mathbb{T}$ is of presheaf type then $\mathbb{R}$ is equal to the theory $\mathbb{T}_{\mathcal{K}}$. $\square$

6.5 Injectivizations of theories

For any geometric theory $\mathbb{T}$, we can slightly modify its syntax so to obtain a geometric theory whose models in $\mathbf{Set}$ are the same as those of $\mathbb{T}$ and whose homomorphisms between them are precisely the sortwise injective $\mathbb{T}$ -model homomorphisms.

This construction is useful in many contexts. For instance, in [9] we showed that if the category $\mathbf{f.p.T-mod}(\mathbf{Set})$ of finitely presentable models of a theory of presheaf type $\mathbb{T}$ satisfies the amalgamation property then the quotient of $\mathbb{T}$ corresponding to the subtopos $\mathbf{Sh}(\mathbf{f.p.T-mod}(\mathbf{Set})^{\text{op}}, J_{at})$ of $[\mathbf{f.p.T-mod}(\mathbf{Set}), \mathbf{Set}]$ (where J_{at} is the atomic topology) via the duality theorem of [9] axiomatizes the homogeneous $\mathbb{T}$ -models (in the sense of [5]) in any Grothendieck topos. Now, the notion of homogeneous $\mathbb{T}$ -model, which is strictly related to the notion of weakly homogeneous model considered in classical Model Theory (cf. [20]), is mostly interesting when the arrows of the category $\mathbf{T-mod}(\mathbf{Set})$ are all monic; indeed, as shown in [6], a necessary condition for $\mathbb{T}$ to admit an associated ‘concrete’ Galois theory, is that all the arrows in $\mathbf{f.p.T-mod}(\mathbf{Set})$ should be strict monomorphisms.

This motivates the following formal definition.

Definition 6.33. Let $\mathbb{T}$ be a geometric theory over a signature Σ . The *injectivization* $\mathbb{T}_m$ of $\mathbb{T}$ is the geometric theory obtained from $\mathbb{T}$ by adding a binary predicate $D_A \multimap A, A$ for each sort A over Σ and the coherent sequents

$$(D_A(x^A, y^A) \wedge x^A = y^A) \vdash_{x^A, y^A} \perp$$

and

$$(\top \vdash_{x^A, y^A} D_A(x^A, y^A) \vee x^A = y^A) .$$

The models of $\mathbb{T}_m$ in an arbitrary topos $\mathcal{E}$ coincide with the models M of $\mathbb{T}$ in $\mathcal{E}$ which are sortwise decidable, in the sense that for every sort A over the signature of $\mathbb{T}$ the object MA of $\mathcal{E}$ is *decidable*, i.e. the diagonal subobject of MA is complemented (by the interpretation of D_A in M).

As shown by the following lemma, the arrows $M \rightarrow N$ in the category $\mathbb{T}_m\text{-mod}(\mathbf{Set})$ are precisely the $\mathbb{T}$ -model homomorphisms $f : M \rightarrow N$ such that for every sort A over the signature of $\mathbb{T}$, $fA : MA \rightarrow NA$ is a monomorphism in $\mathcal{E}$.

Lemma 6.34. *Let A and B be decidable objects in a topos $\mathcal{E}$ and $f : A \rightarrow B$ an arrow in $\mathcal{E}$. Let $D_A \rightarrowtail A \times A$ and $D_B \rightarrowtail B \times B$ denote respectively the complements of the diagonal subobjects $\delta_A : A \rightarrowtail A \times A$ and $\delta_B : B \rightarrowtail B \times B$. Then f is a monomorphism if and only if $f \times f : A \times A \rightarrow B \times B$ restricts to an arrow $D_A \rightarrow D_B$.*

Proof It is immediate to see that $f : A \rightarrow B$ is a monomorphism if and only if the diagram

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow \delta_A & & \downarrow \delta_B \\ A \times A & \xrightarrow{f \times f} & B \times B, \end{array}$$

is a pullback.

Since pullback functors preserve arbitrary unions and intersections of subobjects in a topos (they having both a left and a right adjoint), we have that $(f \times f)^*(D_B) \cong \neg(f \times f)^*(\delta_B)$. Now, $f \times f : A \times A \rightarrow B \times B$ restricts to an arrow $D_A \rightarrow D_B$ if and only if $D_A \leq (f \times f)^*(D_B)$. But this condition holds if and only if $D_A \cap (f \times f)^*(\delta_B) \cong 0$, i.e. if and only if $(f \times f)^*(\delta_B) \leq \delta_A$, which is equivalent to the condition $(f \times f)^*(\delta_B) \cong \delta_A$ (as $\delta_A \leq (f \times f)^*(\delta_B)$). $\square$

Several injectivizations of theories of presheaf type have been considered in the literature, e.g. in [26] and in [21]; see also [7] and [6] for some applications of this type of theories in the context of topos-theoretic Galois-type equivalences.

As we shall see below, under certain conditions, the injectivization of a theory of presheaf type is again of presheaf type.

The following proposition is a corollary of Theorem 4.3.

Proposition 6.35. *Let $\mathbb{T}$ be a geometric theory. Then for any $\mathbb{T}$ -models M and N in a Grothendieck topos $\mathcal{E}$ which are sortwise decidable there exists an object $\text{Hom}_{\mathbb{T}_m\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)$ of $\mathcal{E}$ satisfying the following universal property: for any object E of $\mathcal{E}$ we have an equivalence*

$$\text{Hom}_{\mathcal{E}}(E, \text{Hom}_{\mathbb{T}_m\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)) \cong \text{Hom}_{\mathbb{T}_m\text{-mod}(\mathcal{E}/E)}(!_E^*(M), !_E^*(N))$$

natural in $E \in \mathcal{E}$.

Remarks 6.36. (a) For any $\mathcal{E}$, M and N , $\text{Hom}_{\mathbb{T}_m\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)$ embeds canonically as a subobject of $\text{Hom}_{\mathbb{T}\text{-mod}(\mathcal{E})}^{\mathcal{E}}(M, N)$.

- (b) Let $\mathbb{T}$ be a geometric theory over a signature Σ , c a finitely presentable $\mathbb{T}$ -model and M a sortwise decidable model of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$. Then the subobject

$$Hom_{\underline{\mathbb{T}_m\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M) \hookrightarrow Hom_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$$

can be identified with the interpretation of the formula

$$\chi_c := \bigwedge_{\substack{A \text{ sort over } \Sigma, \\ x, y \in cA, x \neq y}} (\pi_A(f(\gamma_{\mathcal{E}}^*(\bar{x})), \pi_A(f(\gamma_{\mathcal{E}}^*(\bar{y})))) \in D_{MA},$$

written in the internal language of the topos $\mathcal{E}$, where

$$\pi_A : Hom_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(M, N) \rightarrow NA^{MA}$$

is the arrow defined in Remark 4.4(b) and $\bar{x}, \bar{y} : 1 \rightarrow c$ are the arrows in **Set** corresponding respectively to the elements x and y of cA . Indeed, a $\mathbb{T}$ -model homomorphism $f : \gamma_{\mathcal{E}}^*(c) \rightarrow M$ is sortwise monic if and only if for every sort A over Σ and any distinct elements $x, y \in cA$, $fA(\gamma_{\mathcal{E}}^*(\bar{x}))$ and $fA(\gamma_{\mathcal{E}}^*(\bar{y}))$ are disjoint, in other words they satisfy the relation D_{MA} .

It follows that if for every finitely presentable $\mathbb{T}$ -model c , the formula χ_c is equivalent to a geometric formula over the signature of $\mathbb{T}_m$, for instance when Σ only contains a finite number of sorts and for any sort A and any finitely presentable $\mathbb{T}$ -model the set cA is finite, then the theory $\mathbb{T}_m$ satisfies condition (iii)(a) of Theorem 5.1 with respect to the category of finitely presentable $\mathbb{T}$ -models if $\mathbb{T}$ does.

- (c) Let $\mathbb{T}$ be a geometric theory over a signature Σ , c a finitely presentable $\mathbb{T}$ -model and M a sortwise decidable model of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$. If c is strongly finitely presented by a geometric formula $\phi(\vec{x})$ over Σ then the subobject $Hom_{\underline{\mathbb{T}_m\text{-mod}_m(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ of $Hom_{\underline{\mathbb{T}\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M) \cong [[\vec{x} \cdot \phi]]_M$ can be identified with the intersection of the interpretations in M of the formulae of the form $\neg(\exists x^A)(\theta_1(\vec{x}, x^A) \wedge \theta_2(\vec{x}, x^A))$, where A is a sort over Σ and θ_1, θ_2 are $\mathbb{T}$ -provably functional formulae from $\{\vec{x} \cdot \phi\}$ to $\{x^A \cdot \top\}$.

Lemma 6.37. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , c a set-based $\mathbb{T}$ -model and M a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$. A generalized element $x : E \rightarrow G_M(c) = Hom_{\underline{\mathbb{T}_m\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^*(c), M)$ can be identified with a Σ -homomorphism $\xi_x : c \rightarrow Hom_{\mathcal{E}}(E, M)$ which is sortwise disjunctive in the sense that for every sort A over Σ , the function $\xi_x A : cA \rightarrow Hom_{\mathcal{E}}(E, MA)$ has the property that for any distinct elements $z, w \in cA$, the arrows $\xi_x A(z), \xi_x A(w) : E \rightarrow MA$ have equalizer zero in $\mathcal{E}$.*

Proof The thesis follows from Proposition 4.5, observing that, by Lemmas 6.41 and 6.43, an arrow $\tau_A : \gamma_{\mathcal{E}/E}^*(cA) \rightarrow !_E^*(MA)$ in $\mathcal{E}/E$ is monic in $\mathcal{E}/E$ if and only if the corresponding arrow $\xi_A : cA \rightarrow \text{Hom}_{\mathcal{E}}(E, MA)$ satisfies the property that for any distinct elements $z, w \in cA$, the arrows $\xi_A(z) : E \rightarrow MA$ and $\xi_A(w) : E \rightarrow MA$ have equalizer zero (notice that two arrows $s, t : (a : A \rightarrow E) \rightarrow (b : B \rightarrow E)$ in $\mathcal{E}/E$ have equalizer zero in $\mathcal{E}/E$ if and only if the arrows $s : A \rightarrow B$ and $t : A \rightarrow B$ have equalizer zero in $\mathcal{E}$). $\square$

Remark 6.38. Suppose that the formula $\{x^A . \top\}$ strongly presents a $\mathbb{T}$ -model F_A . Then a Σ -structure homomorphism $s : F_A \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, corresponding to a generalized element $z : E \rightarrow MA$, is sortwise disjunctive if and only if for any sort B over Σ and any two $\mathbb{T}$ -provably inequivalent $\mathbb{T}$ -provably functional geometric formulae θ_1 and θ_2 from $\{x^A . \top\}$ to $\{x^B . \top\}$, the generalized elements $[[\theta_1]]_M \circ z$ and $[[\theta_2]]_M \circ z$ are disjoint. Indeed, from the proof of Theorem 4.1 we know that that for any sort B and $\mathbb{T}$ -provably functional geometric formula θ from $\{x^A . \top\}$ to $\{x^B . \top\}$, the function $[[\theta]]_{F_A} : F_A A = \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{x^A . \top\}, \{x^A . \top\}) \rightarrow F_A B = \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{x^A . \top\}, \{x^B . \top\})$ coincides with $[\theta] \circ - : \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{x^A . \top\}, \{x^A . \top\}) \rightarrow \text{Hom}_{\mathcal{C}_{\mathbb{T}}}(\{x^A . \top\}, \{x^B . \top\})$, while the generator u_A of F_A is precisely the identity arrow on $\{x^A . \top\}$. Now, it is immediate to see that for any Σ -structure homomorphism $s : F_A \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and any sort B over Σ , the diagram

$$\begin{array}{ccc} F_A A & \xrightarrow{sA} & \text{Hom}_{\mathcal{E}}(E, MA) \\ \downarrow [\theta] \circ - & & \downarrow [[\theta]]_M \circ - \\ F_A B & \xrightarrow{sB} & \text{Hom}_{\mathcal{E}}(E, MB) \end{array}$$

commutes, i.e. $sA(u_A) = sB(\theta)$. From these remarks, our claim immediately follows.

Note that if $\mathbb{T}$ is a universal Horn theory (in the sense of [3]) then we can suppose without loss of generality θ_1 and θ_2 to be functional formulae of the form $x^B = t(x^A)$, where t is a term of type $A \rightarrow B$ over Σ (cf. Remark 4.2(a)).

6.5.1 Condition (iii) of Theorem 5.1 for injectivizations

In this section we shall establish a general result about injectivizations of theories of presheaf type, namely that for any theory of presheaf type $\mathbb{T}$ such that all the monic arrows in the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ are sortwise monic, the injectivization of $\mathbb{T}$ satisfies condition (iii) of Theorem 5.1 with respect to the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$.

Before proving this theorem, we need a series of preliminary results.

Lemma 6.39. *Let $r : R \rightarrowtail A \times A$ a subobject in a Grothendieck topos $\mathcal{E}$, $e : E \rightarrow A \times A$ an arrow in $\mathcal{E}$ and $\{f_i : E_i \rightarrow E \mid i \in I\}$ an epimorphic family in $\mathcal{E}$. If for every $i \in I$ the arrow $e \circ f_i$ factors through r then e factors through r .*

Proof The arrow $\coprod_{i \in I} f_i : \coprod_{i \in I} E_i \rightarrow E$ induced by the universal property of the coproduct is an epimorphism, since by our hypothesis the family $\{f_i : E_i \rightarrow E \mid i \in I\}$ is epimorphic. The factorizations $b_i : E_i \rightarrow R$ of the arrows $e \circ f_i$ through r induce an arrow $b := \coprod_{i \in I} b_i : \coprod_{i \in I} E_i \rightarrow R$, such that $r \circ b = e \circ \coprod_{i \in I} f_i$. Consider the epi-mono factorization $b = h \circ k$ in $\mathcal{E}$ of the arrow b , where $k : \coprod_{i \in I} E_i \rightarrowtail U$ and $h : U \rightarrowtail R$, and the epi-mono factorization $e = m \circ n$ of the arrow e in $\mathcal{E}$, where $n : E \rightarrow T$ and $m : T \rightarrowtail A$. Clearly, the arrow $e \circ \coprod_{i \in I} f_i$ factorizes both as $m \circ (n \circ \coprod_{i \in I} f_i)$ and as $(r \circ h) \circ k$. Now, as $m, r \circ h$ are monic and $n \circ \coprod_{i \in I} f_i, k$ are epic, the uniqueness of the epi-mono factorization of a given arrow in a topos implies that there exists an isomorphism $i : T \cong U$ such that $r \circ h \circ i = m$ and $i \circ n \circ \coprod_{i \in I} f_i = k$. The arrow $h \circ i \circ n$ thus provides a factorization of e through r , as required. $\square$

Lemma 6.40. *Let $\{f_i : A_i \rightarrow B \mid i \in I\}$ be a family of arrows in a Grothendieck topos $\mathcal{E}$. Then the arrow $\coprod_{i \in I} f_i : \coprod_{i \in I} A_i \rightarrow B$ is monic if and only if for every $i \in I$, f_i is monic and for every $i, i' \in I$, either $i = i'$ or the subobjects $f_i : A_i \rightarrowtail B$ and $f_{i'} : A_{i'} \rightarrowtail B$ are disjoint.*

Proof The ‘if’ direction follows from Proposition IV 7.6 [26], while the ‘only if’ one follows from the fact that coproducts in a topos are always disjoint (cf. Corollary IV 10.5 [26]) and the composite of a given monomorphism with two disjoint subobjects yields two disjoint subobjects. $\square$

Remark 6.41. If in the statement of Lemma 6.40 the objects A_i are all equal to the terminal object $1_{\mathcal{E}}$ of $\mathcal{E}$ then the arrows f_i are automatically monic and any two of them are disjoint if and only if their equalizer is zero.

The notation employed in the statements and proofs of the following results is borrowed from section 3.2.

Lemma 6.42. *Let $F : \mathcal{D}^{op} \rightarrow \mathcal{E}$ be a flat functor from a subcategory $\mathcal{D}$ of a small category $\mathcal{C}$ to a Grothendieck topos $\mathcal{E}$ and $x : E \rightarrow \tilde{F}(d)$ a generalized element which factors through $\chi_d^F : F(d) \rightarrow \tilde{F}(d)$ as $\chi_d^F \circ x'$. Then the natural transformation $\alpha_x : \gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{C}d} \rightarrow !_E^* \circ \tilde{F}$ corresponding to x is equal to $\tilde{\alpha}_{x'}$, where $\alpha_{x'}$ is the natural transformation $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{D}d} \rightarrow !_E^* \circ F$ corresponding to x' .*

Proof Straightforward from the results of section 3.2. $\square$

Given a small category $\mathcal{C}$, we shall write $\mathcal{C}_m$ for the category whose objects are the objects of $\mathcal{C}$ and whose arrows are the monic arrows in $\mathcal{C}$ between them. The following results concern extensions $\tilde{F}$ of flat functors F along the embedding $\mathcal{C}_m \hookrightarrow \mathcal{C}$.

Below, for a given flat functor $H : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$ with values in a Grothendieck topos $\mathcal{E}$ and any object E of $\mathcal{E}$, we write H_E for the flat functor $!_E^* \circ H : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}/E$.

Proposition 6.43. *Let $\mathcal{C}$ be a small category and and $F : \mathcal{C}_m^{\text{op}} \rightarrow \mathcal{E}$ a flat functor. Then for any object $c \in \mathcal{C}$ and any generalized element $x : E \rightarrow \tilde{F}(c)$, x factors through $\chi_c^F : F(c) \rightarrow \tilde{F}(c)$ if and only if the corresponding natural transformation $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{C}} c \rightarrow !_E^* \circ F$ is pointwise monic.*

Proof The ‘only if’ direction follows at once from Lemma 6.42 and Theorem 6.8. To prove the ‘if’ one, thanks to the localization technique, we can suppose without loss of generality $E = 1_{\mathcal{E}}$.

Suppose that the natural transformation $\alpha_x : \gamma_{\mathcal{E}}^* \circ y_{\mathcal{C}} c \rightarrow \tilde{F}$ corresponding to the generalized element $x : 1 \rightarrow \tilde{F}(c)$ is pointwise monic. We want to prove that x factors through $\chi_c^F : F(c) \rightarrow \tilde{F}(c)$. Recall that $\alpha_x(d) :$

$\coprod_{f \in \text{Hom}_{\mathcal{C}}(d, c)} 1_{\mathcal{E}} \rightarrow \tilde{F}(c)$ is defined as the arrow which sends the component of f to the generalized element $\tilde{F}(f) \circ x : 1 \rightarrow \tilde{F}(d)$.

By Lemma 6.40, if α_x is pointwise monic then for any object $d \in \mathcal{C}$ and arrows $f, g : d \rightarrow c$ in $\mathcal{C}$, either $f = g$ or the equalizer of $\tilde{F}(f) \circ x$ and $\tilde{F}(g) \circ x$ is zero. Consider the pullbacks of the jointly epimorphic arrows $\kappa_{(a, z)} : F(a) \rightarrow \tilde{F}(c)$ along the arrow $x : 1_{\mathcal{E}} \rightarrow \tilde{F}(c)$:

$$\begin{array}{ccc} E_{(a, z)} & \xrightarrow{h_{(a, z)}} & F(a) \\ \downarrow e_{(a, z)} & & \downarrow \kappa_{(a, z)} \\ 1 & \xrightarrow{x} & \tilde{F}(c) \end{array}$$

We shall prove that the epimorphic family $\{e_{(a, z)} : E_{(a, z)} \rightarrow 1_{\mathcal{E}} \mid (a, z) \in \mathcal{A}_c\}$ satisfies the condition that for any $(a, z) \in \mathcal{A}_c$, the composite arrow $x \circ e_{(a, z)}$ factors through $\chi_c^F : F(c) \rightarrow \tilde{F}(c)$; this will imply our thesis by Lemma 6.39. As if $E_{(a, z)} \cong 0$ then $x \circ e_{(a, z)}$ factors through $\chi_c^F : F(c) \rightarrow \tilde{F}(c)$, we can suppose $E_{(a, z)} \not\cong 0$. Under this hypothesis, the arrow $z : c \rightarrow a$ is monic in $\mathcal{C}$. Indeed, for any two arrows $f, g : b \rightarrow c$ such that $z \circ f = z \circ g$, $\tilde{F}(f) \circ x \circ e_{(a, z)} = \tilde{F}(f) \circ \kappa_{(a, z)} \circ h_{(a, z)} = \tilde{F}(f) \circ \tilde{F}(z) \circ \chi_a^F \circ h_{(a, z)} = \tilde{F}(g) \circ \tilde{F}(z) \circ \chi_a^F \circ h_{(a, z)} = \tilde{F}(g) \circ \kappa_{(a, z)} \circ h_{(a, z)} = \tilde{F}(g) \circ x \circ e_{(a, z)}$, whence either $f = g$ or the the equalizer of $\tilde{F}(f) \circ x$ and $\tilde{F}(g) \circ x$ is zero. But if the equalizer of $\tilde{F}(f) \circ x$ and $\tilde{F}(g) \circ x$ were zero then $E_{(a, z)}$ would also be isomorphic to zero (since it would admit an arrow to zero), contrary to our hypothesis; so $f = g$, as required. $\square$

Remark 6.44. By Proposition 6.35, Proposition 6.43 can be reformulated as follows: for any object $c \in \mathcal{C}$, $F(c) \cong \text{Hom}_{\mathbb{T}_{\mathcal{C}_m\text{-mod}(\mathcal{E})}}^{\mathcal{E}}(\gamma_{\mathcal{E}}^* \circ y_{\mathcal{C}}c, \tilde{F})$, where $\mathbb{T}_{\mathcal{C}_m}$ is the geometric theory of flat functors on $\mathcal{C}_m^{\text{op}}$.

Theorem 6.45. *Let $\mathcal{C}$ be a small category and $\mathbf{Flat}_m(\mathcal{C}^{\text{op}}, \mathcal{E})$ the subcategory of $\mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E})$ whose objects are the same as $\mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E})$ and whose arrows are the natural transformations between them which are pointwise monic. Then the extension functor*

$$\xi_{\mathcal{E}} : \mathbf{Flat}(\mathcal{C}_m^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\mathcal{C}^{\text{op}}, \mathcal{E}),$$

along the embedding $\mathcal{C}_m \hookrightarrow \mathcal{C}$, which by Theorem 6.8 takes values into $\mathbf{Flat}_m(\mathcal{C}^{\text{op}}, \mathcal{E})$, is full on this latter category.

Proof Let F, G be flat functors $\mathcal{C}_m^{\text{op}} \rightarrow \mathcal{E}$ with values in a Grothendieck topos $\mathcal{E}$, and let $\beta : \tilde{F} \rightarrow \tilde{G}$ be a pointwise monic natural transformation between them. We want to prove that there exists a natural transformation $\alpha : F \rightarrow G$ such that $\beta = \tilde{\alpha}$. It suffices to show that for any $c \in \mathcal{C}$, $\beta(c) : \tilde{F}(c) \rightarrow \tilde{G}(c)$ restricts (along the arrows $\chi_c^F : F(c) \rightarrow \tilde{F}(c)$ and $\chi_c^G : G(c) \rightarrow \tilde{G}(c)$) to an arrow $F(c) \rightarrow G(c)$. To this end, we define a function $\gamma_E : \text{Hom}_{\mathcal{E}}(E, F(c)) \rightarrow \text{Hom}_{\mathcal{E}}(E, G(c))$ natural in $E \in \mathcal{E}$. By Remark 3.2(iii), the set $\text{Hom}_{\mathcal{E}}(E, F(c))$ (resp. the set $\text{Hom}_{\mathcal{E}}(E, G(c))$) can be identified with the set of arrows $E \rightarrow \tilde{F}(c)$ (resp. with the set of arrows $E \rightarrow \tilde{G}(c)$) which factor through χ_c^F (resp. through χ_c^G). By Proposition 6.43, for any flat functor $H : \mathcal{C}^{\text{op}} \rightarrow \mathcal{E}$, the arrows $E \rightarrow \tilde{H}(c)$ which factor through χ_c^H correspond precisely to the natural transformations $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{C}}c \rightarrow \tilde{H}_E$ which are pointwise monic. Now, since β is pointwise monic then $!_E^* \circ \beta : \tilde{F}_E \rightarrow \tilde{G}_E$ is also pointwise monic (since the functor $!_E^*$ preserves monomorphisms, it being the inverse image of a geometric morphism); hence any generalized element $E \rightarrow \tilde{F}(c)$ which factors through χ_c^F gives rise, by composition with $!_E^* \circ \beta$ of the corresponding, pointwise monic, natural transformation $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{C}}c \rightarrow \tilde{F}_E$, to a pointwise monic natural transformation $\gamma_{\mathcal{E}/E}^* \circ y_{\mathcal{C}}c \rightarrow \tilde{G}_E$, that is to a generalized element $E \rightarrow \tilde{G}(c)$ which factors through χ_c^G . But this generalized element is precisely $\beta_c \circ x$. So $\beta(c) : \tilde{F}(c) \rightarrow \tilde{G}(c)$ restricts to an arrow $F(c) \rightarrow G(c)$, as required. $\square$

We can now prove the following

Theorem 6.46. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ . Then the injectivization of $\mathbb{T}$ satisfies condition (iii)(b)-(1) of Theorem 5.1 with respect to the category $f.p.\mathbb{T}\text{-mod}(\mathbf{Set})_m$, and condition (iii)(b)-(2) with respect to the same category if every monic arrow in $f.p.\mathbb{T}\text{-mod}(\mathbf{Set})$ is sortwise monic (for instance, by Proposition 6.47, if for every sort A over Σ there exists the free $\mathbb{T}$ -model on A).*

Proof By Theorem 6.7, the composite functor

$$\xi_{\mathcal{E}} : \mathbf{Flat}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})_m^{\text{op}}, \mathcal{E}) \rightarrow \mathbf{Flat}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E}),$$

takes values in the subcategory $\mathbb{T}_m\text{-mod}(\mathcal{E})$ of $\mathbb{T}\text{-mod}(\mathcal{E})$. The functor $\xi_{\mathcal{E}}$ is faithful by Proposition 3.2(iii) and full on $\mathbb{T}_m\text{-mod}(\mathcal{E})$ by Theorem 6.45 in view of the fact that for any sortwise monic $\mathbb{T}$ -model homomorphism $f : M \rightarrow N$ in $\mathcal{E}$, the natural transformation $\alpha_f : F_M \rightarrow F_N$ corresponding to it under the Morita-equivalence

$$\mathbf{Flat}(\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, \mathcal{E}) \simeq \mathbb{T}\text{-mod}(\mathcal{E})$$

for $\mathbb{T}$ is pointwise monic. This latter fact can be proved as follows. For any object D of the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$, the value of α_f at D can be identified with the arrow $[[\vec{x} \cdot \phi]]_M \rightarrow [[\vec{x} \cdot \phi]]_N$ canonically induced by f , where $\phi(\vec{x})$ is ‘the’ formula which presents the model D ; therefore if f is sortwise monic then $\alpha_f(D)$ is monic in $\mathcal{E}$, it being the restriction of a monic arrow (namely $fA_1 \times \cdots \times fA_n$, where $\vec{x} = (x^{A_1}, \dots, x^{A_n})$) along two subobjects. $\square$

The following proposition identifies a class of theories of presheaf type $\mathbb{T}$ with the property that the monic arrows of the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ are sortwise monic.

Proposition 6.47. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ in which for every sort A over Σ the formula $\{x^A \cdot \top\}$ presents a $\mathbb{T}$ -model, and let $f : M \rightarrow N$ be a homomorphism of finitely presentable $\mathbb{T}$ -models M and N . Then f is monic as an arrow of $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ (equivalently, as an arrow of $\mathbb{T}\text{-mod}(\mathbf{Set})$) if and only if it is sortwise monic.*

Proof Let A be a sort over Σ . As $\{x^A \cdot \top\}$ presents a $\mathbb{T}$ -model F_A then for any model P of $\mathbb{T}$ in $\mathbf{Set}$ we have an equivalence $\text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, P) \cong PA$ natural in P . In particular we have equivalences $\text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, M) \cong MA$ and $\text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, N) \cong NA$ under which the function $f \circ - : \text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, M) \rightarrow \text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, N)$ corresponds to the function $fA : MA \rightarrow NA$. Now, if f is monic then the function $f \circ - : \text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, M) \rightarrow \text{Hom}_{\mathbb{T}\text{-mod}(\mathbf{Set})}(F_A, N)$ is injective, equivalently $fA : MA \rightarrow NA$ is injective, as required. $\square$

6.5.2 A criterion for injectivizations

In this section we shall establish a result providing a sufficient condition for the injectivization of a theory of presheaf type of a certain form to be again of presheaf type. Before stating it, we need some preliminaries.

The following definition gives a natural topos-theoretic generalization of the standard notion of congruence on a set-based structure.

Definition 6.48. Let Σ be a one-sorted first-order signature and M a Σ -structure in a Grothendieck topos $\mathcal{E}$. An equivalence relation $R \rightrightarrows M \times M$ on M in $\mathcal{E}$ is said to be a *congruence* if for any function symbol f over Σ of arity n , we have a commutative diagram

$$\begin{array}{ccc} R^n & \longrightarrow & (M \times M)^n \cong M^n \times M^n \\ \downarrow & & \downarrow f_M \times f_M \\ R & \longrightarrow & M \times M. \end{array}$$

Proposition 6.49. Let Σ be a one-sorted first-order signature and M a Σ -structure in a Grothendieck topos $\mathcal{E}$. For any congruence R on M there exists a Σ -structure M/R on $\mathcal{E}$ whose underlying object is the quotient in $\mathcal{E}$ of M by the relation R , and a Σ -structure epimorphism $p_R : M \rightarrow M/R$ given by the canonical projection. Conversely, for any Σ -structure epimorphism $q : M \rightarrow N$, the kernel pair R_q of q is a congruence on M such that q is isomorphic to M/R_q .

Proof The proof of the proposition is immediate by using the exactness properties of Grothendieck toposes relating epimorphisms and equivalence relations. $\square$

Proposition 6.50. Let $\mathbb{T}$ be a geometric theory over a one-sorted signature Σ and M a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$. If the axioms of $\mathbb{T}$ are all of the form $(\phi \vdash_{\vec{x}} \psi)$, where ϕ does not contain any conjunctions, then for any congruence R on M the Σ -structure M/R is a $\mathbb{T}$ -model.

Proof The thesis can be easily proved by induction on the structure of geometric formulae over Σ , using the fact that the action of the canonical projection homomorphism $p_R : M \rightarrow M/R$ on subobjects (of powers of M) preserves the top subobject, the natural order on subobjects, image factorizations and arbitrary unions. $\square$

The following lemma shows that one can always perform image factorization of homomorphisms of structures in regular categories.

Lemma 6.51. Let Σ be a first-order signature and $\mathcal{C}$ a regular category. Then any Σ -structure homomorphism $f : M \rightarrow N$ in $\mathcal{C}$ can be factored as $h \circ g$, where $h : N' \rightarrow N$ is a Σ -substructure of N and $g : M \rightarrow N'$ is sortwise a cover.

Proof First, we notice that in any regular category finite products of covers are covers; indeed, composite of covers are covers (this can be easily proved by using the definition of cover as an arrow orthogonal to the class of monomorphisms), and the product of two covers $f \times g$, where $f : A \rightarrow B$

and $g : C \rightarrow D$, is equal to the composite $(1_B \times g) \circ (f \times 1_C)$ of two arrows which are pullbacks of covers.

For every sort A over Σ we set $N'A$ equal to $Im(fA)$, gA equal to the canonical cover $MA \rightarrow Im(fA)$ and hA equal to the canonical subobject $N'A \rightarrow NA$. For any function symbol $\xi : A_1, \dots, A_n \rightarrow B$ over Σ , we set $N'\xi$ equal to the restriction $N'A_1 \times \dots \times N'A_n \rightarrow N'B$ of $N\xi : NA_1 \times \dots \times NA_n \rightarrow NB$. This restriction actually exists (and is unique) since f is a Σ -structure homomorphism and $gA_1 \times \dots \times gA_n$ is a cover. For any relation symbol R over Σ of type $A_1, \dots, A_n$, we set $N'R$ equal to the intersection of NR with the canonical subobject $N'A_1 \times \dots \times N'A_n \rightarrow NA_1 \times \dots \times NA_n$. It is clear that $f = h \circ g$, that g is sortwise a cover and that h is a substructure homomorphism, as required. $\square$

The following proposition, giving an explicit characterization of decidable objects in terms of their generalized elements, was stated in [26] as Exercise VIII.8(a).

Proposition 6.52. *Let $\mathcal{E}$ be a cocomplete (in particular, a Grothendieck) topos and A an object of $\mathcal{E}$. Then A is decidable if and only if for any generalized elements $x, y : E \rightarrow A$ there exists an epimorphic family (possibly consisting of just two elements) $\{e_i : E_i \rightarrow E \mid i \in I\}$ such that for any $i \in I$, either $x \circ e_i = y \circ e_i$ or the equalizer of $x \circ e_i$ and $y \circ e_i$ is zero.*

Proof Let us suppose that A is decidable. Let $p : P \rightarrow A \times A$ be the complement of the diagonal subobject $\Delta : A \rightarrow A \times A$. Consider the pullback of $\langle x, y \rangle : E \rightarrow A \times A$ along p :

$$\begin{array}{ccc} E' & \xrightarrow{u} & P \\ \downarrow s & & \downarrow p \\ E & \xrightarrow{\langle x, y \rangle} & A \times A \end{array}$$

The equalizer $i : R \rightarrow E'$ of $x \circ s$ and $y \circ s$ is zero; indeed, by definition of P , the diagram

$$\begin{array}{ccc} 0 & \longrightarrow & A \\ \downarrow & & \downarrow \Delta \\ P & \xrightarrow{p} & A \times A \end{array}$$

is a pullback, and the arrows $z := x \circ s \circ i = y \circ s \circ i : R \rightarrow A$ and $u \circ i : R \rightarrow P$ satisfy the condition $\Delta \circ z = p \circ u \circ i$.

Let us denote by $t : E'' \rightarrow E$ the pullback of the subobject $\Delta : A \rightarrow A \times A$ along $\langle x, y \rangle$. The arrows $s : E' \rightarrow E$ and $t : E'' \rightarrow E$ are jointly epimorphic, they being the pullbacks of arrows which are jointly epimorphic, namely Δ and p .

Therefore, by setting $I = \{0, 1\}$, $E_0 = E'$, $E_1 = E''$, $e_0 : E_0 \rightarrow E$ equal to s and $e_1 : E_1 \rightarrow E$ equal to t , we have that the family $\{e_i : E_i \rightarrow E \mid i \in I\}$ satisfies the condition in the statement of the proposition.

Conversely, let us suppose that the condition in the statement of the proposition is satisfied. To prove that A is decidable, we have to show that the subobject $\Delta \vee \neg\Delta \rightarrow A \times A$ is an isomorphism, in other words that any subobject $\langle x, y \rangle : E \rightarrow A \times A$ factors through it.

Notice that for any generalized elements $x', y' : E' \rightarrow A$, the arrow $\langle x', y' \rangle : E' \rightarrow A \times A$ factors through $\neg\Delta \rightarrow A \times A$ if and only if the equalizer of x' and y' is zero. Indeed, $\langle x', y' \rangle$ factors through $\neg\Delta$ if and only if its image does, and by definition of $\neg\Delta$ this holds if and only if the pullback of it (or equivalently, of $\langle x', y' \rangle$) along Δ is zero, i.e. if and only if the equalizer of x' and y' is zero.

Now, by our hypotheses, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ such that for any $i \in I$, either $x \circ e_i = y \circ e_i$ or the equalizer of $x \circ e_i$ and $y \circ e_i$ is zero. By Lemma 6.39, to prove that $\langle x, y \rangle$ factors through $\Delta \vee \neg\Delta \rightarrow A \times A$ it suffices to prove that for any $i \in I$, $\langle x \circ e_i, y \circ e_i \rangle$ factors through $\Delta \vee \neg\Delta \rightarrow A \times A$. But by our hypothesis, for any given $i \in I$, either $x \circ e_i = y \circ e_i$ (which implies that $\langle x \circ e_i, y \circ e_i \rangle$ factors through $\Delta : A \rightarrow A \times A$) or the equalizer of $x \circ e_i$ and $y \circ e_i$ is zero (which implies, as we have just seen, that $\langle x \circ e_i, y \circ e_i \rangle$ factors through $\neg\Delta \rightarrow A \times A$); therefore for any $i \in I$, $\langle x \circ e_i, y \circ e_i \rangle$ factors through $\Delta \vee \neg\Delta \rightarrow A \times A$, as required. $\square$

Proposition 6.53. *Let Σ be a one-sorted signature, c a finite Σ -structure in **Set**, M a Σ -structure in a Grothendieck topos $\mathcal{E}$ whose underlying object is decidable and E an object of $\mathcal{E}$. Then for any Σ -structure homomorphism $f : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a quotient map $q_i : c \rightarrow c_i$, where c_i is a finite Σ -structure, and a disjunctive Σ -structure homomorphism (in the sense of Lemma 6.37) $J_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $J_i \circ q_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ f$ for all $i \in I$:*

$$\begin{array}{ccc} c & \xrightarrow{f} & \text{Hom}_{\mathcal{E}}(E, M) \\ \downarrow q_i & & \downarrow f \\ c_i & \xrightarrow{J_i} & \text{Hom}_{\mathcal{E}}(E_i, M) . \end{array}$$

Proof Let us suppose that c has n elements $x_1, \dots, x_n$. We know from Proposition 6.52 that for any pair (i, j) where $i, j \in \{1, \dots, n\}$, there exist arrows $e_{(i,j)} : E_{(i,j)} \rightarrow E$ and $e'_{(i,j)} : E'_{(i,j)} \rightarrow E$ such that $e_{(i,j)}$ and $e'_{(i,j)}$ are jointly epimorphic, $f(x_i) \circ e_{(i,j)} = f(x_j) \circ e_{(i,j)}$ and $f(x_i) \circ e'_{(i,j)}, f(x_j) \circ e'_{(i,j)}$ are disjoint. The iterated fibered product of all these epimorphic families (corresponding to the pairs (i, j) such that $i, j \in \{1, \dots, n\}$) thus yields an

epimorphic family $\{u_k : U_k \rightarrow E \mid k \in K\}$ such that for every $k \in K$ there exists a subset $S_k \subseteq \{1, \dots, n\} \times \{1, \dots, n\}$ such that for every $(i, j) \in S_k$, $f(x_i) \circ u_k = f(x_j) \circ u_k$ and for every $(i, j) \notin S_k$, $f(x_i) \circ u_k$ and $f(x_j) \circ u_k$ are disjoint. For each $k \in K$, consider the quotient $q_k : c \rightarrow c_q$ of c by the congruence generated by the pairs of the form (x_i, x_j) for $(i, j) \in S_k$. By definition of this congruence relation, the Σ -structure homomorphism $\text{Hom}_{\mathcal{E}}(e_k, M) \circ f$ factors through q_k , and the resulting factorization is a disjunctive Σ -structure homomorphism. We have thus found a set of data satisfying the condition in the statement of the proposition, as required. $\square$

Proposition 6.54. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ such that the finitely presentable $\mathbb{T}$ -models coincide with the finitely presentable $\mathbb{T}_m$ -models. Suppose that the following condition is satisfied: for any Grothendieck topos $\mathcal{E}$, object E of $\mathcal{E}$ and Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c is a finitely presentable $\mathbb{T}$ -model and M is a sortwise decidable $\mathbb{T}$ -model, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{T}$ -model homomorphism $f_i : c \rightarrow c_i$ of finitely presentable $\mathbb{T}$ -models and a sortwise disjunctive Σ -structure homomorphism $x_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $x_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ x$ for all $i \in I$. Then the injectification $\mathbb{T}_m$ of $\mathbb{T}$ satisfies conditions (i) and (ii) of Theorem 5.1 (with respect to its category of finitely presentable models).*

Proof The proposition represents the particular case of Theorem 6.3 for the faithful interpretation of $\mathbb{T}$ into its injectivization, with $\mathcal{K} = \mathcal{H}$ equal to the category of finitely presentable $\mathbb{T}$ -models. The hypotheses of the theorem are satisfied since the functor $\int a$ is full (cf. Remark 6.4). $\square$

Remark 6.55. Any geometric theory $\mathbb{T}$ such that the finitely presentable $\mathbb{T}$ -models are precisely the finitely generated ones and all its axioms are of the form $(\phi \vdash_{\vec{x}} \psi)$, where ψ is a quantifier-free geometric formula, satisfies the first of the hypotheses of the Corollary. Indeed, every finitely presentable $\mathbb{T}$ -model is clearly finitely presentable as a $\mathbb{T}_m$ -model, and since by Proposition 6.11 the substructures of models of $\mathbb{T}$ are again models of $\mathbb{T}$, every model of $\mathbb{T}$ can be expressed as the directed union of its finitely generated submodels; so the finitely presentable $\mathbb{T}_m$ -models are precisely the finitely generated $\mathbb{T}$ -models.

Corollary 6.56. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ such that the finitely presentable $\mathbb{T}$ -models coincide with the finitely presentable $\mathbb{T}_m$ -models and the monic $\mathbb{T}$ -model homomorphisms in **Set** are precisely the homomorphisms which are sortwise monic. Suppose that the following condition is satisfied: for any Grothendieck topos $\mathcal{E}$, object E of $\mathcal{E}$ and Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c is a finitely presentable $\mathbb{T}$ -model and M is a sortwise decidable $\mathbb{T}$ -model, there exists an epimorphic*

family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{T}$ -model homomorphism $f_i : c \rightarrow c_i$ of finitely presentable $\mathbb{T}$ -models and a sortwise disjunctive Σ -structure homomorphism $x_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $x_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ x$ for all $i \in I$. Then the injectification $\mathbb{T}_m$ of $\mathbb{T}$ is of presheaf type.

Proof By Theorem 6.46, $\mathbb{T}_m$ satisfies condition (iii) of Theorem 5.1 (since all the monic arrows in the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ are injective functions), while Proposition 6.54 ensures that conditions (i) and (ii) of Theorem 5.1 are satisfied. We can thus conclude that the theory $\mathbb{T}_m$ is of presheaf type. $\square$

As a consequence of Corollary 6.56, we obtain the following result.

Corollary 6.57. *Let $\mathbb{T}$ be a geometric theory over a one-sorted signature Σ such that any quotient of a finitely presentable $\mathbb{T}$ -model is a $\mathbb{T}$ -model (for instance, a theory whose axioms are all of the form $(\phi \vdash_{\vec{x}} \psi)$, where ϕ does not contain any conjunctions - cf. Proposition 6.50). Suppose that the finitely presentable $\mathbb{T}$ -models are exactly the finite $\mathbb{T}$ -models and that all the monic arrows in the category $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$ are injective functions. Then if $\mathbb{T}$ is of presheaf type, $\mathbb{T}_m$ is of presheaf type as well.*

Proof Propositions 6.50 and 6.53 ensure that all the conditions of Corollary 6.56 are satisfied. We can thus conclude that $\mathbb{T}_m$ is of presheaf type, as required. $\square$

7 Expansions of theories

7.1 General theory

In this section we introduce the syntactic notion of expansion of a geometric theory, and show that it corresponds in a natural way to having a geometric morphisms between the respective classifying toposes. Further, we characterize the expansions of theories whose corresponding geometric morphisms are localic and hyperconnected. This section has been inspired by section 6.2 of [17], which contains a brief informal discussion of this topic.

Definition 7.1. Let $\mathbb{T}$ be a geometric theory over a signature Σ .

- (i) A geometric *expansion* of $\mathbb{T}$ is a geometric theory obtained from $\mathbb{T}$ by adding sorts, relation or function symbols to the signature Σ and geometric axioms over the resulting extended signature; equivalently, a geometric expansion of $\mathbb{T}$ is a geometric theory $\mathbb{T}'$ over a signature containing Σ such that every axiom of $\mathbb{T}$, regarded as a geometric sequent in the signature of $\mathbb{T}'$, is provable in $\mathbb{T}'$.

- (ii) A geometric expansion $\mathbb{T}'$ of $\mathbb{T}$ is said to be *localic* if no new sorts are added to Σ to obtain the signature of $\mathbb{T}'$.
- (iii) A geometric expansion $\mathbb{T}'$ of $\mathbb{T}$ is said to be *hyperconnected* if no new function or relation symbols which only involve sorts over Σ are added to Σ to form the signature of $\mathbb{T}'$, and for any geometric sequent σ over Σ , σ is provable in $\mathbb{T}'$ if and only if it is provable in $\mathbb{T}$.

Notice that if $\mathbb{T}'$ is an expansion of a geometric theory $\mathbb{T}$ then there is a canonical morphism of sites $(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}}) \rightarrow (\mathcal{C}_{\mathbb{T}'}, J_{\mathbb{T}'})$ inducing a geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$.

We say that a geometric expansion $\mathbb{T}'$ of $\mathbb{T}$ is *faithful* (resp. *fully faithful*) if for every Grothendieck topos $\mathcal{E}$, the induced functor

$$(p_{\mathbb{T}}^{\mathbb{T}'})_{\mathcal{E}} : \mathbb{T}'\text{-mod}(\mathcal{E}) \rightarrow \mathbb{T}\text{-mod}(\mathcal{E})$$

is faithful (resp. full and faithful).

We know from Theorem 9.1 [9] that the inclusion part of the surjection-inclusion factorization of the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$ can be identified with the classifying topos of the quotient of $\mathbb{T}$ consisting of all the sequents over Σ which are provable in the theory $\mathbb{T}'$.

Recall (cf. for instance section A4.6 of [21]) that a geometric morphism $f : \mathcal{F} \rightarrow \mathcal{E}$ is *localic* if every object of $\mathcal{F}$ is a subquotient (i.e. a quotient of a subobject) of an object of the form $f^*(a)$, where a is an object of $\mathcal{E}$; the morphism f is *hyperconnected* if f^* is full and faithful, and its image is closed under subobjects in $\mathcal{F}$.

The following technical lemma will be useful in the sequel.

Lemma 7.2. *Let $f : \mathcal{F} \rightarrow \mathcal{E}$ a geometric morphism between Grothendieck toposes and $\mathcal{C}$ a full subcategory of $\mathcal{E}$ which is separating for $\mathcal{E}$. Suppose that the following conditions are satisfied:*

- (i) $\mathcal{C}$ is closed in $\mathcal{E}$ under finite products;
- (ii) f satisfies the property that the restriction $f^*|_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{F}$ of f^* to $\mathcal{C}$ is full and faithful;
- (iii) for any family of arrows $\mathcal{T}$ in $\mathcal{C}$ with common codomain, if the image of $\mathcal{T}$ under f^* is epimorphic in $\mathcal{F}$ then $\mathcal{T}$ is epimorphic in $\mathcal{E}$, and
- (iv) every subobject in $\mathcal{F}$ of an object of the form $f^*(c)$ where c is an object of $\mathcal{C}$, is, up to isomorphism, of the form $f^*(m)$ (where m is a subobject in $\mathcal{E}$).

Then f is hyperconnected.

Proof We have to prove that, under the specified hypotheses, $f^* : \mathcal{E} \rightarrow \mathcal{F}$ is full and faithful and its image is closed under subobjects in $\mathcal{F}$.

To prove that f^* is faithful, we have to verify that for any arrows $h, k : u \rightarrow v$ in $\mathcal{E}$, $f^*(h) = f^*(k)$ implies $h = k$. Since $\mathcal{C}$ is a separating set for $\mathcal{E}$, we can suppose without loss of generality u to lie in $\mathcal{C}$. Now, $h = k$ if and only if the equalizer $z : w \rightarrow u$ of h and k is an isomorphism. The family of arrows from objects of $\mathcal{C}$ to w is epimorphic, whence, as $f^*(z)$ is an isomorphism (since $f^*(h) = f^*(k)$), the family formed by the composition of these arrows with z is epimorphic on u ; indeed, the members of this latter family all lie in $\mathcal{C}$ and the image of this family under f^* is epimorphic. It follows that z is an epimorphism, equivalently an isomorphism, that is $h = k$, as required.

To prove the fullness of f^* , we have to verify that for any objects a and b of $\mathcal{E}$ and any arrow $s : f^*(a) \rightarrow f^*(b)$ in $\mathcal{F}$, there exists a (unique) arrow $t : a \rightarrow b$ in $\mathcal{E}$ such that $f^*(t) = s$. As $\mathcal{C}$ is a separating set for $\mathcal{E}$ there exist epimorphic families $\{f_i : c_i \rightarrow b \mid i \in I\}$ and $\{g_j : d_j \rightarrow a \mid j \in J\}$ in $\mathcal{E}$ consisting of arrows whose domains lie in $\mathcal{C}$.

For any $i \in I$ and $j \in J$, consider the pullback square

$$\begin{array}{ccc} r_{i,j} & \xrightarrow{p_i} & f^*(c_i) \\ \downarrow q_j & & \downarrow f^*(f_i) \\ f^*(d_j) & \xrightarrow{s \circ f^*(g_j)} & f^*(b) . \end{array}$$

By the universal property of pullbacks, the arrow $\langle p_i, q_j \rangle : r_{i,j} \rightarrow f^*(c_i) \times f^*(d_j) \cong f^*(c_i \times d_j)$ is a monomorphism. Since $c_i \times d_j$ lies in $\mathcal{C}$ by our hypothesis, $r_{i,j}$ lies in the image of f^* and hence can be covered by arrows $h_k^{i,j} : f^*(e_k^{i,j}) \rightarrow r_{i,j}$ (for $k \in K_{i,j}$) where each object $e_k^{i,j}$ lies in $\mathcal{C}$. Consider the arrows $p_i \circ h_k^{i,j} : f^*(e_k^{i,j}) \rightarrow f^*(c_i)$ and $q_j \circ h_k^{i,j} : f^*(e_k^{i,j}) \rightarrow f^*(d_j)$. Since the family $\{f^*(g_j \circ m_k^{i,j}) \mid i \in I, j \in J, k \in K_{i,j}\}$ is epimorphic in $\mathcal{F}$, our hypotheses ensure that the family $\{g_j \circ m_k^{i,j} : e_k^{i,j} \rightarrow A \mid i \in I, j \in J, k \in K_{i,j}\}$ is epimorphic in $\mathcal{E}$. To define an arrow $t : a \rightarrow b$ in $\mathcal{E}$ it is therefore equivalent to specify, for each $i \in I, j \in J$ and $k \in K_{i,j}$, an arrow $v_k^{i,j} : e_k^{i,j} \rightarrow b$ in such a way that the compatibility conditions of Corollary 2.8 are satisfied. Take $v_k^{i,j}$ equal to $f_i \circ l_k^{i,j}$. The compatibility conditions hold since the images of them under the functor f^* do and, as we have proved above, f^* is faithful. Thus we have a unique arrow $t : a \rightarrow b$ in $\mathcal{E}$ such that $t \circ g_j \circ m_k^{i,j} = f_i \circ l_k^{i,j}$. The fact that $f^*(t) = s$ follows at once.

To conclude the proof of the lemma, it remains to show that the image of f^* is closed under subobjects in $\mathcal{F}$. Let a be an object of $\mathcal{E}$ and $k : z \rightarrow f^*(a)$ a subobject in $\mathcal{F}$. Since $\mathcal{C}$ is a separating set for $\mathcal{E}$, the canonical arrow from the coproduct of all objects in $\mathcal{C}$ to a is an epimorphism y ; by pulling back $f^*(y)$ along the monomorphism k we obtain a monomorphism u from z' to $\text{dom}(f^*(y))$ and an epimorphism $e : z' \rightarrow z$. Now, by our hypotheses, the

pullbacks of u along the coproduct arrows belong to the image of f^* ; from the fact that f^* preserves coproducts it thus follows that u itself belongs to the image of f^* , it being of the form $f^*(m)$, where m is a coproduct in $\mathcal{E}$ of subobjects in $\mathcal{C}$. Now, since e is an epimorphism, z is isomorphic to the coequalizer of its kernel pair $r : w \rightrightarrows f^*(b) \times f^*(b) \cong f^*(b \times b)$, where $b = \text{dom}(m)$. Using the coproduct representation of m and the fact that $\mathcal{C}$ is closed under finite products in $\mathcal{E}$, one can prove by considering the pullbacks of r along the images under f^* of the coproduct arrows to $b \times b$, that r lies, up to isomorphism, in the image of f^* ; therefore, as f^* preserves coequalizers, k is isomorphic to a subobject in the image of f^* , as required. $\square$

Theorem 7.3. *Let $\mathbb{T}$ be a geometric theory over a signature Σ and $\mathbb{T}'$ a geometric expansion of $\mathbb{T}$ over a signature Σ' . If $\mathbb{T}'$ is a hyperconnected (resp. a localic) expansion of $\mathbb{T}$ then $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$ is a hyperconnected (resp. a localic) geometric morphism.*

In particular, the hyperconnected-localic factorization of the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$ is given $p_{\mathbb{T}}^{\mathbb{T}'} \circ p_{\mathbb{T}'}^{\mathbb{T}''}$, where $\mathbb{T}''$ is the intermediate expansion of $\mathbb{T}$ obtained by adding to the signature Σ of $\mathbb{T}$ no new sorts and all the relation and function symbols over Σ' which only involve the sorts over Σ , and all the sequents over this extended signature which are provable in $\mathbb{T}'$.

Proof Suppose that $\mathbb{T}'$ is localic over $\mathbb{T}$. We have to prove that every object of $\mathbf{Set}[\mathbb{T}']$ is a quotient of a subobject of an object of the form $f^*(H)$ where H is an object of $\mathbf{Set}[\mathbb{T}]$. Since the signature of $\mathbb{T}'$ does not contain any sorts already present in the signature of $\mathbb{T}$ and the objects of the category $\mathcal{C}_{\mathbb{T}'}$ form a separating set for the topos $\mathbf{Set}[\mathbb{T}']$, we can conclude that the subobjects of the objects of the form $p_{\mathbb{T}}^{\mathbb{T}'}(\{\vec{x} . \top\})$ form a separating set for $\mathbf{Set}[\mathbb{T}']$. Therefore any object of $\mathbf{Set}[\mathbb{T}']$ is a quotient of a coproduct of subobjects of objects of the form $p_{\mathbb{T}}^{\mathbb{T}'}(\{\vec{x} . \top\}) \cong \{\vec{x} . \top\}$; but the fact that $p_{\mathbb{T}}^{\mathbb{T}'*}$ preserves coproducts immediately implies that any such coproduct is a subobject of an object in the image of $p_{\mathbb{T}}^{\mathbb{T}'*}$.

Suppose instead that $\mathbb{T}'$ is hyperconnected over $\mathbb{T}$. We have to prove that $p_{\mathbb{T}}^{\mathbb{T}'*}$ is hyperconnected. It suffices to apply Lemma 7.2 by taking $\mathcal{E}$ equal to $\mathbf{Set}[\mathbb{T}]$ and $\mathcal{C}$ equal to the syntactic category $\mathcal{C}_{\mathbb{T}}$; the fact that the hypotheses of the lemma are satisfied follows immediately from the fact that $\mathbb{T}'$ is hyperconnected over $\mathbb{T}$. $\square$

Theorem 7.4. *Let $\mathbb{T}$ be a geometric theory, $\mathcal{E}$ a Grothendieck topos and M a model of $\mathbb{T}$ in $\mathcal{E}$. Then $\mathcal{E}$ is the classifying topos for $\mathbb{T}$ and M is a universal model for $\mathbb{T}$ if and only if the following conditions are satisfied:*

- (i) *The family $\mathcal{F}$ of objects which can be built from the interpretations in M of the sorts, function symbols and relation symbols over the signature*

of $\mathbb{T}$ by using geometric logic constructions (i.e. the objects given by the interpretations in M of geometric formulae over the signature of $\mathbb{T}$) is separating for $\mathcal{E}$;

- (ii) The model M is conservative for $\mathbb{T}$; that is, for any geometric sequent σ over the signature of $\mathbb{T}$, σ is valid in M if and only if it is provable in $\mathbb{T}$;
- (iii) Any arrow k in $\mathcal{E}$ between objects A and B in the family $\mathcal{F}$ of point (i) is definable; that is, if A (resp. B) is equal to the interpretation of a geometric formula $\phi(\vec{x})$ (resp. $\psi(\vec{y})$) over the signature of $\mathbb{T}$, there exists a $\mathbb{T}$ -provably functional formula θ from $\phi(\vec{x})$ to $\psi(\vec{x})$ such that the interpretation of θ in M is equal to k .

Proof By the universal property of the geometric syntactic category $\mathcal{C}_{\mathbb{T}}$ of $\mathbb{T}$, the $\mathbb{T}$ -model M corresponds to a geometric functor $F_M : \mathcal{C}_{\mathbb{T}} \rightarrow \mathcal{E}$ assigning to each object (or arrow) of $\mathcal{C}_{\mathbb{T}}$ its interpretation in M . Condition (ii) in the statement of the theorem is equivalent to the assertion that the functor F_M is faithful, while condition (iii) is equivalent to saying that F_M is full. Therefore under conditions (ii) and (iii), $\mathcal{C}_{\mathbb{T}}$ embeds as a full subcategory of $\mathcal{E}$. Now, condition (i) ensures that $\mathcal{C}_{\mathbb{T}}$ is dense in $\mathcal{E}$, whence the Comparison Lemma yields an equivalence $\mathcal{E} \simeq \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J)$, where J is the Grothendieck topology on $\mathcal{C}_{\mathbb{T}}$ induced by the canonical topology on $\mathcal{E}$, that is the geometric topology on $\mathcal{C}_{\mathbb{T}}$. By the syntactic construction of classifying toposes and universal models, we can thus conclude that $\mathcal{E}$ is ‘the’ classifying topos for $\mathbb{T}$ and M is a universal model for $\mathbb{T}$. $\square$

As immediate corollaries of Theorem 7.4, one recovers the following known results:

- (i) Let $\mathcal{C}$ be a separating set of objects for a Grothendieck topos $\mathcal{E}$, and $\Sigma_{\mathcal{C}}$ the signature consisting of one sort ‘ c ’ for each object c of $\mathcal{C}$ and one function symbol ‘ f ’ for each arrow f in $\mathcal{E}$ between objects in $\mathcal{C}$. Then there exists a geometric theory $\mathbb{T}$ over the signature $\Sigma_{\mathcal{C}}$ classified by $\mathcal{E}$, whose universal model is given by the ‘tautological’ $\Sigma_{\mathcal{C}}$ -structure in $\mathcal{E}$ (cf. p. 837 [21]);
- (ii) Let B be a pre-bound for $\mathcal{E}$ over $\mathbf{Set}$ (that is, an object such that the subobjects of its finite powers form a generating set for $\mathcal{E}$); then there exists a one-sorted geometric theory $\mathbb{T}$ classified by $\mathcal{E}$ and a universal model for $\mathbb{T}$ whose underlying object is B (cf. Theorem D3.2.5 [21]).

The first result can be obtained from Theorem 7.4 by observing that the theory $\mathbb{T}$ of the tautological $\Sigma_{\mathcal{C}}$ -structure obviously satisfies all the hypotheses of the theorem. That the first two conditions are satisfied is obvious,

while the fact that the third holds can be proved as follows. Since every object of the syntactic category of $\mathbb{T}$ is a subobject of a finite product of objects of the form $\{x^{\lceil c \rceil} . \top\}$ (for $c \in \mathcal{C}$), it is enough to prove that every arrow $k : R \rightarrow c$ in $\mathcal{E}$ having as codomain an object c in $\mathcal{C}$ is definable. Suppose that $R = [[\vec{x} . \phi]]_M$, where the sorts of the variables in $\vec{x} = (x_1, \dots, x_n)$ are respectively $\lceil c'_1 \rceil, \dots, \lceil c'_n \rceil$, and denote by $r : R \rightarrow c'_1 \times \dots \times c'_n$ the corresponding subobject. Since $\mathcal{C}$ is a separating set for $\mathcal{E}$, the family of arrows $\{f_i : c_i \rightarrow R \mid i \in I\}$ from objects of $\mathcal{C}$ to R is epimorphic; hence the geometric formula

$$\bigvee_{i \in I} (\exists z_i^{\lceil c_i \rceil}) (x_1 = \lceil g_1 \rceil(z_i) \wedge \dots \wedge x_n = \lceil g_n \rceil(z_i) \wedge x^{\lceil c \rceil} = \lceil k \circ f_i \rceil(z_i)),$$

where $r \circ f_i = \langle g_1, \dots, g_n \rangle$, is $\mathbb{T}$ -provably functional from $\{\vec{x} . \phi\}$ to $\{x^{\lceil c \rceil} . \top\}$ and its interpretation coincides with k . By Diaconescu's theorem, the theory $\mathbb{T}$ can be explicitly characterized as the theory of flat $J_{\mathcal{E}}^{\mathcal{C}}$ -continuous functors on $\mathcal{C}$, where $J_{\mathcal{E}}^{\mathcal{C}}$ is the Grothendieck topology on $\mathcal{C}$ induced by the canonical topology on $\mathcal{E}$.

The second result can be deduced from Theorem 7.4 by taking $\mathbb{T}$ to be the theory of the tautological structure over the one-sorted signature Σ_B consisting of an n -ary relation symbol $\lceil R \rceil$ for each subobject $R \rightarrowtail B^n$ in $\mathcal{E}$. The fact that $\mathbb{T}$ satisfies the first two conditions of the theorem is obvious, while the validity of the third condition follows from the fact that the graphs of morphisms $B^n \rightarrow B$ in $\mathcal{E}$ are interpretations of $(n+1)$ -relation symbols over Σ_B .

The following theorem provides a converse to Theorem 7.3.

Theorem 7.5. *Let $p : \mathcal{E} \rightarrow \mathbf{Set}[\mathbb{T}]$ be a geometric morphism to the classifying topos of a geometric theory $\mathbb{T}$. Then p is, up to isomorphism, of the form $p_{\mathbb{T}}^{\mathbb{T}'}$ for some geometric expansion $\mathbb{T}'$ of $\mathbb{T}$. If p is hyperconnected (resp. localic) then we can take $\mathbb{T}'$ to be a hyperconnected (resp. localic) expansion of $\mathbb{T}$.*

Proof Choose a triplet $\mathcal{T} = (\mathcal{C}_{ob}, \mathcal{C}_{arr}, \mathcal{C}_{rel})$ consisting of a set $\mathcal{C}_{ob}$ of objects of $\mathcal{E}$, of a set $\mathcal{C}_{arr}$ of arrows in $\mathcal{E}$ from finite products of objects of $\mathcal{C}_{ob}$ to objects of $\mathcal{C}_{ob}$ and of a set $\mathcal{C}_{rel}$ of subobjects of finite products of objects of $\mathcal{C}_{ob}$ with the property that the family of objects of $\mathcal{E}$ which can be built out of objects in $\mathcal{C}_{ob}$, arrows in $\mathcal{C}_{arr}$ and subobjects in $\mathcal{C}_{rel}$ by using geometric logic constructions is separating for $\mathcal{E}$. By definition of Grothendieck topos, such a triplet always exists. Let us define an expansion $\mathbb{T}_{\mathcal{T}}$ of $\mathbb{T}$ as follows: the signature $\Sigma_{\mathbb{T}_{\mathcal{T}}}$ of $\mathbb{T}_{\mathcal{T}}$ is obtained by adding to the signature of $\mathbb{T}$ one sort $\lceil c \rceil$ for each object c in $\mathcal{C}_{ob}$ which is not of the form $f^*(H)$ for some object H of $\mathcal{C}_{\mathbb{T}} \hookrightarrow \mathbf{Set}[\mathbb{T}]$, one function symbol $\lceil f \rceil$ for each arrow f in $\mathcal{C}_{arr}$ whose domain or codomain is not of the form $f^*(H)$ (with the obvious sorts), one relation symbol for each subobject in $\mathcal{C}_{rel}$ (with the obvious sorts, the

ones corresponding to an object of the form $f^*(\{\vec{x} . \phi\})$ being the sorts of the variables $\vec{x}$ and an additional relation symbol $\ulcorner R \urcorner$ for any subobject $R \mapsto c_1 \times \cdots \times c_n$ in $\mathcal{E}$ (where $c_1, \dots, c_n$ are objects in $\mathcal{C}_{ob}$) which cannot be obtained from the data in $\mathcal{T}$ by means of geometric logic constructions (whose sorts are the obvious ones).

Consider the tautological $\mathbb{T}_{\mathcal{T}}$ -structure M in $\mathcal{E}$ obtained by interpreting each sort $\ulcorner c \urcorner$ by the corresponding object c (and similarly for the function and relation symbols added to the signature of $\mathbb{T}$), and each sort A over the signature of $\mathbb{T}$ by the object $f^*(\{x^A . \top\})$ (and similarly for the function and relation symbols over the signature of $\mathbb{T}$). Define $\mathbb{T}'$ to be the theory of M over the signature $\mathbb{T}_{\mathcal{T}}$. The theory $\mathbb{T}'$ satisfies the conditions of Theorem 7.4; the validity of the first two conditions is obvious, while the validity of the third follows from the fact that any subobject of a product of objects in $\mathcal{C}_{ob}$ is definable in $\mathbb{T}'$ and the model M is conservative for $\mathbb{T}'$, whence the formula defining the graph of the given arrow is $\mathbb{T}'$ -provably functional from the formula defining the domain to the formula defining the codomain. Therefore $\mathbb{T}'$ is classified by the topos $\mathcal{E}$ with universal model M . Notice that $\mathbb{T}'$ is an expansion of $\mathbb{T}$. This proves the first part of the theorem.

Suppose now that f is localic. We can define a triplet $\mathcal{T} = (\mathcal{C}_{ob}, \mathcal{C}_{arr}, \mathcal{C}_{rel})$ satisfying the conditions specified above as follows: we set $\mathcal{C}_{ob}$ equal to the set of objects of the form $f^*(H)$ where H is an object of $\mathcal{C}_{\mathbb{T}} \hookrightarrow \mathbf{Set}[\mathbb{T}]$, $\mathcal{C}_{arr}$ equal to the empty set and $\mathcal{C}_{rel}$ equal to the set of all subobjects of (finite products of) objects in $\mathcal{C}_{ob}$. Alternatively, one can take $\mathcal{C}_{ob}$ to be the set of objects of the form $f^*(\{x^A . \top\})$ (where A is any sort over the signature of $\mathbb{T}$), $\mathcal{C}_{arr}$ to be the set of arrows of the form $f^*([\theta])$, where $[\theta]$ is an arrow in $\mathcal{C}_{\mathbb{T}}$, and $\mathcal{C}_{rel}$ to be the set of all subobjects of (finite products of) objects in $\mathcal{C}_{ob}$. In either case, the signature of the resulting expansion will contain no new sorts with respect to the signature of $\mathbb{T}$ and hence it will be localic.

Suppose instead that f is hyperconnected. Given any set of objects $\mathcal{K}$ of $\mathcal{E}$ which, together with the objects of the form $f^*(H)$ (where H is an object of $\mathcal{C}_{\mathbb{T}} \hookrightarrow \mathbf{Set}[\mathbb{T}]$), form a separating set of $\mathcal{E}$, we can define a triplet $\mathcal{T} = (\mathcal{C}_{ob}, \mathcal{C}_{arr}, \mathcal{C}_{rel})$ satisfying the required conditions by setting $\mathcal{C}_{ob} = \mathcal{K}$, $\mathcal{C}_{arr} = \emptyset$ and $\mathcal{C}_{rel} = \emptyset$. Since f is hyperconnected, f^* is full and the image of f^* is closed under subobjects; hence the signature of $\mathbb{T}_{\mathcal{T}}$ does not contain any relation or function symbol only involving the sorts of $\mathbb{T}$. On the other hand, any geometric sequent over the signature of $\mathbb{T}$ is provable in $\mathbb{T}'$ if and only if it is provable in $f^*(M_{\mathbb{T}})$, where $M_{\mathbb{T}}$ is the universal model of $\mathbb{T}$ lying in $\mathbf{Set}[\mathbb{T}]$ (since f^* is full and faithful), i.e. if and only if it is provable in $\mathbb{T}'$. Hence $\mathbb{T}'$ is a hyperconnected expansion of $\mathbb{T}$, as required.

This completes our proof. $\square$

Theorem 7.5 yields, in view of the equivalence between conditions (iii)(b) and (iii)(c) of Theorem 5.1 and Remark 5.2(d), the following reformulation of a particular case of Theorem 6.46.

Theorem 7.6. *Let $\mathbb{T}$ be a theory of presheaf type such that the finitely presentable models of $\mathbb{T}$ coincide with the finitely presentable models of $\mathbb{T}_m$. Then there is a faithful expansion (in the sense of section 7.1) of the injectivization of $\mathbb{T}$ which is classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})_m, \mathbf{Set}]$. If moreover all the monic homomorphisms in $\mathbb{T}\text{-mod}(\mathbf{Set})$ are all sortwise monic then this expansion can be taken to be fully faithful.*

□

A simple example of a theory satisfying both of the hypotheses of the theorem is the theory $\mathbb{A}$ of commutative rings with unit. Indeed, the finitely presented commutative rings with unit coincide precisely with the finitely generated ones, that is with the finitely presentable models of $\mathbb{A}_m$; also, the monic ring homomorphisms are precisely the injective ones.

Another consequence of Theorem 7.4 is the following criterion for a geometric theory to be of presheaf type.

Theorem 7.7. *Let $\mathbb{T}$ be a geometric theory over a signature Σ . Then $\mathbb{T}$ is of presheaf type if and only if the following conditions are satisfied:*

- (i) *Every finitely presentable model is presented by a geometric formula over Σ ;*
- (ii) *Every property of finite tuples of elements of a (finitely presentable) $\mathbb{T}$ -model which is preserved by $\mathbb{T}$ -model homomorphisms is definable by a geometric formula over Σ ;*
- (iii) *The finitely presentable $\mathbb{T}$ -models are jointly conservative for $\mathbb{T}$.*

Proof The fact that any theory of presheaf type satisfies the given conditions was established in [13]. It thus remains to prove the ‘if’ part of the theorem.

Consider the Σ -structure U in the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ given by the forgetful functors at each sort. Clearly, U is a $\mathbb{T}$ -model.

To deduce our thesis, we shall verify that $\mathbb{T}$ satisfies the conditions of Theorem 7.4 with respect to the model U .

Since every finitely presentable $\mathbb{T}$ -model is presented by a geometric formula over Σ , the first condition of the theorem is satisfied; indeed, any representable functor $\text{Hom}_{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})}(c, -)$ is isomorphic to the interpretation of a formula $\phi(\vec{x})$ in the model U (take as $\phi(\vec{x})$ any formula presenting c). The second condition of the theorem follows immediately from the fact that the finitely presentable $\mathbb{T}$ -models are jointly conservative for $\mathbb{T}$. It remains to show that the third condition of the theorem is satisfied. To this end, we observe that for any geometric formulae $\phi(\vec{x})$ and $\psi(\vec{y})$ over the signature of $\mathbb{T}$, the graph of any arrow $[[\vec{x} \cdot \phi]]_U \rightarrow [[\vec{y} \cdot \psi]]_U$ in the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ is a subobject of the product $[[\vec{x} \cdot \phi]]_U \times [[\vec{y} \cdot \psi]]_U$ in $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ and hence it defines a property of tuples of elements

of finitely presentable models of $\mathbb{T}$ which is preserved by $\mathbb{T}$ -model homomorphisms; therefore, by our assumptions, there exists a formula $\theta(\vec{x}, \vec{y})$ over Σ such that its interpretation in U coincides with such subobject. Since U is conservative and such subobject is the graph of an arrow in the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$, the formula $\theta(\vec{x}, \vec{y})$ is $\mathbb{T}$ -provably functional from $\phi(\vec{x})$ to $\psi(\vec{y})$, as required. $\square$

7.2 Expanding a geometric theory to a theory of presheaf type

In this section we shall discuss the problem of expanding a geometric theory $\mathbb{T}$ to a theory of presheaf type classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$. We shall say that a geometric theory is a *presheaf completion* of a geometric theory $\mathbb{T}$ if it is an expansion of $\mathbb{T}$ such that the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$ is isomorphic to the canonical geometric morphism $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \rightarrow \mathbf{Set}[\mathbb{T}]$.

The results of section 7.1 show that in order to obtain a presheaf completion of a given geometric theory $\mathbb{T}$ we can add a new sort $\ulcorner c \urcorner$ for each finitely presentable $\mathbb{T}$ -model c which is not presented by a geometric formula over the signature of $\mathbb{T}$, and a relation symbol for each subobject of a finite product of objects of $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ which are either of the form $\text{Hom}_{\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})}(c, -)$ (where c is not presented by any geometric formula over the signature of $\mathbb{T}$), or of the form U_A (evaluation functor $\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}) \rightarrow \mathbf{Set}$ at the sort A) where A is a sort over the signature of $\mathbb{T}$ (the sort corresponding to such a representable $\text{Hom}(c, -)$ being $\ulcorner c \urcorner$ and to a functor U_A being A). Indeed, by Theorem 7.4, the theory of the tautological structure over this extended signature will be an expansion of the theory $\mathbb{T}$ classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$. Notice that if $\mathbb{T}$ satisfies the property that every finitely presentable model of $\mathbb{T}$ is presented by a geometric formula over its signature then this expansion of $\mathbb{T}$ is localic over $\mathbb{T}$.

Of course, as it is clear from the results of section 7.1, there are in general many syntactic ways of ‘completing’ a given geometric theory to a theory of presheaf type; the procedure described above represents just a particular choice which is by no means canonical. In fact, what is most interesting in practice is to obtain explicit axiomatizations of presheaf-type completions of a given theory $\mathbb{T}$ directly from the axioms of $\mathbb{T}$ (cf. section 8 below for some examples). Nonetheless, the results established above provide a useful guide in seeking such axiomatizations, as they indicate that in order to complete a geometric theory $\mathbb{T}$ to a theory classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$ one should expand the language of $\mathbb{T}$ in order to make each finitely presentable $\mathbb{T}$ -model presented by a formula in the extended signature and every property of finite tuples of elements of finitely presentable models of $\mathbb{T}$ definable by a geometric formula in the extended signature; we shall see con-

crete applications of these general remarks in section 8. In fact, Theorem 7.7 ensures that making every finitely presentable model $\mathbb{T}$ finitely presented by a formula written in a possibly expanded signature is a necessary condition for the resulting theory to be classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$. Moreover, if one is able to prove that any finitely presentable $\mathbb{T}$ -model is strongly finitely presented as a model of the theory $\mathbb{T}$, when the latter is considered over an extended signature Σ' , condition (iii) of Theorem 5.1 is automatically satisfied, while conditions (i) and (ii) can be made to hold at the cost of adding further axioms to $\mathbb{T}$ over the signature Σ' (cf. Theorem 6.29). On the other hand, by Theorem 7.7, in order to complete $\mathbb{T}$ to a theory of presheaf type classified by the topos $[\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$, one should also expand the signature of $\mathbb{T}$ by adding relation symbols for making any property of finitely presentable $\mathbb{T}$ -models which is preserved by homomorphisms of models definable over the extended signature (if it is not already definable over the original signature).

The following lemma shows that, under appropriate conditions, models which are finitely presented over a given signature remain finitely presented over a larger signature obtained from the former by adding relation symbols characterized by disjunctive sequents of a certain form.

Lemma 7.8. *Let $\mathbb{T}$ be a geometric theory over a signature Σ , Σ' a signature obtained from Σ by only adding relation symbols R and $\mathbb{S}$ a geometric theory over Σ' obtained from $\mathbb{T}$ by adding pairs of axioms of the form $(\top \vdash_{\vec{z}} R(\vec{z}) \vee \bigvee_{i \in I} \phi_i^R(\vec{z}))$ and $(R \wedge \bigvee_{i \in I} \phi_i^R(\vec{z}) \vdash_{\vec{z}} \perp)$, where for each $i \in I$ ϕ_i^R is a geometric formula over Σ such that there exists a geometric formula $\psi_i^R(\vec{z})$ over Σ with the property that the sequents $(\phi_i^R \wedge \psi_i^R \vdash_{\vec{z}} \perp)$ and $(\top \vdash_{\vec{z}} \phi_i^R \vee \psi_i^R)$ are provable in $\mathbb{S}$.*

Let $\mathbb{R}$ be the cartesianization of $\mathbb{S}$ (in the sense of Remark 6.23(d)) and $\phi(\vec{x}) = \phi(x_1, \dots, x_n)$ a $\mathbb{R}$ -cartesian formula over Σ' with the property that there exists a $\mathbb{R}$ -model M_ϕ with n generators $\vec{\xi} = (\xi_1, \dots, \xi_n) \in [[\vec{x} \cdot \phi]]_{M_\phi}$ such that for any $\mathbb{R}$ -model N , the elements of the interpretation $[[\vec{x} \cdot \phi]]_N$ are in natural bijective correspondence with the Σ -structure homomorphisms $f : M_\phi \rightarrow N$ such that $f(\vec{\xi}) \in [[\vec{x} \cdot \phi]]_N$ through the assignment $f \mapsto f(\vec{\xi})$. Then M_ϕ is finitely presented by the formula $\{\vec{x} \cdot \phi\}$ as a $\mathbb{R}$ -model.

Proof We have to prove that, for any $\mathbb{R}$ -model N and any tuple $\vec{a} \in [[\vec{x} \cdot \phi]]_N$, the unique Σ -structure homomorphism $f : M_\phi \rightarrow N$ such that $f(\vec{\xi}) = \vec{a}$ preserves the satisfaction of all the relation symbols R in Σ' , i.e. that for any such symbol R of arity m and any m -tuple $(y_1, \dots, y_m)$ of elements of M_ϕ satisfying R_{M_ϕ} , the m -tuple $(f(y_1), \dots, f(y_m))$ satisfies R_N .

As M_ϕ is by our hypothesis generated by the elements $\xi_1, \dots, \xi_n$, each y_i is equal to the interpretation in M_ϕ of a term $t_i(\xi_1, \dots, \xi_n)$ evaluated in the tuple $\vec{\xi} = (\xi_1, \dots, \xi_n)$. For each $i \in I$, consider the formula $\psi_i'^R(x_1, \dots, x_n)$ obtained by replacing each of the variables $z_1, \dots, z_m$ in the formula ψ_i^R with

the corresponding term t_i . Let us show that the sequent $\phi \vdash_{\vec{x}} \psi_i'^R$ is valid in every $\mathbb{R}$ -model. This amounts to showing that for any $\mathbb{R}$ -model P and any tuple $\vec{b} \in [[\vec{x} \cdot \phi]]_P$, $\vec{b}$ satisfies the formula $\psi_i'^R$. To prove this, we observe that, since M_ϕ is by our hypotheses a $\mathbb{S}$ -model, the m -tuple $(y_1, \dots, y_m)$ satisfies the formula ψ_i^R (for each $i \in I$). Now, $(y_1, \dots, y_m) \in [[\vec{y} \cdot \psi_i^R]]_{M_\phi}$ means that $(\xi_1, \dots, \xi_n) \in [[\vec{x} \cdot \psi_i'^R]]_{M_\phi}$, which implies, since $f_{\vec{b}} : M_\phi \rightarrow P$ is a Σ -structure homomorphism, that $\vec{b} \in [[\psi_i'^R]]_N$, as required. Since $\mathbb{R}$ has enough models as it is cartesian, we can conclude that the sequent $\phi \vdash_{\vec{x}} \psi_i'^R$ is provable in $\mathbb{R}$ and hence in $\mathbb{S}$ (for each $i \in I$). It follows that the cartesian sequent $\phi \vdash_{\vec{x}} R(t_1/z_1, \dots, t_m/z_m)$ is provable in $\mathbb{R}$. By evaluating this sequent in the model N at the tuple $f(\vec{\xi}) \in [[\vec{x} \cdot \phi]]_N$, and using the fact that Σ -structure homomorphisms commute with the interpretation of Σ -terms, we obtain that $(f(y_1), \dots, f(y_m)) = (f(t_1^{M_\phi}(\vec{\xi})), \dots, f(t_m^{M_\phi}(\vec{\xi}))) = (t_1^N(f(\vec{\xi})), \dots, t_m^N(f(\vec{\xi})))$ satisfies R_N , as required. $\square$

Notice that if a model of a geometric theory $\mathbb{T}$ is finitely presentable as a model of its cartesianization then it is strongly finitely presented. In fact, any structure of the form $\text{Hom}_{\mathcal{E}}(E, M)$, where M is a model of $\mathbb{T}$ in a topos $\mathcal{E}$, is a model of the cartesianization of $\mathbb{T}$, as it is obtained by applying a global section functor, which is cartesian, to a model of $\mathbb{T}$. Conversely, if a theory $\mathbb{T}$ is of presheaf type then any finitely presentable model of $\mathbb{T}$ is strongly finitely presented (cf. Corollary 4.10) and hence finitely presented relatively to whatever sub-theory $\mathbb{S}$ of $\mathbb{T}$ (that is, theory $\mathbb{S}$ of which $\mathbb{T}$ is a quotient) whose set-based models admit representations as global sections $\text{Hom}_{\mathcal{E}}(E, M)$ of models M of $\mathbb{T}$ in Grothendieck toposes (pairs of theories satisfying these conditions are investigated for instance in [14] and [16]). This remark can be often profitably applied to the cartesianization of $\mathbb{T}$, even though it is not known in general if it is always the case that every model of it admits a sheaf representation of the above kind.

Summarizing, we have the following theorem.

Theorem 7.9. *Let $\mathbb{T}$ be a theory of presheaf type and $\mathbb{S}$ be a sub-theory of $\mathbb{T}$ such that every set-based model of $\mathbb{S}$ admits a representation as a structure of global sections $\text{Hom}_{\mathcal{E}}(1, M)$ of a model M of $\mathbb{T}$ in a Grothendieck topos $\mathcal{E}$. Then every finitely presentable model of $\mathbb{T}$ is finitely presented as a model of $\mathbb{S}$.*

$\square$

7.3 Faithful expansions

Let $\mathbb{T}$ be a geometric theory over a signature Σ and $\mathbb{T}'$ a geometric expansion of $\mathbb{T}$ over a signature Σ' .

Suppose that $\mathcal{C}$ is a subcategory of the category of finitely presentable $\mathbb{T}'$ -models such that the canonical functor

$$(p_{\mathbb{T}}^{\mathbb{T}'})_{\mathbf{Set}} : \mathbb{T}'\text{-mod}(\mathbf{Set}) \rightarrow \mathbb{T}\text{-mod}(\mathbf{Set})$$

induced by the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'} : \mathbf{Set}[\mathbb{T}'] \rightarrow \mathbf{Set}[\mathbb{T}]$ restricts to a functor

$$j : \mathcal{C} \rightarrow \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}) .$$

Then we have a commutative diagram

$$\begin{array}{ccc} \mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}, J_{\mathbb{T}'}) & \xrightarrow{p_{\mathbb{T}}^{\mathbb{T}'}} & \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}}) \\ s_{\mathcal{C}}^{\mathbb{T}'} \uparrow & & \uparrow t^{\mathbb{T}} \\ [\mathcal{C}, \mathbf{Set}] & \xrightarrow{[j, \mathbf{Set}]} & [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}], \end{array}$$

where the geometric morphisms

$$s_{\mathcal{C}}^{\mathbb{T}'} : [\mathcal{C}, \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}, J_{\mathbb{T}'})$$

and

$$t^{\mathbb{T}} : [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$$

are the canonical ones and $[j, \mathbf{Set}]$ is the geometric morphism canonically induced by the functor j . We shall refer to this diagram as to $(*)$.

Theorem 7.10. *Let $\mathbb{T}$ be a theory of presheaf type over a signature Σ and $\mathbb{T}'$ a geometric expansion of $\mathbb{T}$ over a signature Σ' . Suppose that $\mathbb{T}'$ is classified by the topos $[\mathcal{C}, \mathbf{Set}]$, where $\mathcal{C}$ is a full subcategory of $\text{f.p.}\mathbb{T}'\text{-mod}(\mathbf{Set})$ such that the functor $(p_{\mathbb{T}}^{\mathbb{T}'})_{\mathbf{Set}} : \mathbb{T}'\text{-mod}(\mathbf{Set}) \rightarrow \mathbb{T}\text{-mod}(\mathbf{Set})$ restricts to a faithful functor $j : \mathcal{C} \rightarrow \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})$. Then for any model c of $\mathbb{T}'$ in $\mathcal{C}$ whose associated $\mathbb{T}$ -model $j(c)$ is finitely presented by a geometric formula $\phi(\vec{x})$ over Σ , there exists a geometric formula $\psi(\vec{x})$ over Σ' in the context $\vec{x}$ which presents the $\mathbb{T}'$ -model c and such that the sequent $\psi \vdash_{\vec{x}} \phi$ is provable in $\mathbb{T}'$.*

Before giving the proof of the theorem, we need to recall the following straightforward lemma, of which we give a proof for the reader's convenience.

Lemma 7.11. *Let $R : \mathcal{A} \rightarrow \mathcal{B}$ and $L : \mathcal{B} \rightarrow \mathcal{A}$ be a pair of adjoint functors, where R is the right adjoint and L is the left adjoint. Let $\eta : 1_{\mathcal{B}} \rightarrow R \circ L$ be the unit of the adjunction and b an object of $\mathcal{B}$. Then $\eta(b)$ is monic if and only if for any arrows f, g in $\mathcal{B}$ with codomain b , $Lf = Lg$ implies $f = g$. In particular, η is pointwise monic if and only if the functor L is faithful.*

Proof Let us denote by $\tau_{a,b}$ the bijection between the sets $\text{Hom}_{\mathcal{A}}(Lb, a)$ and $\text{Hom}_{\mathcal{B}}(b, Ra)$ given by the adjunction. By the naturality in b of $\tau_{a,b}$, for any arrow $h : b' \rightarrow b$ in $\mathcal{B}$, the arrow $\eta_b \circ h$ corresponds under $\tau_{Lb, b'}$ to the arrow Lh . Therefore, as $\tau_{Lb, b'}$ is a bijection, $\eta_b \circ f = \eta_b \circ g$ if and only if $Lg = Lf$. From this the thesis follows at once. $\square$

We can now prove the theorem.

Proof The geometric morphism

$$[j, \mathbf{Set}] : [\mathcal{C}, \mathbf{Set}] \rightarrow [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$$

is essential, that is its inverse image $[j, \mathbf{Set}]^*$ admits a left adjoint $[j, \mathbf{Set}]_!$, namely the left Kan extension along the functor j , and the following diagram commutes:

$$\begin{array}{ccc} [\mathcal{C}, \mathbf{Set}] & \xrightarrow{[j, \mathbf{Set}]_!} & [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}] \\ y_{\mathcal{C}} \uparrow & & \uparrow y \\ \mathcal{C}^{\text{op}} & \xrightarrow{j^{\text{op}}} & \text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set})^{\text{op}}, \end{array}$$

where $y_{\mathcal{C}}$ and y are the Yoneda embeddings.

The functor

$$[j, \mathbf{Set}]_! : [\mathcal{C}, \mathbf{Set}] \rightarrow [\text{f.p.}\mathbb{T}\text{-mod}(\mathbf{Set}), \mathbf{Set}]$$

satisfies the property that for any object c of $\mathcal{C}$ and any arrows $\alpha, \beta : P \rightarrow y_{\mathcal{C}}(c)$, where P is an object of $[\mathcal{C}, \mathbf{Set}]$, $[j, \mathbf{Set}]_!(\alpha) = [j, \mathbf{Set}]_!(\beta)$ implies $\alpha = \beta$. Indeed, this is clearly true for P equal to a representable by the commutativity of the above diagram, the full and faithfulness of the Yoneda embeddings $y_{\mathcal{C}}$ and y' and the fact that the functor j is faithful by our hypotheses, and one can always reduce to this case by considering a covering of P in $[\mathcal{C}, \mathbf{Set}]$ by representables.

Now, by our hypotheses the geometric morphisms $s_{\mathcal{C}}^{\mathbb{T}'}$ and $t^{\mathbb{T}}$ defined above are equivalences. Lemma 7.11 thus implies that the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'}$ is essential and the unit of the adjunction between $(p_{\mathbb{T}}^{\mathbb{T}'})^*$ (right adjoint) and $(p_{\mathbb{T}}^{\mathbb{T}'})_!$ (left adjoint) is monic when evaluated at any object of the form $y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{y} \cdot \chi\})$, where $\chi(\vec{y})$ is a formula presenting a $\mathbb{T}'$ -model. Let c be a $\mathbb{T}'$ -model in $\mathcal{C}$. Since $\mathbb{T}'$ is classified by the topos $[\mathcal{C}, \mathbf{Set}]$, c is finitely presented by a formula $\{\vec{x} \cdot \chi\}$ over the signature Σ' of $\mathbb{T}'$. The commutativity of the above square and of diagram (*) thus implies that that $(p_{\mathbb{T}}^{\mathbb{T}'})_!(y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{y} \cdot \chi\})) \cong y_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\})$, where $\phi(\vec{x})$ is a formula over the signature Σ which presents the model $j(c)$ and $y_{\mathcal{C}_{\mathbb{T}}} : \mathcal{C}_{\mathbb{T}} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}}, J_{\mathbb{T}})$, $y_{\mathcal{C}_{\mathbb{T}'}} : \mathcal{C}_{\mathbb{T}'} \rightarrow \mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}, J_{\mathbb{T}'})$ are the Yoneda embeddings. By definition of the geometric morphism $p_{\mathbb{T}}^{\mathbb{T}'}$, we have that $(p_{\mathbb{T}}^{\mathbb{T}'})^*(y_{\mathcal{C}_{\mathbb{T}}}(\{\vec{x} \cdot \phi\})) = y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{x} \cdot \phi\})$ (where the latter ϕ is considered as a formula over the signature Σ'). The unit of the adjunction between $(p_{\mathbb{T}}^{\mathbb{T}'})^*$ and $(p_{\mathbb{T}}^{\mathbb{T}'})_!$ thus yields a monic arrow $y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{y} \cdot \chi\}) \rightarrow p_{\mathbb{T}}^{\mathbb{T}'*}(p_{\mathbb{T}}^{\mathbb{T}'}_!(y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{y} \cdot \chi\}))) \cong y_{\mathcal{C}_{\mathbb{T}'}}(\{\vec{x} \cdot \phi\})$ in the topos $\mathbf{Sh}(\mathcal{C}_{\mathbb{T}'}, J_{\mathbb{T}'})$, in other words a monic arrow $\{\vec{y} \cdot \chi\} \rightarrow \{\vec{x} \cdot \phi\}$ in the geometric syntactic category $\mathcal{C}_{\mathbb{T}'}$. Therefore $\{\vec{y} \cdot \chi\}$ is isomorphic, as an object of $\mathcal{C}_{\mathbb{T}'}$, to an object $\{\vec{x} \cdot \phi'\}$ such that the sequent $(\phi' \vdash_{\vec{x}} \phi)$ is provable in $\mathbb{T}'$; hence $\phi'(\vec{x})$ presents c as a $\mathbb{T}'$ -model, as required. $\square$

8 Examples

In this section we shall discuss in detail various non-trivial examples of theories of presheaf type in light of the theory developed in the paper.

8.1 Theories whose finitely presentable models are finite

As an application of Corollary 6.25, one can recover at once the well-known results that the following theories are of presheaf type:

1. The theory of decidable objects (cf. [23] and p. 907 of [21]);
2. The theory of decidable Boolean algebras (cf. Example D3.4.12 [21]);
3. The theory of linear orders;
4. The theory of total orders with endpoints (cf. section VIII.8 of [26])

Notice that the former two theories are injectivizations of two cartesian theories, namely the empty theory over a signature consisting exactly of one sort and the algebraic theory of Boolean algebras. The latter two theories clearly satisfy the hypothesis of Corollary 6.57; hence their injectivizations are of presheaf type as well.

8.1.1 The theory of abstract circles

The theory $\mathbb{C}$ of abstract circles has been introduced by I. Moerdijk and shown in [28] to have the property that the points of Connes' topos of cyclic sets (cf. [15]) can be identified with the set-based models of $\mathbb{C}$. In the same paper it is also stated that Connes' topos actually classifies $\mathbb{C}$, but the argument given therein seems incomplete.

We shall prove in this section, by using Corollary 6.22, that $\mathbb{C}$ is of presheaf type classified by Connes' topos. Specifically, we shall show that $\mathbb{C}$ satisfies the hypotheses of the corollary with respect to its Horn part (i.e., the Horn theory consisting of the collection of all Horn sequents which are provable in $\mathbb{C}$). We will also prove that the injectivization of $\mathbb{C}$ is of presheaf type as well.

The signature Σ of the theory $\mathbb{C}$ consists of two sorts P and S (variables of type P will be denoted by letters $x, y, \dots$, while variables of type S will be denoted by letters $a, b, \dots$), two unary function symbols $0 : P \rightarrow S$ and $1 : P \rightarrow S$, two unary function symbols $\delta_0 : S \rightarrow P$ and $\delta_1 : S \rightarrow P$, one unary function symbol $*$: $S \rightarrow S$ and a ternary predicate R of type S . The axioms of $\mathbb{C}$ can be formulated as follows:

1. *Non-triviality axioms:*

$$(\top \vdash_{\square} (\exists x)(x = x));$$

$$(\top \vdash_{x,y} (\exists a)(\delta_0(a) = x \wedge \delta_1(a) = y));$$

$$(0(x) = 1(x) \vdash_x \perp)$$

2. ‘Equational’ axioms:

$$(\top \vdash_a a^{**} = a);$$

$$(\top \vdash_a \delta_0(a^*) = \delta_1(a));$$

$$(\top \vdash_x \delta_0(0(x)) = x \wedge \delta_1(0(x)) = x);$$

$$(\top \vdash_x 0(x)^* = 1(x));$$

$$(\delta_0(a) = x \wedge \delta_1(a) = x \vdash_{x,a} a = 0(x) \vee a = 1(x));$$

3. Axioms for concatenation:

$$(R(a, b, c) \wedge R(a, b, c')) \vdash_{a,b,c,c'} c = c';$$

$$(R(a, b, c) \vdash_{a,b,c} \delta_0(c) = \delta_0(a) \wedge \delta_1(c) = \delta_1(b));$$

$$(R(a, b, c) \vdash_{a,b,c} R(c^*, a, b^*));$$

$$R(a, b, d) \wedge R(c, d, e) \vdash_{a,b,c,d,e} (\exists e')(R(b, c, e') \wedge R(a, e', e));$$

$$(R(a, b, 0(x)) \vdash_{a,b,x} a = 0(x));$$

$$(\delta_0(a) = x \vdash_{a,x} R(0(x), a, a);$$

$$(\delta_1(a) = \delta_0(b) \vdash_{a,b} (\exists c)R(a, b, c) \vee (\exists d)R(b^*, a^*, d)) .$$

A model of $\mathbb{C}$ in **Set** is said to be an *abstract circle*. Any set of points P of the circle S^1 defines an abstract circle S_P whose segments $a \in S$ such that $\delta_0(a) = x$ and $\delta_1(a) = y$ are the oriented arcs on S^1 from the point x to the point y . For any natural number $n > 0$, there is exactly one abstract circle, up to isomorphism, whose set of points has n elements; we shall denote it by the symbol C_n .

To prove that $\mathbb{C}$ is of presheaf type, we first notice that the sequent

$$(\delta_0(a) = \delta_0(b) \wedge \delta_1(a) = \delta_1(b) \vdash_{a,b} a = b)$$

is provable in $\mathbb{C}$. We shall refer to it as to the ‘uniqueness axiom’. This sequent can be easily deduced as a consequence of the seventh axiom of group 3 and the fifth axiom of group 2.

It follows that, modulo the non-triviality axioms and the uniqueness axiom, we can replace any expression of the form $(\exists c)\phi(c)$ arising in an axiom of $\mathbb{C}$ with the requirement that the unique segment d such that $\delta_0(d) = \delta_0(c)$ and $\delta_1(d) = \delta_1(c)$ satisfies ϕ . In particular, the fourth axiom of group 3 is provably equivalent, modulo the non-triviality axioms and the uniqueness axiom, to the following sequent:

$$(\delta_0(u) = \delta_0(b) \wedge \delta_1(u) = \delta_1(c) \wedge R(a, b, d) \wedge R(d, c, e) \vdash_{a,b,c,d,e,u} R(b, c, u) \wedge R(a, u, e)) .$$

Similarly, the seventh axiom of group 3 is provably equivalent, modulo the non-triviality axioms and the uniqueness axiom, to the following sequent:

$$(\delta_0(c) = \delta_0(a) \wedge \delta_1(c) = \delta_1(b) \wedge \delta_1(a) = \delta_0(b) \vdash_{a,b,c} R(a, b, c) \vee R(b^*, a^*, c^*)) .$$

From these remarks we see that $\mathbb{C}$ admits a presentation in which all the axioms do not contain quantifications except for the first and second of group 1.

Let us show that $\mathbb{C}$ satisfies the hypotheses of Corollary 6.22 with respect to its Horn part and the category of finite $\mathbb{C}$ -models.

Given a homomorphism $f : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c is a finite model of the Horn part of $\mathbb{C}$, M is a model of $\mathbb{C}$ in a Grothendieck topos $\mathcal{E}$ and E is an object of $\mathcal{E}$, we can ‘localize’ f (in the sense of section 6.4) a finite number of times (once for the first axiom and once for each point of c for the second axiom) to obtain Σ -substructure homomorphisms $f_i : c_i \hookrightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that the structures c_i are finite models of the Horn part of $\mathbb{C}$ satisfying the non-triviality axioms (notice that any substructure of a structure of the form $\text{Hom}_{\mathcal{E}}(E_i, M)$, where M is a model of $\mathbb{C}$ in $\mathcal{E}$, satisfies all the Horn sequents provable in $\mathbb{C}$). We can clearly further localize each of these homomorphisms so to obtain the satisfaction of all the other axioms of $\mathbb{C}$; since this can be done without modifying their domains, the final result will be a family of homomorphisms whose domains are structures which satisfy all the axioms of $\mathbb{C}$.

Next, we notice that for any set-based model M of the Horn part of $\mathbb{C}$, any point $x \in MP$ and any segment $a \in MS$ there exists a finite substructure N of M such that NP contains x and NS contains a (take N equal to the substructure of M given by the sets

$$NP = \{x, \delta_0(a), \delta_1(a)\}$$

and

$$NS = \{a, 0(x), 1(x), 0(\delta_0(a)), 0(\delta_1(a)), 1(\delta_0(a)), 1(\delta_1(a))\} .$$

Moreover, any two finite substructures N_1 and N_2 of M are contained in a common substructure N of M (take N equal to the substructure of M given by $NP = N_1P \cup N_2P$ and $NS = N_1S \cup N_2S$).

This discussion, combined with the argument above (specialized to the case $\mathcal{E} = \mathbf{Set}$), shows that every model of $\mathbb{C}$ in $\mathbf{Set}$ is a directed union of finite models of $\mathbb{C}$. It follows in particular that every finitely presentable $\mathbb{C}$ -model is finite (it being a retract of a finite model). On the other hand, every finite model of $\mathbb{C}$ is finitely presentable as a model of the Horn part of $\mathbb{C}$ (cf. Theorem 6.4 [6]). We can thus conclude that the hypotheses of Corollary 6.22 are satisfied, whence $\mathbb{C}$ is of presheaf type classified by the topos of covariant set-valued functors on the category of finite models of $\mathbb{C}$.

Due to the presence of conjunctions in the premises of some axioms of $\mathbb{C}$, we cannot directly apply Corollary 6.57 to conclude that the injectivization $\mathbb{C}_m$ of $\mathbb{C}$ is of presheaf type. We shall instead apply Corollary 6.56. Since every $\mathbb{C}$ -model in **Set** is a directed union of finite $\mathbb{C}$ -models, the finitely presentable $\mathbb{C}_m$ -models are exactly the finite $\mathbb{C}$ -models. Moreover, by Proposition 6.47, the monic $\mathbb{C}$ -model homomorphisms in **Set** are precisely the homomorphisms which are sortwise injective; indeed, the formulae $\{x^P \cdot \top\}$ and $\{x^S \cdot \top\}$ strongly present respectively the $\mathbb{C}$ -models C_1 and C_2 . To show that the hypotheses of Corollary 6.56 are satisfied, it remains to verify that for any Grothendieck topos $\mathcal{E}$, object E of $\mathcal{E}$ and Σ -structure homomorphism $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where c is a finite $\mathbb{C}$ -model and M is a sortwise decidable $\mathbb{C}$ -model, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a $\mathbb{C}$ -model homomorphism $f_i : c \rightarrow c_i$ of finite $\mathbb{C}$ -models and a sortwise disjunctive Σ -structure homomorphism $x_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $x_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ x$ for all $i \in I$.

In order to apply Proposition 6.53, we make $\mathbb{C}$ into a one-sorted theory by identifying points x with the segments 0_x and rewriting the axioms appropriately. The proposition yields an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a sortwise surjective homomorphism $q_i : c \rightarrow c_i$, where c_i is a finite Σ -structure, and a disjunctive Σ -structure homomorphism (in the sense of Lemma 6.37) $J_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $J_i \circ q_i = \text{Hom}_{\mathcal{E}}(e_i, M) \circ f$ for all $i \in I$. Now, since c is a $\mathbb{C}$ -model and q_i is sortwise surjective, the structure c_i satisfies the non-triviality axioms. We can clearly suppose $E_i \not\cong 0$ without loss of generality, and hence the arrows J_i to be injective. If we consider the image factorizations of the Σ -structure homomorphisms J_i (in the sense of Lemma 6.51), we thus obtain substructures $c'_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ whose underlying sets are the same as those of c_i (since the J_i are injective) and Σ -structure homomorphisms $q'_i : c \rightarrow c'_i$. Since the c'_i have the same underlying set as c_i , they all satisfy the non-triviality axiom, and, at the cost of refining the epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$, we can suppose them to satisfy all the other axioms of $\mathbb{C}$ (cf. the argument given above for showing that $\mathbb{C}$ satisfies the hypotheses of Corollary 6.22 with respect to its Horn part and the category of finite $\mathbb{C}$ -models). So the hypotheses of Corollary 6.56 are satisfied, and we can conclude that $\mathbb{C}_m$ is of presheaf type classified by the topos of covariant set-valued functors on the category of finite models of $\mathbb{C}$ and injective homomorphisms between them.

8.1.2 The geometric theory of finite sets

In this section we shall revisit, from the point of view of the theory developed in the paper, a well-known example of a non-trivial theory of presheaf type, namely the geometric theory $\mathbb{T}$ of finite sets. Recall from [21] (Example D1.1.7(k)) that the signature Σ of $\mathbb{T}$ consists of one sort A and a n -ary

relation symbol R_n for each $n > 0$. The axioms of $\mathbb{T}$ are the following: for each n , one has the axiom

$$\sigma_n := (R_n(x_1, \dots, x_n) \vdash_{\vec{x}, y} \bigvee_{1 \leq i \leq n} y = x_i),$$

expressing the requirement that if an n -tuple of individuals satisfies the relation R_n then it exhausts the members of (the set interpreting) the sort A . (The case $n = 0$ of this axiom is $(R_0 \vdash_{\square} \perp)$, which says that if R_0 holds then the interpretation of the sort A must be empty.) We also have, for each $n > 0$, the axiom

$$(\top \vdash_{\square} \bigvee_{1 \leq i \leq n} (\exists x_1) \cdots (\exists x_n) R_n(x_1, \dots, x_n)) .$$

Finally, to ensure that the interpretations of the R_n are uniquely determined by that of the sort A (i.e. that R_n holds for all n -tuples which exhaust the elements of the interpretation of sort A , and not just for some of them), one adds the axioms

$$(R_n(x_1, \dots, x_n)) \vdash_{x_1, \dots, x_n} R_m(x_{f(1)}, \dots, x_{f(m)})$$

whenever $f : \{1, 2, \dots, m\} \rightarrow \{1, 2, \dots, n\}$ is a surjection, and

$$(R_n(x_1, \dots, x_n) \wedge x_i = x_j \vdash_{x_1, \dots, x_n} R_{n-1}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n))$$

whenever $1 \leq i < j \leq n$.

We can deduce that $\mathbb{T}$ is of presheaf type as an application of Corollary 6.22.

The models of $\mathbb{T}$ in **Set** can be identified with the finite sets, while the $\mathbb{T}$ -model homomorphism are the precisely the surjective functions between them.

We can regard $\mathbb{T}$ as a quotient of its Horn part (i.e. of the theory consisting of all the Horn sequents over the signature of $\mathbb{T}$ which are provable in $\mathbb{T}$). The criterion for finite presentability given by Lemma 6.2 [6] ensures that every finite set, regarded as a model of $\mathbb{T}$, is finitely presentable as a model of the Horn part of $\mathbb{T}$. Indeed, the finiteness of the structure immediately implies that the second condition of the lemma is satisfied (cf. the proof of Theorem 6.4 [6]), while the satisfaction of the first condition follows from the fact that every function from a finite model of cardinality n of the Horn part of $\mathbb{T}$ to a set-based model M of the Horn part of $\mathbb{T}$ which preserves the predicate R_n preserves the predicate R_m for any $m < n$ (by the ‘introduction’ and ‘elimination’ rules expressed by the last two groups of axioms for $\mathbb{T}$).

To apply Corollary 6.22, it thus remains to prove that the second condition in the statement of the corollary is satisfied. Given a homomorphism

$f : a \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where a is a finite model of the Horn part of $\mathbb{T}$ and M is a $\mathbb{T}$ -model in a Grothendieck topos $\mathcal{E}$, if the cardinality of aA is n then for any $m > n$, the sequent σ_m is satisfied in a provided that σ_n is. Indeed, if $m \geq n$ then for any m -tuple $(y_1, \dots, y_m)$ of elements in aA there exists a sub- n -tuple of elements of aA obtained by removing $m - n$ repetitions, which satisfies the relation R_n if the m -tuple $(y_1, \dots, y_m)$ satisfies the relation R_m , since a , as a model of the Horn part of $\mathbb{T}$, satisfies the Horn sequent expressing the ‘elimination’ rule (i.e., the last group of axioms for $\mathbb{T}$). Similarly, by invoking the ‘introduction’ rules, one can prove that if the cardinality of aA is n then for any $k < n$, the sequent σ_k is satisfied in a provided that σ_n is.

Let us show that we can inductively localize the morphism f to eventually arrive at Σ -structure homomorphisms $f_i : a_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ defined on Σ -structures a_i which satisfy all the axioms of $\mathbb{T}$. Notice that if the Σ -structure homomorphisms $h_i : a \rightarrow a_i$ in such localization are quotient maps (in the sense that a tuple in a_i satisfies a relation if and only if it is the image under h_i of a tuple satisfying that relation in a) then the fact that the a_i satisfy the last two groups of axioms of $\mathbb{T}$ will follow automatically from the fact that a does. It will thus be enough to show, by the above considerations, that each a_i satisfies the sequent σ_{n_i} , where n_i is the cardinality of a_iA , and the second axiom of $\mathbb{T}$.

Starting from a Σ -structure homomorphism $f : a \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, where a is a finite model of the Horn part of $\mathbb{T}$, from the fact that M is a model of $\mathbb{T}$ it follows that there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ and for each $i \in I$ a natural number n_i and a n_i -tuple $(\xi_1, \dots, \xi_{n_i})$ of generalized elements $E_i \rightarrow MA$ such that $\langle \xi_1, \dots, \xi_{n_i} \rangle$ factors through the interpretation of R_{n_i} in M . By taking a_i to be the Σ -substructure of $\text{Hom}_{\mathcal{E}}(E_i, M)$ on the finite subset consisting of the elements $\xi_1, \dots, \xi_{n_i}$ plus all the elements in the image of the homomorphism $\text{Hom}_{\mathcal{E}}(e_i, M) \circ f$, we clearly obtain a structure satisfying the second axiom of $\mathbb{T}$. This structure will also satisfy the last two groups of axioms for $\mathbb{T}$ (the validity of Horn sequents is inherited by substructures). Now, in order to obtain from this family of Σ -structure homomorphisms a localization such that the domains of its homomorphisms satisfy all the axioms of $\mathbb{T}$, for each $i \in I$, it suffices to localize each f_i a finite number of times (one for each $(n_i + 1)$ -tuple $\vec{u}$ of elements of a_i , where n_i is the cardinality of a_i), endowing the sets d_i arising in the surjective-injective factorizations of the homomorphisms with the quotient structure induced by the domains a_i via the relevant quotient map q_i (in the sense that the interpretation of each relation symbol R over the signature of $\mathbb{T}$ in such a set d_i is defined to be equal to the image of the interpretation of R in a_i under the quotient map q_i). By the above remarks, these structures will satisfy all the axioms of $\mathbb{T}$.

This completes the proof of the fact that the hypotheses of Corollary 6.22 are satisfied by the theory $\mathbb{T}$ with respect to its Horn part and the category $\mathcal{C}$ of finite $\mathbb{T}$ -models; therefore the theory $\mathbb{T}$ is of presheaf type classified by

the topos $[\mathcal{C}, \mathbf{Set}]$.

8.2 The theory of Diers' fields

As observed in [22], the coherent theory $\mathbb{T}$ of fields is not of presheaf type. In fact, one can easily identify two properties which are preserved by homomorphisms of fields but which are not definable by geometric formulae in the signature of the coherent theory of fields: the property of a field to have characteristic 0, and the property of a tuple $(x_1, \dots, x_n, x_{n+1})$ of elements of a field that the element x_{n+1} is transcendental over the subfield generated by the elements $x_1, \dots, x_n$. This can be easily seen by arguing as follows. Assuming the axiom of choice, the theory $\mathbb{T}$ has enough $\mathbf{Set}$ -based models (it being coherent). Hence if the property of having characteristic 0 were definable by a geometric sentence ϕ over the signature of fields then the sequent $(\top \vdash_{\square} \phi \vee \bigvee_{p \in \mathbb{P}} \phi_p)$, where $\mathbb{P}$ is the set of prime numbers and ϕ_p (for each $p \in \mathbb{P}$)

is the sentence $p \cdot 1 = 0$ expressing the property of having characteristic p , would be provable in $\mathbb{T}$. But, $\mathbb{T}$ being coherent, the infinitary disjunction on the right-hand side of the sequent would then be provably equivalent to a finite subdisjunction, which is absurd as it would imply that the set of all possible characteristics of a field is finite. A similar argument works for the other property, which, like the former, is the complement of a property definable by a strictly infinitary geometric formula.

In order to make such properties definable and possibly obtain a presheaf completion of the theory $\mathbb{T}$, it is thus necessary to enlarge the signature of $\mathbb{T}$ with new relation symbols. In fact, Johnstone introduces in [22] a 0-ary predicate R_0 , expressing the property of a field to have characteristic 0, and for each natural number $n \geq 0$ a $n + 1$ -predicate $R_{n+1}(x_1, \dots, x_n, x_{n+1})$ expressing the property of x_{n+1} of being transcendental over the subfield generated by the elements $x_1, \dots, x_n$. Let Σ' be the resulting signature. Formally, one has to impose the following axioms over Σ' to ensure that these predicates have indeed the required meaning (below we use the abbreviation $Inv(z)$ for the formula $(\exists x)(x \cdot z = 1 \wedge z \cdot x = 1)$):

$$(\top \vdash R_0 \vee \bigvee_{p \in \mathbb{P}} p \cdot 1 = 0);$$

$$(R_0 \wedge p \cdot 1 = 0 \vdash \perp)$$

(for each $p \in \mathbb{P}$);

$$(\top \vdash_{x_1, \dots, x_{n+1}} R_n(x_1, \dots, x_n, x_{n+1}) \vee (\bigvee_{m \in \mathbb{N}, \vec{c}_1^{\vec{n}}, \dots, \vec{c}_m^{\vec{n}}} (\sum_{j=0}^m P_{\vec{c}_j^{\vec{n}}} x_{n+1}^j = 0) \wedge \bigvee_{i \in \{0, 1, \dots, m\}} Inv(P_{\vec{c}_i^{\vec{n}}}))$$

(for each natural number $n \geq 0$), where the former disjunction is taken over all the natural numbers $m \geq 0$ and all the tuples $\vec{c}_i^{\vec{n}}$ (for $i \in \{0, \dots, m\}$) of

integer coefficients (i.e., coefficients of the form $1 + \dots + 1$ an integer number of times) of polynomials in n variables $x_1, \dots, x_n$ of degree $\leq m$ and the expression $P_{\vec{c}_i}$ (for each $i \in \{0, \dots, m\}$) denotes the polynomial term in the variables $x_1, \dots, x_n$ corresponding to the tuple $\vec{c}_i$, and

$$(R_n(x_1, \dots, x_n, x_{n+1}) \wedge (\bigvee_{m \in \mathbb{N}, \vec{c}_1^m, \dots, \vec{c}_m^m} (\sum_{j=0}^m P_{\vec{c}_j^m} x_{n+1}^j = 0) \wedge \bigvee_{i \in \{0, 1, \dots, m\}} \text{Inv}(P_{\vec{c}_i^m}))) \vdash_{x_1, \dots, x_{n+1}} \perp).$$

Let $\mathbb{D}$ be the theory, called in [22] of *Diers fields*, obtained from $\mathbb{T}$ by adding these new predicates and the above-mentioned axioms.

Note that the geometric formula

$$\bigvee_{m \in \mathbb{N}, \vec{c}_1^m, \dots, \vec{c}_m^m} (\sum_{j=0}^m P_{\vec{c}_j^m} x_{n+1}^j = 0) \wedge \bigvee_{i \in \{0, 1, \dots, m\}} \text{Inv}(P_{\vec{c}_i^m})$$

is $\mathbb{D}$ -provably equivalent to a disjunction of geometric formulae, namely, the formulae $(\sum_{j=0}^m P_{\vec{c}_j^m} x_{n+1}^j = 0) \wedge \text{Inv}(P_{\vec{c}_i^m})$ (for each $m, \vec{c}_1^m, \dots, \vec{c}_m^m$ and $i \in \{0, \dots, m\}$), each of which has a $\mathbb{D}$ -provable complement, namely the formula $\text{Inv}(\sum_{j=0}^m P_{\vec{c}_j^m} x_{n+1}^j = 0) \vee P_{\vec{c}_i^m} = 0$.

Notice also that, for any tuples $\vec{c}_i^m$ (for $i \in \{0, \dots, m\}$) of integer coefficients of polynomial expressions $P_{\vec{c}_i^m}$ in the variables $x_1, \dots, x_n$, the cartesian sequent

$$(*) (R_n(x_1, \dots, x_n, x_{n+1}) \wedge \text{Inv}(P_{\vec{c}_i^m}) \vdash_{x_1, \dots, x_{n+1}} \text{Inv}(\sum_{j=0}^m P_{\vec{c}_j^m} x_{n+1}^j = 0))$$

is provable in $\mathbb{D}$.

Following [22], we observe that the theory $\mathbb{D}$ satisfies the property that every finitely presentable model of it, i.e. any finitely generated field, is finitely presented as a model of its cartesianization. This will follow from Lemma 7.8 once we have proved that every finitely generated field F is presented by a finite set of generators, in the weak sense of the lemma, by a formula over the signature of $\mathbb{D}$. To this end, we regard $\mathbb{T}$ as axiomatized in the signature of von Neumann regular rings, which contains a unary function for the operation of pseudoinverse; indeed, over this signature, every finitely generated field (in the sense of field theory) becomes finitely generated (in the sense of model theory).

Notice that, by the above remarks, $\mathbb{D}$ satisfies the first set of hypotheses of Lemma 7.8 with respect to the theory $\mathbb{T}$.

We shall prove that every finitely generated field F is (weakly) presented (as a model of the cartesianization of $\mathbb{D}$) by a geometric formula over the

signature of $\mathbb{D}$ by induction on the number n of generators of F . If $n = 0$ then F is equal to its prime field; so, either F has characteristic p , in which case it is equal to $\mathbb{Z}_p$, or F has characteristic 0 , in which case it is equal to $\mathbb{Q}$. Now, the field $\mathbb{Z}_p$ is clearly (weakly) presented by the formula $p \cdot 1 = 0$, while the field $\mathbb{Q}$ is (weakly) presented by the formula R_0 since the sequent $R_0 \vdash \text{Inv}(n \cdot 1)$ is provable in the cartesianization of $\mathbb{D}$ for each non-zero natural number n . Now, consider a field F generated by $n + 1$ elements $x_1, \dots, x_n, x_{n+1}$. If we denote by F_0 its prime field, we have that $F = F_0(x_1, \dots, x_n)(x_{n+1})$.

Suppose that $F_0(x_1, \dots, x_n)$ is (weakly) presented by a formula in n variables $\phi(\vec{x})$ with the elements $x_1, \dots, x_n$ as generators.

There are two cases: either the element x_{n+1} is transcendental over the field $F_0(x_1, \dots, x_n)$ or not.

In the first case, F is isomorphic to the field of rational functions in one variable with coefficients in $F_0(x_1, \dots, x_n)$, and it is (weakly) presented by the formula $\phi \wedge R_n(x_1, \dots, x_n, x_{n+1})$ with generators $x_1, \dots, x_n, x_{n+1}$; in other words, for any model $(A, \{(R_n)_A \mid n \in \mathbb{N}\})$ of the cartesianization of $\mathbb{D}$, the function which assigns a ring homomorphism $f : F \rightarrow A$ with the property that $f(x_1, \dots, x_n, x_{n+1}) \in (R_{n+1})_A \cap [[\vec{x} \cdot \phi]]_A$ to the element $f(x_1, \dots, x_n, x_{n+1})$ is injective and surjective on $(R_{n+1})_A \cap [[\vec{x} \cdot \phi]]_A$. This can be proved as follows. Since F is generated by the elements $x_1, \dots, x_{n+1}$, the injectivity is clear, so it remains to prove the surjectivity, i.e. that for any $(n+1)$ -tuple $(a_1, \dots, a_n, a_{n+1}) \in (R_{n+1})_A \cap [[\vec{x} \cdot \phi]]_A$ there exists a ring homomorphism $F \rightarrow A$ which sends x_i to a_i for each $i \in \{1, \dots, n+1\}$. By the induction hypothesis, since $(a_1, \dots, a_n) \in [[\vec{x} \cdot \phi]]_A$, there exists a unique ring homomorphism $g : F_0(x_1, \dots, x_n) \rightarrow A$ such that $g(x_i) = a_i$ for each $i \in \{1, \dots, n\}$. By definition of F , there exists a ring homomorphism $f : F \rightarrow A$ which extends g and sends x_{n+1} to a_{n+1} if and only if for every polynomial P with a non-zero coefficient $P_{c_i^n}(x_1, \dots, x_n)$ in $F_0(x_1, \dots, x_n)$, $\sum_{j=0}^m g(P_{c_j^n})a^j$ is invertible in A . But since $P_{c_i^n}(x_1, \dots, x_n)$ is non-zero (equivalently, invertible) in the field $F_0(x_1, \dots, x_n)$, its image $g(P_{c_i^n}(x_1, \dots, x_n)) = P_{c_i^n}(g(x_1), \dots, g(x_n))$ under the homomorphism g is invertible in A ; therefore, since the sequent $(*)$ holds in A , the condition $(f(x_1), \dots, f(x_n), a) \in (R_n)_A$ entails the fact that $\sum_{j=0}^m g(P_{c_j^n})a^j$ is invertible in A , as required.

In the second case, consider the minimal polynomial P for x_{n+1} over $F_0(x_1, \dots, x_n)$; then F is isomorphic to the quotient of $F_0(x_1, \dots, x_n)$ by the ideal generated by the polynomial P . It is immediate to see that F is (weakly) presented by the formula in $n + 1$ variables $\phi \wedge P(x_{n+1}) = 0$ with generators $x_1, \dots, x_n, x_{n+1}$.

These arguments show that the theories $\mathbb{T}$ and $\mathbb{D}$ satisfy the hypotheses

of Lemma 7.8. It follows that all the finitely generated fields are finitely presented models of the cartesianization of $\mathbb{D}$, as required. Condition (iii) of Theorem 5.1 is therefore satisfied (cf. Proposition 5.11(i)). Alternatively, we could have deduced the fact that the theory $\mathbb{D}$ satisfies condition (iii) of Theorem 5.1 from Corollary 5.16 and Theorem 5.21.

In [22], Johnstone shows that $\mathbb{D}$ is a theory of presheaf type by assuming a form of the axiom of choice to ensure that $\mathbb{D}$ has enough set-based models. Theorem 5.1 allows to prove that $\mathbb{D}$ is of presheaf type directly, without assuming any non-constructive principles. Having already proved that condition (iii) of the theorem is satisfied, it remains to see that conditions (i) and (ii) hold as well.

The fact that condition (ii)(a) of Theorem 5.7 holds follows immediately from the above-mentioned discussion in view of Remark 5.8(b). Condition (ii)(c) of Theorem 5.3 is automatically satisfied while condition (ii)(a) of Theorem 5.7 follows from condition (ii)(a) of Theorem 5.7 (cf. Remarks 5.4(b)-(c)). By Remark 5.8(a), to prove that condition (ii)(b) of Theorem 5.7 holds, it suffices to verify that condition (ii)(b) of Theorem 5.3 does. It thus remains to verify that condition (ii)(b) of Theorem 5.3 holds, i.e. that for any finitely generated fields c and d , $\mathbb{D}$ -model M in a Grothendieck topos $\mathcal{E}$ and Σ' -structure homomorphisms $x : c \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$ and $y : d \rightarrow \text{Hom}_{\mathcal{E}}(E, M)$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a finitely generated field b_i , field homomorphisms $u_i : c \rightarrow b_i$, $v_i : d \rightarrow b_i$ and a Σ' -structure homomorphism $z_i : b_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $\text{Hom}_{\mathcal{E}}(e_i, M) \circ x = z_i \circ u_i$ and $\text{Hom}_{\mathcal{E}}(e_i, M) \circ y = z_i \circ v_i$.

We can prove this by induction on the sum n of the minimal number of generators of c and of d .

Before proceeding with the proof, it is convenient to remark the following fact: for any non-zero model $(A, \{(R_n)_A \mid n \in \mathbb{N}\})$ of the cartesianization of $\mathbb{D}$ (for instance, a Σ' -structure of the form $\text{Hom}_{\mathcal{E}}(E, M)$, where M is a model of $\mathbb{D}$ in the topos $\mathcal{E}$ and E is a non-zero object of $\mathcal{E}$) and any field e , considered as a model of $\mathbb{D}$, all the Σ' -structure homomorphisms $e \rightarrow A$ reflect the satisfaction of the relations R_n , i.e. $f(x_1, \dots, x_n) \in (R_n)_A$ implies $(x_1, \dots, x_n) \in (R_n)_e$. This easily follows from the disjunctive axioms of $\mathbb{D}$ defining R_n and the cartesian axiom (*). Note also that such homomorphisms are always injective (since their domain is a field and their codomain is a non-zero ring).

In proving our claim, we can suppose without loss of generality all the objects E arising in Σ' -structure homomorphisms to structures of the form $\text{Hom}_{\mathcal{E}}(E, M)$ to be non-zero (since removing zero arrows from an epimorphic family leaves the family epimorphic).

If $n = 0$ (that is, if both c and d are equal to their prime fields) then c and d have the same characteristic; indeed, equalities of the form $n \cdot 1 = 0$ are preserved and reflected by the homomorphisms x and y (cf. the above remarks). So c and d are isomorphic, whence the claim is trivially satisfied.

Let us now assume that the condition is true for all $k \leq n$ and prove it for $n + 1$. We can represent c as $c'(x)$ in such a way that a set of generators for c can be obtained by adding x to a set of generators for c' ; then by the induction hypothesis there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ and for each $i \in I$ a finitely generated field u_i , field homomorphisms $f_i : c' \rightarrow u_i$ and $g_i : d \rightarrow u_i$ and a Σ' -structure homomorphism $r_i : u_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, M)$ such that $r_i \circ f_i = \text{Hom}_{\mathcal{E}}(E, M) \circ f|_{c'}$ and $r_i \circ g_i = \text{Hom}_{\mathcal{E}}(E, M) \circ g$. Now, consider for each $i \in I$ the element $f(x) \circ e_i \in \text{Hom}_{\mathcal{E}}(E_i, M)$. Suppose that c' has m generators $\chi_1, \dots, \chi_m$; then for each $i \in I$, by the disjunctive axiom of $\mathbb{D}$ involving R_m , there exists an epimorphic family $\{f_{i,j} : F_{i,j} \rightarrow E_i \mid j \in J_i\}$ such that for any $j \in J_i$, either $R_m((r_i \circ f_i)(\chi_1) \circ e_i \circ f_{i,j}, \dots, (r_i \circ f_i)(\chi_m) \circ e_i \circ f_{i,j}, f(x) \circ e_i \circ f_{i,j})$ or $f(x) \circ e_i \circ f_{i,j}$ is the root of a non-zero polynomial P with coefficients belonging to the von Neumann regular subring of $\text{Hom}_{\mathcal{E}}(F_{i,j}, M)$ generated by the elements $(r_i \circ f_i)(\chi_1) \circ e_i \circ f_{i,j}, \dots, (r_i \circ f_i)(\chi_m) \circ e_i \circ f_{i,j}$. In the first case, $f(x) \circ e_i \circ f_{i,j}$ is transcendental over c' via the embedding $\text{Hom}_{\mathcal{E}}(f_{i,j}, M) \circ r_i \circ f_i$ and over u_i via the embedding $\text{Hom}_{\mathcal{E}}(f_{i,j}, M) \circ r_i$ (apply the above remarks to these two embeddings); it follows that the homomorphism f_i extends to a homomorphism from $c = c'(x)$ to u_i .

In the second case, the homomorphism $\text{Hom}_{\mathcal{E}}(e_i \circ f_{i,j}, M) \circ f$ being injective, there exists a non-zero polynomial P with coefficients in c' such that $f(x) \circ e_i \circ f_{i,j}$ is a root of the image of P under $\text{Hom}_{\mathcal{E}}(f_{i,j}, M) \circ r_i \circ f_i = \text{Hom}_{\mathcal{E}}(e_i \circ f_{i,j}, M) \circ f|_{c'}$. It follows that x is a root of P in c . We can thus clearly suppose P to be irreducible without loss of generality, and represent $c = c'(x)$ as $c = c'[z]/P(z)$, via an isomorphism sending z to x ; denoting P' the image of P under the homomorphism f_i , we thus obtain that the arrow f_i yields an arrow $c = c'[z]/P(z) \rightarrow u_i[w]/P'(w)$ and the homomorphism $\text{Hom}_{\mathcal{E}}(f_{i,j}, M) \circ r_i$ factors through the quotient map $u_i \rightarrow u_i[w]/P'(w)$ yielding a ring homomorphism which is in fact a Σ' -structure homomorphism (by Lemma 7.8, cf. the argument given above for proving that every finitely generated field is finitely presented as a model of the cartesianization of $\mathbb{D}$).

From these remarks it is now straightforward to obtain a set of data satisfying the requirements of our condition.

We could have alternatively proved that $\mathbb{D}$ satisfies condition (ii)(b) of Theorem 5.3 either by using Remark 5.4(d) or by applying Theorem 5.5.

8.3 The theory of algebraic extensions of a given field

Let F be a field. We define the theory $\mathbb{T}_F$ of *algebraic extensions* of F as the expansion of the coherent theory of fields obtained by adding one constant symbol $\bar{a}$ for each element $a \in F$ and the following axioms:

$$(\top \vdash \overline{1_F} = 1);$$

$$(\top \vdash \overline{0_F} = 0);$$

$$(\top \vdash \overline{a} + \overline{b} = \overline{a +_F b}),$$

for any elements $a, b \in F$ (the symbol $+_F$ denotes the addition operation in F);

$$(\top \vdash \overline{a} \cdot \overline{b} = \overline{a \cdot_F b}),$$

for any elements $a, b \in F$ (the symbol $\cdot_F$ denotes the multiplication operation in F), plus the algebraicity axiom

$$(\top \vdash_x \bigvee_{n \in \mathbb{N}, a_0, \dots, a_{n-1}, a_n \in F} \overline{a_n} x^n + \overline{a_{n-1}} x^{n-1} + \dots + \overline{a_0} = 0).$$

We shall prove that $\mathbb{T}_F$ is of presheaf type. Clearly, the finitely presentable models of $\mathbb{T}_F$ are exactly the finitely generated algebraic extensions of F , that is the finite extensions of F . One can prove, by adapting the argument used in the proof of the result that every finitely generated field is finitely presented as a model of the cartesianization of the theory of Diers' fields established in section 8.2, that every finite extension of F is finitely presented as a model of the cartesianization of $\mathbb{T}_F$. Specifically, every finite extension $F(x_1, \dots, x_n)$ of F is presented by the conjunction of the formulae of the form $P_i(x_1, \dots, x_{i+1})(x_i) = 0$ (for $i = 0, \dots, n-1$), where P is the minimal polynomial for the element x_{i+1} over the field $F(x_1, \dots, x_i)$.

Condition (iii) of Theorem 5.1 is thus satisfied by the theory $\mathbb{T}_F$ with respect to the category of finite extensions of F (cf. Proposition 5.11(i)).

In verifying that conditions (i) and (ii) of Theorem 5.1 are satisfied, one is reduced as in section 8.2 to check that condition (ii)(b) of Theorem 5.3 holds; this can be done again by adapting the argument given in section 8.2 to this case. We can thus conclude that $\mathbb{T}_F$ is of presheaf type classified by the topos of covariant set-valued functors on the category of finite extensions of F .

Next, let us consider the theory $\mathbb{S}_F$ of $\mathbb{T}_F$ of *separable extensions* of F , that is the quotient of $\mathbb{T}_F$ obtained by adding the following sequent:

$$(\top \vdash_x \bigvee_{n \in \mathbb{N}, (a_0, \dots, a_{n-1}, a_n) \in \mathcal{S}_F^n} \overline{a_n} x^n + \overline{a_{n-1}} x^{n-1} + \dots + \overline{a_0} = 0),$$

where $\mathcal{S}_F^n$ is the set of n -tuples of elements $a_1, \dots, a_n$ of F such that the polynomial $a_n Z^n + a_{n-1} Z^{n-1} + \dots + a_0 \in F[Z]$ is irreducible and separable.

Clearly, the finitely presentable $\mathbb{S}_F$ -models are precisely the finite separable extensions of F . In particular, every finitely presentable $\mathbb{S}_F$ -model is finitely presentable as a $\mathbb{T}_F$ -model. In fact, by Artin's primitive element theorem, every finite separable extension of F is presented by a formula of the form $\{x \cdot \overline{a_n} x^n + \overline{a_{n-1}} x^{n-1} + \dots + \overline{a_0} = 0\}$.

The hypotheses of Corollary 6.22 are trivially satisfied; indeed, for any $\mathbb{T}_F$ -model homomorphism $M \rightarrow N$ (in an arbitrary Grothendieck topos), if

N is separable (i.e., a model of $\mathbb{S}_F$) then M is *a fortiori* separable as well. We can thus conclude that also the theory $\mathbb{S}_F$ is of presheaf type, classified by the category of covariant set-valued functors on the category of finite separable extensions of F .

Remark 8.1. The theory of fields of a fixed finite characteristic p which are algebraic over their prime field, which was proved in [11] to be of presheaf type classified by the category of covariant set-valued functors on the category of finite fields of characteristic p , is (trivially) Morita-equivalent to the theory $\mathbb{T}_{\mathbb{Z}_p}$ introduced above.

8.4 Groups with decidable equality

In this section we shall study the injectivization $\mathbb{G}$ of the (algebraic) theory of groups.

Clearly, the finitely presentable $\mathbb{G}$ -models are precisely the finitely generated groups.

Even if $\mathbb{G}$ satisfies condition (iii) of Theorem 5.1 with respect to the category of finitely generated groups and injective homomorphisms between them (by Theorem 5.21 and Corollary 5.16), $\mathbb{G}$ is not of presheaf type. To see this, consider the property of an element x of a group G to be non-nilpotent. This property is clearly preserved by injective homomorphisms of groups, so if $\mathbb{G}$ were of presheaf it would be definable by a geometric formula $\phi(x)$ over the signature of $\mathbb{G}$. Then the sequent

$$(\top \vdash_x \bigvee_{n \in \mathbb{N}} \phi \vee (x^n = 1)),$$

would be provable in $\mathbb{G}$ (since $\mathbb{G}$ has enough set-based models and this sequent is valid in every set-based $\mathbb{G}$ -model by definition of ϕ) and hence, as $\mathbb{G}$ is coherent, the disjunction on the right hand side would be $\mathbb{G}$ -provably equivalent to a finite sub-disjunction; but this is absurd since it implies that there exists a natural number n such that every nilpotent element x of a group satisfies $x^n = 1$.

To obtain a presheaf completion of the theory $\mathbb{G}$, we add to the signature of $\mathbb{G}$ a relation symbol R_N^n for each natural number n and any normal subgroup of the free group F_n on n generators, and the following axioms (where the symbol $\neq$ denotes the predicate of $\mathbb{G}$ which is $\mathbb{G}$ -provably complemented to the equality relation):

$$(\top \vdash_{\vec{x}} R_N^n(\vec{x}) \vee (\bigvee_{w(\vec{x}), w'(\vec{x}) \in F_n \mid ww'^{-1} \in N} w \neq w') \vee (\bigvee_{w(\vec{x}), w'(\vec{x}) \in F_n \mid ww'^{-1} \notin N} w = w'))$$

and

$$(R_N^n(\vec{x}) \wedge (\bigvee_{w(\vec{x}), w'(\vec{x}) \in F_n \mid ww'^{-1} \in N} w \neq w' \vee \bigvee_{w(\vec{x}), w'(\vec{x}) \in F_n \mid ww'^{-1} \notin N} w = w')) \vdash_{\vec{x}} \perp$$

(for any $n \in \mathbb{N}$ and any normal subgroup N of F_n), and

$$(\top \vdash_{\vec{x}} \bigvee_{N \in \mathcal{N}_n} R_N^n(\vec{x})),$$

where $\mathcal{N}_n$ is the set of normal subgroups of F_n (for any $n \in \mathbb{N}$).

Let $\mathbb{G}_p$ be the resulting theory; we shall prove that it is of presheaf type.

Notice that the theories $\mathbb{G}$ and $\mathbb{G}_p$ satisfy the first set of hypotheses of Lemma 7.8.

Let us first show that every finitely generated group is finitely presented as a model of the cartesianization of $\mathbb{G}_p$. By Lemma 7.8, to prove that F_n/N is presented by the formula $R_N^n(\vec{x})$ as a model of the cartesianization of $\mathbb{G}_p$, it suffices to verify that for any set-based model $(G', \{(R_N^n)_G \mid n \in \mathbb{N}, N \in \mathcal{N}_n\})$ of the cartesianization of $\mathbb{G}_p$, denoting by $\vec{\xi} = (\xi_1, \dots, \xi_n)$ the generators of the group F_n , the group homomorphisms $f : F_n/N \rightarrow G'$ such that $f(\vec{\xi}) \in [[\vec{x} \cdot R_N^n]]_{G'}$ correspond exactly to the n -tuples $\vec{y}$ of elements of G' which belong to the interpretation in G' of the formula R_N^n (via the assignment $f \rightarrow f(\vec{\xi})$). Given a n -tuple $\vec{y}$ of elements of G' which belong to the interpretation of R_N^n in G' , we can define a function $f : F_n/N \rightarrow G'$ by setting $f([w]) = w_{G'}(\vec{y})$. This is a well-defined injective group homomorphism since the following sequents are provable in the cartesianization of $\mathbb{G}_p$ and hence are valid in G' :

$$(R_N^n \vdash_{\vec{x}} w = w')$$

for any $w, w' \in F_n$ such that $ww'^{-1} \in N$, and

$$(R_N^n \vdash_{\vec{x}} w \neq w')$$

for any $w, w' \in F_n$ such that $ww'^{-1} \notin N$. Since every finitely generated group is, up to isomorphism, of the form F_n/N for some natural number n and a normal subgroup N of F_n , we can conclude that the theory $\mathbb{G}_p$ satisfies condition (iii) of Theorem 5.1 (cf. Proposition 5.11(i)). An alternative way to prove this would have been to invoke Corollary 5.16 and Theorem 5.21.

As in the case of Diers fields treated in section 8.2, in order to verify that the theory $\mathbb{G}_p$ satisfies conditions (i) and (ii) of Theorem 5.1, one is reduced to show that condition (ii)(b) of Theorem 5.3 is satisfied; but this follows from Remark 5.4(d). The theory $\mathbb{G}_p$ is thus of presheaf type classified by the topos $[\text{f.g.}\mathbf{Grp}, \mathbf{Set}]$, where f.g. $\mathbf{Grp}$ is the category of finitely generated groups and injective homomorphisms between them.

The category f.g. $\mathbf{Grp}$ is cocartesian; indeed, it has an initial object (namely, the trivial group) and pushouts (given by the free product with amalgamation construction, cf. [30]). It follows the topos $[\text{f.g.}\mathbf{Grp}, \mathbf{Set}]$ is coherent. The theory $\mathbb{G}_p$, in spite of being infinitary, is thus classified by a coherent topos. This fact has various implications for $\mathbb{G}_p$. For instance,

the coherence of the classifying topos for $\mathbb{G}_p$ implies that every formula over the signature of $\mathbb{G}_p$ presenting a finitely generated group (regarded as a $\mathbb{G}_p$ -model) is not only $\mathbb{G}_p$ -irreducible, but also a coherent object of the classifying topos; in particular, for any $\mathbb{G}_p$ -compact formula $\{\vec{x} . \phi\}$ over the signature of $\mathbb{G}_p$, the formula $\{x, y . \phi(x) \wedge \phi(y)\}$ is also $\mathbb{G}_p$ -compact (recall that a formula $\{\vec{z} . \chi\}$ over the signature Σ of a geometric theory $\mathbb{T}$ is $\mathbb{T}$ -compact if whenever $\{\vec{z} . \chi\}$ $\mathbb{T}$ -provably entails a disjunction of geometric formulas over Σ , $\{\vec{z} . \chi\}$ $\mathbb{T}$ -provably entails a finite sub-disjunction of it). Semantically, a formula $\{\vec{x} . \phi\}$ is $\mathbb{G}_p$ -compact if whenever $\{S_i \mid i \in I\}$ is a family of assignments $G \rightarrow S_i^G \subseteq G^n$ sending each finitely generated group G to a subset $S_i^G \subseteq [[\vec{x} . \phi]]_G$ in such a way that every injective homomorphism $f : G \rightarrow G'$ of groups sends tuples in S_i^G to tuples in $S_i^{G'}$, if $[[\vec{x} . \phi]]_G = \bigcup_{i \in I} S_i^G$ for all G then there exists a finite subset $J \subseteq I$ such that $[[\vec{x} . \phi]]_G = \bigcup_{i \in J} S_i^G$. The fact that the classifying topos of $\mathbb{G}_p$ is coherent also implies, by Deligne's theorem (assuming the axiom of choice), the existence of set-based models for any non-contradictory quotient of $\mathbb{G}_p$ whose associated Grothendieck topology on f.g. $\mathbf{Grp}^{\text{op}}$ is of finite type.

8.5 Locally finite groups

Let $\mathbb{A}$ be the algebraic theory of groups. Since every finite group is finitely presented as a $\mathbb{A}$ -model (cf. Theorem 6.4 [6]) and $\mathbb{A}$ is of presheaf type, Theorem 6.29 ensures that there exists a quotient $\mathbb{U}$ of $\mathbb{A}$ classified by the topos $[\mathcal{C}, \mathbf{Set}]$, where $\mathcal{C}$ is the category of finite groups and homomorphisms between them, which can be characterized as the set of all geometric sequents over the signature of $\mathbb{A}$ that are valid in every finite group. On the other hand, since $\mathbb{U}$ is classified by the topos $[\mathcal{C}, \mathbf{Set}]$, the set-based models of $\mathbb{U}$ are exactly the groups which can be expressed as filtered colimits of finite groups, such groups are exactly the groups which validate all the geometric sequents over the signature of $\mathbb{A}$ which hold in every finite group. As a by-product, we obtain the following characterization of locally finite groups (equivalently, of the groups which can be expressed as filtered colimits of finite groups).

Proposition 8.2. *The locally finite groups are exactly the groups which satisfy all the geometric sequents over the signature of the theory of groups which hold for all finite groups.*

Proof In view of the above remarks, it remains to verify that a group is locally finite (in the sense that all its finitely generated subgroups are finite) if and only if it is a filtered colimit of finite groups. This can be proved as follows. If a group is locally finite then it is the filtered union of all its finitely generated (and hence finite) subgroups. Conversely, suppose that G is a filtered colimit of finite groups. Then G is the directed union of the

images in G of these finite subgroups, which are again finite. It follows that every finitely generated subgroup H of G is contained in one of them (notice that, since the union is filtered, there exists one of them which contains all the generators of H) and hence it is *a fortiori* finite, as required. $\square$

The injectivization of $\mathbb{A}$ is also of presheaf type (by Corollary 6.57) and can be characterized as the quotient of $\mathbb{G}$ consisting of all the geometric sequents over the signature of the injectivization $\mathbb{G}$ of the theory of groups which hold in all finite groups.

8.6 Vector spaces

Let $\mathbb{V}_K$ be the expansion of the algebraic theory of vector spaces over a field K obtained by adding a n -ary predicate R_n for each natural number n expressing the property of a n -tuple of elements to be linearly independent, i.e. the following sequents:

$$(\top \vdash_{\vec{x}} R_n(\vec{x}) \vee \bigvee_{\substack{(k_1, \dots, k_n) \in K^n \\ k_i \neq 0 \text{ for some } i}} k_1 x_1 + \dots + k_n x_n = 0),$$

and

$$(R_n(\vec{x}) \wedge (\bigvee_{\substack{(k_1, \dots, k_n) \in K^n \\ k_i \neq 0 \text{ for some } i}} k_1 x_1 + \dots + k_n x_n = 0) \vdash_{\vec{x}} \perp).$$

The category of models of the theory $\mathbb{V}_K$ in **Set** has as objects the vector spaces over K and as arrows the injective homomorphisms between them. The finitely presentable $\mathbb{V}_K$ -models are precisely the finite-dimensional vector spaces over K .

By using techniques analogous to those employed in section 8.2, one can prove that $\mathbb{V}_K$ is of presheaf type.

Also, by using Corollary 6.22, one can easily prove that for any fixed natural number n , both the expansion of the theory of vector spaces over a field K and of the theory $\mathbb{V}_K$ obtained by adding the sequent

$$\top \vdash_{(x_1, \dots, x_{n+1})} \bigvee_{\substack{(k_1, \dots, k_{n+1}) \in K^{n+1} \\ k_i \neq 0 \text{ for some } i}} k_1 x_1 + \dots + k_{n+1} x_{n+1} = 0)$$

are of presheaf type.

The models in **Set** of the latter theory are precisely the vector spaces over K of dimension $\leq n$.

8.7 The theory of abelian l -groups with strong unit

Recall that an abelian l -group with strong unit is a lattice-ordered group $(G, 0, \leq)$ with a distinguished element u , called the unit of the group, such

that for any element $x \in G$ such that $x \geq 0$ there exists a natural number n such that $x \leq nu$, where $nu = u + \dots + u$ n times. We refer the reader to chapter 2 of [31] as an introduction to the theory of lattice-ordered groups.

We can axiomatize the theory $\mathbb{L}_u$ of abelian l -groups with strong unit over a signature Σ consisting of four binary function symbols $+$, $-$, $\inf$, $\sup$, two constants 0 and u and a binary relation symbol $\leq$, by using Horn sequents to formalize the notion of abelian l -group and the following geometric sequent to express the property of strong unit:

$$(x \geq 0 \vdash_x \bigvee_{n \in \mathbb{N}} x \leq nu) .$$

The following lemma will be useful in showing that the theory $\mathbb{L}_u$ is of presheaf type.

Lemma 8.3. *Let G be an abelian group with a distinguished element u and generators $x_1, \dots, x_n$. If for every $i \in \{1, \dots, n\}$ there exists a natural number k_i such that $|x_i| \leq k_i u$ then u is a strong unit for G .*

Proof Recall that the absolute value $|x|$ of an element x of an abelian l -group with unit $(G, 0, +, -, \leq, \inf, \sup)$ is the element $\sup(x, -x)$. For any $x \in G$, $|x| \geq 0$ $|x| = |-x|$, and for any $x, y \in G$, the triangular inequality $|x + y| \leq |x| + |y|$ holds.

Since G is generated by elements $x_1, \dots, x_n$, every element x of G can be expressed as the interpretation $t(x_1, \dots, x_n)$ of a term t over the signature Σ . We shall prove that there exists a natural number n such that $|x| \leq nu$ by induction on the structure of t . This will clearly imply our thesis, since if $x \geq 0$ then $|x| = x$. If t is a variable then the claim is clearly true by our hypothesis. If $x = x' + x''$ with $|x'| \leq n'u$ and $|x''| \leq n''u$ then by the triangular inequality we have $|x| \leq n' + n''$, and similarly for the subtraction. The $\inf$ and $\sup$ cases are similarly straightforward. $\square$

Let us now verify that the theory $\mathbb{L}_u$ satisfies the hypotheses of Corollary 6.22 with respect to its Horn part $\mathbb{H}$.

We have to prove that for any finitely presentable $\mathbb{H}$ -model c , any model G of $\mathbb{L}_u$ in a Grothendieck topos $\mathcal{E}$, any object E of $\mathcal{E}$ and any Σ -structure homomorphism $f : c \rightarrow \text{Hom}_{\mathcal{E}}(E, G)$, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $i \in I$ a finitely presentable model c_i of $\mathbb{L}_u$ and Σ -structure homomorphisms $f_i : c \rightarrow c_i$ and $u_i : c_i \rightarrow \text{Hom}_{\mathcal{E}}(E_i, G)$ such that $\text{Hom}_{\mathcal{E}}(e_i, G) \circ f = u_i \circ f_i$ for all $i \in I$. Let us suppose that c is presented as a $\mathbb{H}$ -model by a cartesian formula $\phi(\vec{y})$ with generators $x_1, \dots, x_n$. Since G is a l -group with strong unit, there exists an epimorphic family $\{e_i : E_i \rightarrow E \mid i \in I\}$ in $\mathcal{E}$ and for each $k \in \{1, \dots, n\}$ and $i \in I$ a natural number $m_{k,i}$ such that $f(|x_k|) \circ e_i \leq m_{k,i} u_{\text{Hom}_{\mathcal{E}}(E_i, G)}$ (where $u_{\text{Hom}_{\mathcal{E}}(E_i, G)}$ denotes the unit of the l -group $\text{Hom}_{\mathcal{E}}(E_i, G)$). For each $i \in I$, let c_i the $\mathbb{H}$ -model presented

by the cartesian formula $\phi(x_1, \dots, x_n) \wedge |x_1| \leq m_{1,i} \wedge \dots \wedge |x_n| \leq m_{n,i}$. For each $i \in I$, we have a natural quotient homomorphism $f_i : c \rightarrow c_i$ through which $\text{Hom}_{\mathcal{E}}(e_i, G) \circ f$ factors; the resulting factorization u_i satisfies the required property $\text{Hom}_{\mathcal{E}}(e_i, G) \circ f = u_i \circ f_i$. Since $\mathbb{H}$ is a Horn theory, each c_i is generated by the n -tuple $f_i(x_1), \dots, f_i(x_n)$ which presents it as a $\mathbb{H}$ -model (cf. Remark 4.2(a)). Therefore the c_i are ℓ -groups with strong unit by Lemma 8.3.

This argument also shows that the finitely presentable $\mathbb{L}_u$ -models are exactly the finitely presented $\mathbb{H}$ -models whose unit is strong (cf. Theorem 6.26).

Therefore all the hypotheses of Corollary 6.22 are satisfied and we can conclude that the theory $\mathbb{L}_u$ is of presheaf type. In fact, $\mathbb{L}_u$ is Morita-equivalent to the (algebraic) theory of MV-algebras (cf. [12]).

References

- [1] M. Artin, A. Grothendieck and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas*, Séminaire de Géométrie Algébrique du Bois-Marie, année 1963-64; second edition published as Lecture Notes in Math., vols. 269, 270 and 305(Springer-Verlag, 1972).
- [2] T. Beke, Theories of presheaf type, *J. Symbolic Logic* 69 (3), 923-934 (2004).
- [3] A. Blass and A. Šcedrov, Classifying topoi and finite forcing, *Journal of Pure and Applied Algebra* 28 (2), 111-140 (1983).
- [4] F. Borceux, *Handbook of categorical algebra* vol. 1, (Cambridge University Press, 1994).
- [5] O. Caramello, Yoneda representations of flat functors and classifying toposes, *Theory and Applications of Categories* 26 (21), 538-553 (2012).
- [6] O. Caramello, Topological Galois Theory, *arXiv:math.CT/1301.0300*.
- [7] O. Caramello, Fraïssé's construction from a topos-theoretic perspective, *arXiv:math.CT/0805.2778*, to appear in *Logica Universalis*.
- [8] O. Caramello, Syntactic characterizations of properties of classifying toposes, *Theory and Applications of Categories* 26 (6), 176-193 (2012).
- [9] O. Caramello, Lattices of theories, *arXiv:math.CT/0905.0299*.
- [10] O. Caramello, The unification of Mathematics via Topos Theory, *arXiv:math.CT/1006.3930v1*.
- [11] O. Caramello and P. T. Johnstone, De Morgan's law and the theory of fields, *Advances in Mathematics* 222 (6), 2145-2152 (2009).
- [12] O. Caramello and A. C. Russo, The Morita-equivalence between MV-algebras and abelian ℓ -groups with strong unit, *arXiv:math.CT/arXiv:1312.1272*.
- [13] O. Caramello, Universal models and definability, *Math. Proc. Cam. Phil. Soc.* 152 (2), 279-302 (2012).

- [14] J. Cole, The bicategory of toposes, and spectra, unpublished manuscript.
- [15] A. Connes, Cohomologie cyclique et foncteur Ext^n , *C.R. Acad. Sci. Paris, S  r. I Math* 296, 953-958 (1983).
- [16] M. Coste, Localisation, spectra and sheaf representation, *Applications of sheaves*, Lecture Notes in Math. vol. 753 (Springer-Verlag, 1979), 212-238.
- [17] D. Coumans, Generalising canonical extension to the categorical setting, *Annals of Pure and Applied Logic*, 163 (12), 1940-1961 (2012).
- [18] J. Giraud, Analysis situs, *S  minaire Bourbaki*, expos   256 (1963).
- [19] M. Hakim, *Topos annel  s et sch  mas relatifs* (Springer-Verlag, 1972).
- [20] W. Hodges, *Model theory*, Encyclopedia of Mathematics and its Applications vol. 42 (Cambridge University Press, 1993).
- [21] P. T. Johnstone, *Sketches of an Elephant: a topos theory compendium*, Vols. 1 and 2, vols. 43 and 44 of *Oxford Logic Guides* (Oxford University Press, 2002).
- [22] P. T. Johnstone, A syntactic approach to Diers' localizable categories, in *Applications of sheaves*, Lecture Notes in Math. vol. 753 (Springer-Verlag, 1979), 466-478.
- [23] P. T. Johnstone and G. Wraith, Algebraic theories in toposes, in *Indexed categories and their applications*, Lecture Notes in Mathematics (Springer-Verlag) vol. 661 141-242 (1978)
- [24] A. Joyal and G. Wraith, Eilenberg-MacLane toposes and cohomology, *Mathematical applications of category theory*, Contemporary Mathematics, vol. 30 (American Mathematical Society, 1984), 117-131.
- [25] W. Lawvere, Functorial Semantics of Algebraic Theories, 1963, republished in *Reprints in Theory and Applications of Categories* 5, 1-121 (2004).
- [26] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic: a first introduction to topos theory* (Springer-Verlag, 1992).
- [27] M. Makkai and G. Reyes, *First-order categorical logic*, Lecture Notes in Mathematics (Springer-Verlag) vol. 611 (1977).
- [28] I. Moerdijk, Cyclic sets as a classifying topos (Utrecht, 1996), available at <http://ncatlab.org/nlab/files/MoerdijkCyclic.pdf>.
- [29] R. Par   and D. Schumacher, Abstract families and the Adjoint Functor Theorems, in *Indexed categories and their applications*, Lecture Notes in Math. vol. 661 (Springer-Verlag, 1978), 1-125.
- [30] O. Schreier, Die Untergruppen der freien Gruppen, *Abh. Math. Sem. Univ. Hamburg* vol 5, 161-183 (1927).
- [31] S. A. Steinberg, *Lattice-ordered Rings and Modules*, Springer (2010).
- [32] S. Vickers, Strongly algebraic = SFP (topically), *Mathematical Structures in Computer Science* 11 (6), 717-742 (2001).