

# Fractal Zeta Functions and Complex Dimensions: A General Higher-Dimensional Theory

Michel L. LAPIDUS, Goran RADUNOVIC and Darko ZUBRINIC



Institut des Hautes Études Scientifiques  
35, route de Chartres  
91440 – Bures-sur-Yvette (France)

Juillet 2015

IHES/M/15/16

# Fractal zeta functions and complex dimensions: A general higher-dimensional theory

Michel L. Lapidus, Goran Radunović and Darko Žubrinčić

**Abstract.** In 2009, the first author introduced a class of zeta functions, called ‘distance zeta functions’, which has enabled us to extend the existing theory of zeta functions of fractal strings and sprays (initiated by the first author and his collaborators in the early 1990s) to arbitrary bounded (fractal) sets in Euclidean spaces of any dimensions. A closely related tool is the class of ‘tube zeta functions’, defined using the tube function of a fractal set. These zeta functions exhibit deep connections with Minkowski contents and upper box (or Minkowski) dimensions, as well as, more generally, with the complex dimensions of fractal sets. In particular, the abscissa of (Lebesgue, i.e., absolute) convergence of the distance zeta function coincides with the upper box dimension of a set. We also introduce a class of transcendently quasiperiodic sets, and describe their construction based on a sequence of carefully chosen generalized Cantor sets with two auxiliary parameters. As a result, we obtain a family of “maximally hyperfractal” compact sets and relative fractal drums (i.e., such that the associated fractal zeta functions have a singularity at every point of the critical line of convergence). Finally, we discuss the general fractal tube formulas and the Minkowski measurability criterion obtained by the authors in the context of relative fractal drums (and, in particular, of bounded subsets of  $\mathbb{R}^N$ ).

**Keywords.** Fractal set, fractal zeta functions, distance zeta function, tube zeta function, geometric zeta function of a fractal string, Minkowski content, Minkowski measurability, upper box (or Minkowski) dimension, complex dimensions of a fractal set, holomorphic and meromorphic functions, abscissa of convergence, quasiperiodic function, quasiperiodic set, relative fractal drum, fractal tube formulas.

---

The research of Michel L. Lapidus was partially supported by the National Science Foundation under grants DMS-0707524 and DMS-1107750, as well as by the Institut des Hautes Études Scientifiques (IHÉS) where the first author was a visiting professor in the Spring of 2012 while part of this research was completed. This paper is based on a plenary lecture given by the first author (MLL) at the International Conference on “Fractal Geometry and Stochastics” held in Tabarz, Germany, in March 2014. Goran Radunović and Darko Žubrinčić express their gratitude to the Ministry of Science of the Republic of Croatia for its support.

## 1. Introduction

This article provides a short survey of some of the recent advances in the theory of fractal zeta functions and the associated higher-dimensional theory of complex dimensions, valid for arbitrary bounded subsets of Euclidean spaces and developed in the forthcoming research monograph [LapRaŽu1], entitled *Fractal Zeta Functions and Fractal Drums: Higher-Dimensional Theory of Fractal Dimensions*. (See also the research articles [LapRaŽu2–6] and the survey article [LapRaŽu7].)

The theory of zeta functions of fractal strings, initiated by the first author in the early 1990s and described in an extensive research monograph [Lap-vFr3], joint with M. van Frankenhuijsen (see also the references therein), was given an unexpected impetus in 2009, when a new class of zeta functions, called ‘distance zeta functions’, was discovered (also by the first author).<sup>1</sup> Since distance zeta functions are associated with arbitrary bounded (fractal) sets in Euclidean spaces of any dimension (see Definition 2.1), they clearly represent a valuable tool connecting the geometry of fractal sets with complex analysis. This interplay is described in [LapRaŽu1–7], where the foundations of the theory of fractal zeta functions have been laid. In this paper, by ‘fractal zeta functions’ we mean the following three classes of zeta functions: zeta functions of fractal strings (and, more generally, of fractal sprays), distance zeta functions and tube zeta functions of bounded subsets of  $\mathbb{R}^N$ , with  $N \geq 1$ , although some other classes may appear as well, like zeta functions of relative fractal drums in  $\mathbb{R}^N$  and spectral zeta functions; see §6 below and [LapRaŽu1, Chap. 4]. The theory of fractal zeta functions exhibits very interesting connections with the Minkowski contents and dimensions of fractal sets; see Theorems 2.6 and 2.8.

Like fractal string theory, which the present theory of fractal zeta functions extends to arbitrary dimensions (as well as to “relative fractal drums” in  $\mathbb{R}^N$ ), the work described here should eventually have applications to various aspects of harmonic analysis, fractal geometry, dynamical systems, geometric measure theory and analysis on nonsmooth spaces, number theory and arithmetic geometry, mathematical physics and, more speculatively, to aspects of condensed matter physics and cosmology. Some of the more mathematical applications of the theory are described in [LapRaŽu1], as well as in [LapRaŽu2–7], but a variety of potential applications remain to be explored or even imagined.

The basic property of the distance zeta function of a fractal set, described in Theorem 2.4, is that its abscissa of (absolute or Lebesgue) convergence is equal to the upper box dimension  $D$  of the set. Under some mild hypotheses,  $D$  is always a

---

<sup>1</sup>For fractal string theory and the associated one-dimensional theory of complex dimensions, as well as for the extensions to higher-dimensional fractal sprays (in the sense of [LapPo3]), we refer the reader to the research monographs [Lap-vFr1–3] along, for example, with the articles [Lap1–6, LapPo1–3, LapMa1–2, Fal1, HeLap, HamLap, Tep1–2, LapRo, LapLéRo, LéMen, LapLu, LapLu-vFr1–2, LapPe1–2, LapPeWi1–2, LalLap1–2, LapRoŽu, RaWi, ElLapMacRo, HerLap1–2]. We refer, in particular, to [Lap-vFr3, §12.2.1 and Chap. 13] for a survey of some of the recent developments of the theory, prior to [LapRaŽu1–7].

singularity; see part (b) of Theorem 2.4. Furthermore, assuming that  $D$  is a pole, then it is simple. Moreover, the residue of the distance zeta function computed at  $D$  is, up to a multiplicative constant, between the corresponding upper and lower Minkowski contents. A similar statement holds for the tube zeta function. (See Theorems 2.6 and 2.8, respectively.)

In addition, according to part (b) of Theorem 2.4, under some mild assumptions on a bounded set  $A$ , the abscissa of (Lebesgue, i.e., *absolute*) convergence of its distance zeta function coincides not only with  $D$ , but also with the abscissa of *holomorphic continuation* of the zeta function.

We stress that if  $D := \overline{\dim}_B A < N$ , all the results concerning the distance zeta functions have exact counterparts for the tube zeta functions, and vice versa. In other words, the fractal zeta functions introduced in [LapRaŽu1–7] contain essentially the same information. In practice, however, it is often the case that one of the fractal zeta functions is better suited for the given situation under consideration.

In §3, we discuss the existence and the construction of a suitable meromorphic continuation of the distance (or tube) zeta function of a fractal set, both in the Minkowski measurable case (Theorem 3.1) and a frequently encountered instance of Minkowski nonmeasurable case (Theorem 3.2). We will illustrate the latter situation by computing the fractal zeta function and the associated complex dimensions of the Sierpiński carpet; see Proposition 3.5 and Example 7.3 when  $N = 2$  or  $3$ , respectively. Many other examples are provided in [LapRaŽu1] and [LapRaŽu2–6], where are calculated, in particular, the complex dimensions of the higher-dimensional Sierpiński gaskets and carpets in  $\mathbb{R}^N$ , for any  $N \geq 2$ .

In §4, we introduce the so-called transcendently  $n$ -quasiperiodic sets, for any integer  $n \geq 2$  (that is, roughly speaking, the sets possessing  $n$  quasiperiods; see Definition 4.4), and describe the construction of 2-quasiperiodic sets, based on carefully chosen generalized Cantor sets with two parameters, introduced in Definition 4.1; see Theorem 4.5. It is also possible to construct  $n$ -quasiperiodic sets, for any  $n \geq 2$ , and even  $\infty$ -quasiperiodic sets, that is, sets which possess infinitely many quasiperiods; see §5 below and [LapRaŽu1, §4.6].

In §6, we introduce the notion of a relative fractal drum  $(A, \Omega)$  (which represents a natural extension of the notion of bounded fractal string and of bounded set). We also introduce the corresponding relative distance and tube zeta functions  $\zeta_A(\cdot, \Omega)$  and  $\tilde{\zeta}_A(\cdot, \Omega)$ , and study their properties. It is noteworthy that the relative box dimension  $\dim_B(A, \Omega)$  can be naturally defined as a real number, which may also assume negative values, including  $-\infty$ .

In §7, we address the question of reconstructing the tube function  $t \mapsto |A_t \cap \Omega|$  of a relative fractal drum  $(A, \Omega)$ , and thereby of obtaining a general “fractal tube formula” expressed in terms of the complex dimensions of  $(A, \Omega)$  (defined as the poles of a suitable meromorphic extension of the relative distance zeta function  $\zeta_A(\cdot, \Omega)$ ). The corresponding tube formulas are obtained in [LapRaŽu1, Chap. 5]

and [LapRaŽu6] (announced in [LapRaŽu5]), as well as illustrated by a variety of examples. The example of the three-dimensional Sierpiński carpet is given in Example 7.3. Moreover, towards the end of §7, we explain how to deduce from our general tube formulas (and significantly extend) earlier results obtained for fractal strings (in [Lap-vFr1–3]) and, especially, for fractal sprays and self-similar tilings (in [LapPe2] and [LapPeWi1]).

In closing this introduction, we recall some basic notation and terminology which will be needed in the sequel. First of all, in order to avoid trivial special cases, we assume implicitly that all bounded subsets of  $\mathbb{R}^N$  under consideration in the statements of the theorems are nonempty. Assume that  $A$  is a given bounded subset of  $\mathbb{R}^N$  and let  $r$  be a fixed real number. We define the *upper* and *lower  $r$ -dimensional Minkowski contents* of  $A$ , respectively, by

$$\mathcal{M}^{*r}(A) := \limsup_{t \rightarrow 0^+} \frac{|A_t|}{t^{N-r}}, \quad \mathcal{M}_*^r(A) := \liminf_{t \rightarrow 0^+} \frac{|A_t|}{t^{N-r}},$$

where  $A_t$  denotes the Euclidean  $t$ -neighborhood of  $A$  (namely,  $A_t := \{x \in \mathbb{R}^N : d(x, A) < t\}$ ) and  $|A_t|$  is the  $N$ -dimensional Lebesgue measure of  $A_t$ . The *upper* and *lower box* (or *Minkowski*) *dimensions* of  $A$  are then defined, respectively, by

$$\overline{\dim}_B A := \inf\{r \in \mathbb{R} : \mathcal{M}^{*r}(A) = 0\}, \quad \underline{\dim}_B A := \inf\{r \in \mathbb{R} : \mathcal{M}_*^r(A) = 0\}.$$

It is easy to check that  $0 \leq \underline{\dim}_B A \leq \overline{\dim}_B A \leq N$ . Furthermore, if  $A$  is such that  $\underline{\dim}_B A = \overline{\dim}_B A$ , then this common value is denoted by  $\dim_B A$  and is called the *box* (or *Minkowski*) *dimension* of  $A$ . Moreover, if  $A$  is such that, for some  $D \in [0, N]$ , we have  $0 < \mathcal{M}_*^D(A) \leq \mathcal{M}^{*D}(A) < \infty$  (in particular, then  $\dim_B A$  exists and  $D = \dim_B A$ ), we say that  $A$  is *Minkowski nondegenerate*. If  $\mathcal{M}_*^D(A) = \mathcal{M}^{*D}(A)$ , then this common value is denoted by  $\mathcal{M}^D(A)$  and called the *Minkowski content* of  $A$ . Finally, assuming that  $A$  is such that  $\mathcal{M}^D(A)$  exists and  $0 < \mathcal{M}^D(A) < \infty$ , we say that  $A$  is *Minkowski measurable*.<sup>2</sup>

Throughout this paper, given  $\alpha \in \mathbb{R} \cup \{\pm\infty\}$ , we denote by  $\{\operatorname{Re} s > \alpha\}$  the corresponding open right half-plane in the complex plane, defined by  $\{s \in \mathbb{C} : \operatorname{Re} s > \alpha\}$ . (In particular, if  $\alpha = \pm\infty$ ,  $\{\operatorname{Re} s > \alpha\}$  is equal to  $\emptyset$  or  $\mathbb{C}$ , respectively.) Similarly, given any  $\alpha \in \mathbb{R}$ , we denote by  $\{\operatorname{Re} s = \alpha\}$  the corresponding vertical line  $\{s \in \mathbb{C} : \operatorname{Re} s = \alpha\}$ .

---

<sup>2</sup>We note that the notion of Minkowski dimension was introduced (for noninteger values) by Bouligand [Bou] in the late 1920s (without making a clear distinction between the lower and upper limits), while the notions of (lower and upper) Minkowski content, Minkowski measurability and Minkowski nondegeneracy were introduced, respectively, in [Fed2], [Sta] and [Žu1]. (See also [Lap1, LapPo1] and, especially, [LapPo2–3], along with [Lap-vFr3], for the latter notions.) For general references on the notion of Minkowski (or box) dimension (from different points of view), we refer, for example, to [Fed2], [Fal2], [Tri], [Mat] and [Lap-vFr3].

## 2. Distance and tube zeta functions

Let us introduce a new class of zeta functions, defined by the first author in 2009, which extends the notion of geometric zeta functions of bounded fractal strings to bounded subsets of Euclidean spaces of arbitrary dimensions.

*Definition 2.1* ([LapRaŽu1,2]). Let  $A$  be a bounded subset of  $\mathbb{R}^N$  and let  $\delta$  be a fixed positive real number. Then, the *distance zeta function*  $\zeta_A$  of  $A$  is defined by

$$\zeta_A(s) := \int_{A_\delta} d(x, A)^{s-N} dx, \quad (1)$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large. Here,  $d(x, A) := \inf\{|x - y| : y \in A\}$  denotes the usual Euclidean distance from  $x$  to  $A$ . Furthermore, the integral is taken in the sense of Lebesgue, and hence, is absolutely convergent.<sup>3</sup>

*Remark 2.2.* Since the difference of any two distance zeta functions of the same set  $A$  corresponding to two different values of  $\delta$  is an entire function,<sup>4</sup> it follows that the dependence of the distance zeta function  $\zeta_A$  on  $\delta > 0$  is inessential, in the sense that the poles (of meromorphic extensions) of  $\zeta_A$ , as well as their multiplicities, do not depend on the choice of  $\delta$ .

The key for understanding the behavior of the distance zeta function  $\zeta_A$  consists in understanding the Lebesgue integrability of the function  $A_\delta \ni x \mapsto d(x, A)^{\operatorname{Re} s - N}$ , where  $s \in \mathbb{C}$  is fixed.<sup>5</sup> (We shall soon see that  $\operatorname{Re} s$  should be sufficiently large.) More precisely, we are interested in the Lebesgue integrability of the function  $x \mapsto d(x, A)^{-\gamma}$  defined on  $A_\delta$ , where  $\gamma := N - \operatorname{Re} s$  and  $s$  is a fixed complex number. Since the function is clearly bounded (and hence, integrable) for  $\gamma \leq 0$ , it suffices to consider the case when  $\gamma > 0$ , that is, when  $\operatorname{Re} s < N$ .

Let us recall a useful and little known result due to Harvey and Polking, stated implicitly on page 42 of [HarPol], in which a sufficient condition for Lebesgue integrability is expressed in terms of the upper box dimension. *If  $A$  is any nonempty bounded subset of  $\mathbb{R}^N$ , then the following implication holds:*<sup>6</sup>

$$\gamma < N - \overline{\dim}_B A \implies \int_{A_\delta} d(x, A)^{-\gamma} dx < \infty. \quad (2)$$

*Remark 2.3.* The sufficient condition for the Lebesgue (i.e., absolute) integrability of the function  $A_\delta \ni x \mapsto d(x, A)^{s-N}$  in the Harvey–Polking result in (2), becomes  $\gamma := N - \operatorname{Re} s < N - \overline{\dim}_B A$ , that is,  $\operatorname{Re} s > \overline{\dim}_B A$ . In other words,  $\zeta_A(s)$  is well defined for all  $s \in \mathbb{C}$  in the open right half-plane  $\{\operatorname{Re} s > \overline{\dim}_B A\}$ .

<sup>3</sup>For simplicity, we implicitly assume throughout this paper that  $|A| = 0$ ; the case when  $|A| > 0$  is discussed in [LapRaŽu1].

<sup>4</sup>This is an easy consequence of the fact that  $d(x, A) \in [\delta_1, \delta_2]$  for all  $x \in A_{\delta_2} \setminus A_{\delta_1}$  with  $0 < \delta_1 < \delta_2 < \infty$ .

<sup>5</sup>Indeed, note that  $|d(x, A)^{s-N}| = d(x, A)^{\operatorname{Re} s - N}$  for all  $x \in A_\delta$ .

<sup>6</sup>Moreover, if we assume that  $D := \dim_B A$  exists,  $D < N$  and  $\mathcal{M}_*^D(A) > 0$ , then the converse implication holds as well; see [Žu1, Thm. 4.3]. (See also [Žu2, Thm. 4.1(b)].)

The distance zeta function of a bounded set represents a natural extension of the notion of *geometric zeta function*  $\zeta_{\mathcal{L}}$ , associated with a bounded fractal string  $\mathcal{L} = (\ell_j)_{j \geq 1}$  (introduced by the first author and his collaborators<sup>7</sup> in the early 1990s and extensively studied in [Lap-vFr1–3] and the relevant references therein):

$$\zeta_{\mathcal{L}}(s) := \sum_{j=1}^{\infty} (\ell_j)^s, \quad (3)$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large. Here, a *bounded fractal string*  $\mathcal{L}$  is defined as a nonincreasing infinite sequence of positive real numbers  $(\ell_j)_{j \geq 1}$  such that  $\ell := \sum_{j \geq 1} \ell_j < \infty$ . Alternatively,  $\mathcal{L}$  can be viewed as a bounded open subset  $\Omega$  of  $\mathbb{R}$ , in which case the  $\ell_j$ s are the lengths of the connected components (open intervals) of  $\Omega$ , written in nonincreasing order (so that  $\ell_j \downarrow 0$  as  $j \rightarrow \infty$ ).

An important first result concerning  $\zeta_{\mathcal{L}}$  (first observed in [Lap2–3], using a result from [BesTay]) is that its abscissa of (absolute) convergence coincides with  $\bar{D}$  (the inner Minkowski dimension of  $\mathcal{L}$  or, equivalently, of its fractal boundary  $\partial\Omega$ ), defined by  $\bar{D} := \bar{\dim}_B(\partial\Omega, \Omega)$ ; see definition (41) below. For a direct proof of this statement, see [Lap-vFr3, Thm. 1.10] or [Lap-vFr3, Thm. 13.111] and [LapLu-vFr2]. In light of the next comment, it can be readily shown that part (a) of Theorem 2.4 below extends this result to arbitrary compact subsets of Euclidean spaces in any dimension; see [LapRaŽu1,2].

It is easy to see that the distance zeta function  $\zeta_{A_{\mathcal{L}}}$  of the set

$$A_{\mathcal{L}} := \left\{ a_k := \sum_{j=k}^{\infty} \ell_j : k \geq 1 \right\} \subseteq [0, \ell],$$

associated with  $\mathcal{L}$ , and the geometric zeta function  $\zeta_{\mathcal{L}}$  are connected by the following simple relation:

$$\zeta_{A_{\mathcal{L}}}(s) = u(s) \zeta_{\mathcal{L}}(s) + v(s), \quad (4)$$

for all complex numbers  $s$  such that  $\operatorname{Re} s$  is sufficiently large, where  $u$  and  $v$  are holomorphic on  $\mathbb{C} \setminus \{0\}$  and  $u$  is nowhere vanishing. In particular, due to Theorem 2.4 below, it follows that the abscissae of convergence of the distance zeta function  $\zeta_{A_{\mathcal{L}}}$  and of the geometric zeta function  $\zeta_{\mathcal{L}}$  coincide, and that the corresponding poles located on the critical line  $\{\operatorname{Re} s = \bar{\dim}_B A_{\mathcal{L}}\}$  (called *principal complex dimensions* of  $\mathcal{L}$  or, equivalently, of  $A_{\mathcal{L}}$ ), as well as their multiplicities, also coincide. The exact same results hold if  $A_{\mathcal{L}}$  is replaced by  $\partial\Omega$ , the boundary of  $\Omega$ , where  $\Omega$  is any geometric realization of  $\mathcal{L}$  by a bounded open subset of  $\mathbb{R}$ . For more details, see [LapRaŽu1,2].

Before stating Theorem 2.4, we need to introduce some terminology and notation, which will also be used in the remainder of the paper.

Given a meromorphic function (or, more generally, an arbitrary complex-valued function)  $f = f(s)$ , initially defined on some domain  $U \subseteq \mathbb{C}$ , we denote by  $D_{\text{hol}}(f)$  the unique extended real number (i.e.,  $D_{\text{hol}}(f) \in \mathbb{R} \cup \{\pm\infty\}$ ) such that

<sup>7</sup>See, especially, [Lap2–3], [LapPo1–3], [LapMa1–2] and [HeLap].

$\{\operatorname{Re} s > D_{\text{hol}}(f)\}$  is the *maximal* open right half-plane (of the form  $\{\operatorname{Re} s > \alpha\}$ , for some  $\alpha \in \mathbb{R} \cup \{\pm\infty\}$ ) to which the function  $f$  can be holomorphically extended.<sup>8</sup> This maximal (i.e., largest) half-plane is denoted by  $\mathcal{H}(f)$  and called the *half-plane of holomorphic continuation* of  $f$ .

If, in addition, the function  $f = f(s)$  is assumed to be given by a *tamed Dirichlet-type integral* (or DTI, in short),<sup>9</sup> of the form

$$f(s) := \int_E \varphi(x)^s d\mu(x), \quad (5)$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large, where  $\mu$  is a (positive or complex) local (i.e., locally bounded) Borel measure on a given (measurable) space  $E$  and

$$0 \leq \varphi(x) \leq C \quad \text{for } |\mu|\text{-a.e. } x \in E, \quad (6)$$

where  $C \geq 0$ ,<sup>10</sup> then  $D(f)$ , the *abscissa of (absolute or Lebesgue) convergence* of  $f$ , is defined as the unique extended real number (i.e.,  $D(f) \in \mathbb{R} \cup \{\pm\infty\}$ ) such that  $\{\operatorname{Re} s > D(f)\}$  is the *maximal* open right half-plane (of the form  $\{\operatorname{Re} s > \alpha\}$ , for some  $\alpha \in \mathbb{R} \cup \{\pm\infty\}$ ) on which the Lebesgue integral initially defining  $f$  in (5) is convergent (or, equivalently, is absolutely convergent), with  $\mu$  replaced by  $|\mu|$ , the total variation measure of  $\mu$ . (Recall that  $|\mu| = \mu$  if  $\mu$  is positive.) In short,  $D(f)$  is called the *abscissa of convergence* of  $f$ . Furthermore, the aforementioned maximal right half-plane is denoted by  $\Pi(f)$  and is called the *half-plane of (absolute or Lebesgue) convergence* of (the Dirichlet-type integral)  $f$ . It is shown in [LapRaŽu1, §2.1] that under mild hypotheses (which are always satisfied in our setting),  $D(f)$  is well defined and (with the notation of (5) just above) we have, equivalently:<sup>11</sup>

$$D(f) = \inf \left\{ \alpha \in \mathbb{R} : \int_E \varphi(x)^\alpha d|\mu|(x) < \infty \right\}, \quad (7)$$

where (as above)  $|\mu|$  is the total variation (local) measure of  $\mu$ . Under the stated conditions on  $f$ , we have  $\Pi(f) \subseteq \mathcal{H}(f)$ ; that is,  $-\infty \leq D_{\text{hol}}(f) \leq D(f) \leq +\infty$ .

Note that the distance zeta function  $\zeta_A$ , defined by (1), is a tamed DTI of the form (5), with  $E := A_\delta$ ,  $\varphi(x) := d(x, A)$  and  $d\mu(x) := d(x, A)^{-N} dx$ . Furthermore, we can clearly take  $C := \delta$  in (6).

The following key result describes some of the basic properties of distance zeta functions.

<sup>8</sup>By using the principle of analytic continuation, it is easy to check that  $D_{\text{hol}}(f)$  and  $\mathcal{H}(f)$  are well defined; see [LapRaŽu1, §2.1].

<sup>9</sup>This is the case of the classic (generalized) Dirichlet series and integrals [Ser, Pos], the classic arithmetic zeta functions (see, e.g., [Lap-vFr3, App. A] and [Lap 4, Apps. B, C & E]), as well as of the geometric zeta functions of fractal strings studied in [Lap-vFr1–3] and of all the fractal zeta functions considered in this paper and in [LapRaŽu1–7].

<sup>10</sup>Such functions  $f$  are called *tamed* DTIs in [LapRaŽu1–7]; see esp. [LapRaŽu1, App. A] for a development of their general theory.

<sup>11</sup>Let  $D := \overline{\dim}_B A$ , for brevity. In light of Theorem 2.4, for this alternative definition of  $D(\zeta_A)$  (or of  $D(\tilde{\zeta}_A)$ ), with  $A \subseteq \mathbb{R}^N$  bounded (as in the present situation), it would suffice to restrict oneself to  $\alpha \geq 0$  in the right-hand side of (7); this follows since  $D(\zeta_A) = \overline{\dim}_B A \geq 0$  and (if  $D < N$ ),  $D(\zeta_A) = D(\tilde{\zeta}_A)$ . Here,  $\tilde{\zeta}_A$  stands for the tube zeta function of  $A$ , defined by Equation (12).

**Theorem 2.4** ([LapRaŽu1,2]). *Let  $A$  be an arbitrary bounded subset of  $\mathbb{R}^N$  and let  $\delta$  be a fixed positive real number. Then:*

(a) *The distance zeta function  $\zeta_A$  is holomorphic on  $\{\operatorname{Re} s > \overline{\dim}_B A\}$ . Moreover,  $\Pi(\zeta_A) = \{\operatorname{Re} s > \overline{\dim}_B A\}$ ; that is,*

$$D(\zeta_A) = \overline{\dim}_B A. \quad (8)$$

(b) *If the box (or Minkowski) dimension  $D := \dim_B A$  exists,  $D < N$  and  $\mathcal{M}_*^D(A) > 0$ , then  $\zeta_A(s) \rightarrow +\infty$  as  $s \in \mathbb{R}$  converges to  $D$  from the right. In particular,  $\mathcal{H}(\zeta_A) = \Pi(\zeta_A) = \{\operatorname{Re} s > \dim_B A\}$ ; that is,*

$$D_{\text{hol}}(\zeta_A) = D(\zeta_A) = \dim_B A. \quad (9)$$

*Remark 2.5.* (a) It would be of interest to construct (if possible) a class of non-trivial bounded subsets  $A$  of  $\mathbb{R}^N$  such that  $D_{\text{hol}}(\zeta_A) < D(\zeta_A)$ . A trivial example is given by  $A = [0, 1]$ , since then  $D_{\text{hol}}(\zeta_A) = 0$  and  $D(\zeta_A) = 1$ .

(b) The analog of Theorem 2.4 holds for the tube zeta function  $\tilde{\zeta}_A$  (to be introduced in Definition 2.7 below), except for the fact that in part (b), one no longer needs to assume that  $D < N$ .

Given a bounded set  $A$ , it is of interest to know the corresponding poles of the associated distance zeta function  $\zeta_A$ , meromorphically extended (if possible) to a neighborhood of the critical line  $\{\operatorname{Re} s = D(\zeta_A)\}$ . Following the terminology of [Lap-vFr3], these poles are called the *complex dimensions* of  $A$  and we denote the resulting set of complex dimensions by  $\mathcal{P}(\zeta_A)$ .<sup>12</sup> We pay particular attention to the set of complex dimensions of  $A$  located on the critical line  $\{\operatorname{Re} s = D(\zeta_A)\}$ , which we call the set of *principal complex dimensions* of  $A$  and denote by  $\dim_{PC} A$ .

For example, it is well known that for the ternary Cantor set  $C^{(1/3)}$ ,  $\dim_B C^{(1/3)} = \log_3 2$  and, moreover (see [Lap-vFr3, §1.2.2 and §2.3.1]), with  $\omega := \sqrt{-1}$ ,

$$\dim_{PC} C^{(1/3)} := \log_3 2 + \frac{2\pi}{\log 3} \mathbb{Z}.$$

The following result provides an interesting connection between the residue of the distance zeta function of a fractal set at  $D := \dim_B A$  and its Minkowski contents.

**Theorem 2.6** ([LapRaŽu1,2]). *Assume that  $A$  is a bounded subset of  $\mathbb{R}^N$  which is nondegenerate (that is,  $0 < \mathcal{M}_*^D(A) \leq \mathcal{M}^{*D}(A) < \infty$  and, in particular,  $\dim_B A = D$ ), and  $D < N$ . If the distance zeta function  $\zeta_A(\cdot, A_\delta) := \zeta_A$ , initially defined*

<sup>12</sup>Strictly speaking, one should talk about the set  $\mathcal{P}(\zeta_A, U)$  of *visible complex dimensions* relative to a domain  $U \subseteq \mathbb{C}$  to which  $\zeta_A$  can be meromorphically extended; see [LapRaŽu1-4] (along with [Lap-vFr3]). In the examples described in this paper, we have  $U := \mathbb{C}$ .

by (1), can be meromorphically extended<sup>13</sup> to a neighborhood of  $s = D$ , then  $D$  is necessarily a simple pole of  $\zeta_A(\cdot, A_\delta)$ , and

$$(N - D) \mathcal{M}_*^D(A) \leq \operatorname{res}(\zeta_A(\cdot, A_\delta), D) \leq (N - D) \mathcal{M}^{*D}(A). \quad (10)$$

Furthermore, the value of  $\operatorname{res}(\zeta_A(\cdot, A_\delta), D)$  does not depend on  $\delta > 0$ . In particular, if  $A$  is Minkowski measurable, then

$$\operatorname{res}(\zeta_A(\cdot, A_\delta), D) = (N - D) \mathcal{M}^D(A). \quad (11)$$

The distance zeta function defined by (1) is closely related to the tube zeta function of a fractal set, which, in turn, is defined via the tube function  $t \mapsto |A_t|$ , for  $t > 0$ , of the fractal set  $A$ , as we now explain.

*Definition 2.7* ([LapRaŽu1,2]). Let  $\delta$  be a fixed positive number, and let  $A$  be a bounded subset of  $\mathbb{R}^N$ . Then, the *tube zeta function* of  $A$ , denoted by  $\tilde{\zeta}_A$ , is defined (for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large) by

$$\tilde{\zeta}_A(s) := \int_0^\delta t^{s-N-1} |A_t| dt. \quad (12)$$

For any fixed positive real number  $\delta > 0$ , the distance and tube zeta functions associated with a given fractal set  $A$  are connected as follows:<sup>14</sup>

$$\zeta_A(s, A_\delta) = \delta^{s-N} |A_\delta| + (N - s) \tilde{\zeta}_A(s, \delta), \quad (13)$$

for  $\operatorname{Re} s > \overline{\dim}_B A$ ,<sup>15</sup> see [LapRaŽu1,2].<sup>16</sup> Using this result, it is easy to obtain the analog of Theorem 2.4 for  $\tilde{\zeta}_A$  (as was stated in Remark 2.5(b) above) and to reformulate Theorem 2.6 in terms of the tube zeta functions. In particular, we conclude that the residue of the tube zeta function of a fractal set, computed at  $s = D$ , is equal to its Minkowski content, provided the set is Minkowski measurable.

**Theorem 2.8** ([LapRaŽu1,2]). *Assume that  $A$  is a nondegenerate bounded subset of  $\mathbb{R}^N$  (so that  $D := \dim_B A$  exists), and there exists a meromorphic extension of  $\tilde{\zeta}_A$  to a neighborhood of  $D$ . Then,  $D$  is a simple pole of  $\tilde{\zeta}_A$ , and for any positive  $\delta$ ,  $\operatorname{res}(\tilde{\zeta}_A, D)$  is independent of  $\delta$ . Furthermore, we have*

$$\mathcal{M}_*^D(A) \leq \operatorname{res}(\tilde{\zeta}_A, D) \leq \mathcal{M}^{*D}(A). \quad (14)$$

In particular, if  $A$  is Minkowski measurable, then

$$\operatorname{res}(\tilde{\zeta}_A, D) = \mathcal{M}^D(A). \quad (15)$$

<sup>13</sup>The existence and construction of meromorphic extensions of fractal zeta functions is discussed in §3. It is studied in a variety of situations in [LapRaŽu1–4,6].

<sup>14</sup>We write here  $\zeta_A(\cdot, A_\delta) := \zeta_A$  and  $\tilde{\zeta}_A(\cdot, \delta) := \tilde{\zeta}_A$ , for emphasis.

<sup>15</sup>In light of the principle of analytic continuation, one deduces that identity (13) continues to hold whenever one (and hence, both) of the fractal zeta functions  $\zeta_A$  and  $\tilde{\zeta}_A$  is meromorphic on a given domain  $U \subseteq \mathbb{C}$ .

<sup>16</sup>The case when  $D = N$  in Theorem 2.8 must be treated separately.

A class of fractal sets  $A$  for which we have strict inequalities in (14) (and hence also in (10) of Theorem 2.6 above) is constructed in Theorem 3.2; see (23).

### 3. Meromorphic extensions of fractal zeta functions

Since the definition of the set of principal complex dimensions  $\dim_{PC} A$  of  $A$  requires the existence of a suitable meromorphic extension of the distance zeta function  $\zeta_A$ , it is natural to study this issue in more detail. For simplicity, we formulate the results of this section for  $\tilde{\zeta}_A$ , but we note that the analogs of Theorems 3.1 and 3.2 also hold for  $\zeta_A$ , provided  $D < N$ ; see [LapRaŽu1, §2.3.3] or [LapRaŽu3].

**Theorem 3.1 (Minkowski measurable case, [LapRaŽu1,3]).** *Let  $A$  be a bounded subset of  $\mathbb{R}^N$  such that there exist  $\alpha > 0$ ,  $\mathcal{M} \in (0, +\infty)$  and  $D \geq 0$  satisfying*

$$|A_t| = t^{N-D} (\mathcal{M} + O(t^\alpha)) \quad \text{as } t \rightarrow 0^+. \quad (16)$$

*Then,  $\dim_B A$  exists and  $\dim_B A = D$ . Furthermore,  $A$  is Minkowski measurable with Minkowski content  $\mathcal{M}^D(A) = \mathcal{M}$ . Moreover, the tube zeta function  $\tilde{\zeta}_A$  has for abscissa of convergence  $D(\tilde{\zeta}_A) = \dim_B A = D$  and possesses a (necessarily unique) meromorphic continuation (still denoted by  $\tilde{\zeta}_A$ ) to (at least) the open right half-plane  $\{\operatorname{Re} s > D - \alpha\}$ . The only pole of  $\tilde{\zeta}_A$  in this half-plane is  $s = D$ ; it is simple and, moreover,  $\operatorname{res}(\tilde{\zeta}_A, D) = \mathcal{M}$ .*

Next, we deal with a useful class of Minkowski nonmeasurable sets. Before stating Theorem 3.2, let us first introduce some notation. Given a locally integrable  $T$ -periodic function  $G : \mathbb{R} \rightarrow \mathbb{R}$ , with  $T > 0$ , we denote by  $G_0$  its truncation to  $[0, T]$ , while the Fourier transform of  $G_0$  is denoted by  $\hat{G}_0$ : for all  $t \in \mathbb{R}$ ,

$$\hat{G}_0(t) := \int_{-\infty}^{+\infty} e^{-2\pi t\tau} G_0(\tau) d\tau = \int_0^T e^{-2\pi t\tau} G(\tau) d\tau. \quad (17)$$

**Theorem 3.2 (Minkowski nonmeasurable case, [LapRaŽu1,3]).** *Let  $A$  be a bounded subset of  $\mathbb{R}^N$  such that there exist  $D \geq 0$ ,  $\alpha > 0$ , and  $G : \mathbb{R} \rightarrow (0, +\infty)$  a nonconstant periodic function with period  $T > 0$ , satisfying*

$$|A_t| = t^{N-D} (G(\log t^{-1}) + O(t^\alpha)) \quad \text{as } t \rightarrow 0^+. \quad (18)$$

*Then  $G$  is continuous,  $\dim_B A$  exists and  $\dim_B A = D$ . Furthermore,  $A$  is Minkowski nondegenerate, with upper and lower Minkowski contents respectively given by*

$$\mathcal{M}_*^D(A) = \min G, \quad \mathcal{M}^{*D}(A) = \max G. \quad (19)$$

*Moreover, the tube zeta function  $\tilde{\zeta}_A$  has for abscissa of convergence  $D(\tilde{\zeta}_A) = D$  and possesses a (necessarily unique) meromorphic extension (still denoted by  $\tilde{\zeta}_A$ ) to (at least) the half-plane  $\{\operatorname{Re} s > D - \alpha\}$ .*

In addition, the set of principal complex dimensions of  $A$  is given by

$$\dim_{PC} A = \left\{ s_k = D + \frac{2\pi}{T} k : \hat{G}_0\left(\frac{k}{T}\right) \neq 0, k \in \mathbb{Z} \right\} \quad (20)$$

(see (17)) and there are no other complex dimensions in  $\{\operatorname{Re} s > D - \alpha\}$ ; they are all simple, and the residue at each  $s_k \in \dim_{PC} A$ , with  $k \in \mathbb{Z}$ , is given by

$$\operatorname{res}(\tilde{\zeta}_A, s_k) = \frac{1}{T} \hat{G}_0\left(\frac{k}{T}\right). \quad (21)$$

If  $s_k \in \dim_{PC} A$ , then  $s_{-k} \in \dim_{PC} A$  (in agreement with the ‘reality principle’), and  $|\operatorname{res}(\tilde{\zeta}_A, s_k)| \leq \frac{1}{T} \int_0^T G(\tau) d\tau$ ; furthermore,  $\lim_{k \rightarrow \pm\infty} \operatorname{res}(\tilde{\zeta}_A, s_k) = 0$ .

Moreover, the set of principal complex dimensions of  $A$  contains  $s_0 = D$ , and

$$\operatorname{res}(\tilde{\zeta}_A, D) = \frac{1}{T} \int_0^T G(\tau) d\tau. \quad (22)$$

In particular,  $A$  is not Minkowski measurable and

$$\mathcal{M}_*^D(A) < \operatorname{res}(\tilde{\zeta}_A, D) < \mathcal{M}^{*D}(A). \quad (23)$$

*Example 3.3* ( $a$ -strings). The compact set  $A := \{j^{-a} : j \in \mathbb{N}\} \cup \{0\}$ , where  $a > 0$ , is Minkowski measurable and

$$\mathcal{M}^D(A) = \frac{2^{1-D}}{1-D} a^D, \quad D := \dim_B A = \frac{1}{1+a}. \quad (24)$$

(See [Lap 1, Exple. 5.1 and App. C].) The associated fractal string  $\mathcal{L} = (\ell_j)_{j \geq 1}$ , defined by  $\ell_j = j^{-a} - (j+1)^{-a}$  for all  $j \geq 1$  (or, equivalently, by  $\Omega := [0, 1] \setminus A \subset \mathbb{R}$ , so that  $\partial\Omega = A$ ), is called the  $a$ -string; see [Lap1–3], [LapPo1–2], [HeLap] and [Lap-vFr3, §6.5.1]. In light of (11) and (15), we then know that  $\operatorname{res}(\zeta_A(\cdot, A_\delta), D) = (1-D)\mathcal{M}^D(A)$  and  $\operatorname{res}(\tilde{\zeta}_A, D) = \mathcal{M}^D(A)$ .

*Example 3.4* (fractal nests). Let  $a > 0$  and let  $A$  be the countable union of concentric circles in  $\mathbb{R}^2$ , centered at the origin and of radii  $r = k^{-a}$ , where  $k \in \mathbb{N}$ . According to the terminology introduced in [LapRaŽu1–4],  $A$  is called the *fractal nest of inner type* generated by the  $a$ -string from the preceding example. Then, using the the distance zeta function of  $A$  it is possible to show that

$$D := \overline{\dim}_B A = \max\left\{1, \frac{2}{1+a}\right\}. \quad (25)$$

(See [LapRaŽu1, Chap. 3] and [LapRaŽu2–4].) The set  $A$  is closely related to the planar spiral  $\Gamma$  defined in polar coordinates by  $r = \theta^{-a}$ ,  $\theta \geq \theta_0$ , where  $\theta_0 > 0$ , and the value of  $\dim_B \Gamma$  is the same as for  $A$ ; see [Tri]. We mention in passing that for  $a \neq 1$ , the fractal nest  $A$  (as well as the corresponding spiral  $\Gamma$ ) is Minkowski measurable and for every  $a \in (0, 1)$ , the value of its Minkowski content is independent of  $\theta_0$  and given by

$$\mathcal{M}^D(A) = \mathcal{M}^D(\Gamma) = \pi(\pi a)^{-2a/(1+a)} \frac{1+a}{1-a}. \quad (26)$$

(See [Žu1].) Using (26), along with with Eq. (11) from Theorem 2.6, we conclude that the residue of the distance zeta function  $\zeta_A$ , as well as of  $\zeta_\Gamma$ , computed at  $s = D$ , is given by

$$\operatorname{res}(\zeta_A, D) = \operatorname{res}(\zeta_\Gamma, D) = \pi(\pi a)^{-2a/(1+a)} \frac{2a}{1-a}, \quad (27)$$

provided  $a \in (0, 1)$ . For  $a = 1$ , we have  $\mathcal{M}^1(A) = \mathcal{M}^1(\Gamma) = +\infty$ . These and related results are useful in the study of fractal properties of spiral trajectories of planar vector fields; see, e.g., [ŽuŽup].

More generally, if we consider the fractal nest  $A_N$  defined as the countable union of concentric spheres in  $\mathbb{R}^N$ , centered at the origin and of radii  $r = k^{-\alpha}$ , where  $k \in \mathbb{N}$ , then using the distance zeta function  $\zeta_{A_N}$ , it can be shown (see [LapRaŽu1, §3.4] and [LapRaŽu2-4]) that

$$\overline{\dim}_B A_N = \max \left\{ N - 1, \frac{N}{1 + a} \right\}. \quad (28)$$

Note that for  $N = 1$  and  $N = 2$ , we recover the box dimension of the  $a$ -string and of the fractal nest, respectively; see [LapRaŽu2-4] and Eqs. (24)–(25) above.

In the following result, we provide the distance zeta function of the Sierpiński carpet and the corresponding principal complex dimensions. It is well known that the Sierpiński carpet is not Minkowski measurable. See, e.g., [Lap-vFr3], as well as [HorŽu] for explicit values of its upper and lower Minkowski contents. A similar result can be obtained for the Sierpiński gasket (and its higher-dimensional analogs); see [LapRaŽu1, §3.2.2] and [LapRaŽu2-4].

**Proposition 3.5 (Distance zeta function of the Sierpiński carpet).** *Let  $A$  be the Sierpiński carpet in  $\mathbb{R}^2$ , constructed in the usual way inside the unit square. Let  $\delta$  be a fixed positive real number. We assume without loss of generality that  $\delta > 1/6$  (so that for this choice of  $\delta$ ,  $A_\delta$  coincides with the  $\delta$ -neighborhood of the unit square  $[0, 1]^2$ ). Then, for all  $s \in \mathbb{C}$ , the distance zeta function  $\zeta_A$  of the Sierpiński carpet is given by*

$$\zeta_A(s) = \frac{8}{2^s s(s-1)(3^s-8)} + 2\pi \frac{\delta^s}{s} + 4 \frac{\delta^{s-1}}{s-1}, \quad (29)$$

which is meromorphic on the whole complex plane. In particular, the set of complex dimensions and of principal complex dimensions of the Sierpiński carpet are given, respectively, by

$$\mathcal{P}(\zeta_A) = \{0, 1\} \cup \dim_{PC} A, \quad \dim_{PC} A = \log_3 8 + \frac{2\pi}{\log 3} \mathbb{Z}. \quad (30)$$

Furthermore, each of the complex dimensions (i.e., each of the poles of  $\zeta_A$ ) is simple. Moreover, the residues of the distance zeta function  $\zeta_A$  computed at the principal poles  $s_k := \log_3 8 + \frac{2\pi}{\log 3} k$ , with  $k \in \mathbb{Z}$ , are given by

$$\operatorname{res}(\zeta_A, s_k) = \frac{2^{-s_k}}{(\log 3) s_k (s_k - 1)}. \quad (31)$$

Finally, the approximate values of the lower and upper  $D$ -dimensional Minkowski contents are given by  $\mathcal{M}_*^D(A) \approx 1.350670$  and  $\mathcal{M}^{*D}(A) \approx 1.355617$ . (The precise values can be found in [HorŽu].)

*Sketch of the proof.* In order to compute the distance zeta function

$$\zeta_A(s) := \int_{A_\delta} d((x, y), A)^{s-2} dx dy$$

of the Sierpiński carpet  $A$ , we first have to calculate

$$\zeta_{A_k}(s, \Omega_k) := \int_{\Omega_k} d((x, y), A_k) dx dy, \quad (32)$$

where  $\Omega_k$  is a square of the  $k$ -th generation (of side lengths  $a_k = 3^{-k}$ ) and  $A_k$  is its boundary. (Here, we deal in fact with ‘relative distance zeta functions’, which are discussed in Remark 3.6 just below; see (42) and [LapRaŽu1–4].) This can be easily done by splitting  $\Omega_k$  into the disjoint union of eight congruent right-angle triangles, and we obtain after a short computation that  $\zeta_{A_k}(s, \Omega_k) = 8 \cdot 2^{-s} a_k^s s^{-1} (s-1)^{-1}$ . Since the  $k$ -th generation consists of  $8^{k-1}$  squares congruent to  $\Omega_k$ , we deduce that

$$\zeta_A(s, [0, 1]^2) = \sum_{k=1}^{\infty} 8^{k-1} \zeta_{A_k}(s, \Omega_k) = \frac{8}{2^s s (s-1) (3^s - 8)}, \quad (33)$$

for  $\operatorname{Re} s > \log_3 8$ . The last expression in (33) is meromorphic in all of  $\mathbb{C}$ . Hence, upon analytic continuation,  $\zeta_A(s, [0, 1]^2)$  is given by that expression for all  $s \in \mathbb{C}$ . Note that the value of  $\zeta_A(s, [0, 1]^2)$  is precisely equal to the first term on the right-hand side of (29). The remaining two terms are obtained by considering  $\zeta_A(s, A_\delta \setminus [0, 1]^2)$ , which can be easily reduced to considering a disk  $B_\delta(0)$  of radius  $\delta$  with respect to its origin  $0 \in \mathbb{R}^2$ , and two rectangles that are congruent to  $\Omega_0 := (0, 1) \times (-\delta, \delta)$  with respect to its middle section  $A_0 := (0, 1) \times \{0\}$ .  $\square$

*Remark 3.6.* Eq. (32) is a very special case of the zeta function of a *relative fractal drum*  $(A, \Omega)$  in  $\mathbb{R}^N$ , a notion which will be briefly discussed in §6 and is the object of [LapRaŽu4] and [LapRaŽu1, Chap. 4]; see the first equality in Eq. (42) below.

#### 4. Transcendentally quasiperiodic sets

In this section, we define a class of *quasiperiodic fractal sets*. The simplest of such sets has two incommensurable periods. Moreover, using suitable generalized Cantor sets, it is possible to ensure that the quotient of their periods be a transcendental real number. Our construction of such sets is based on a class of generalized Cantor sets with two parameters, which we now introduce.

*Definition 4.1* ([LapRaŽu1,2]). The generalized Cantor sets  $C^{(m,a)}$  are determined by an integer  $m \geq 2$  and a real number  $a \in (0, 1/m)$ . In the first step of the analog of Cantor's construction, we start with  $m$  equidistant, closed intervals in  $[0, 1]$  of length  $a$ , with  $m-1$  'holes', each of length  $(1-ma)/(m-1)$ . In the second step, we continue by scaling by the factor  $a$  each of the  $m$  intervals of length  $a$ ; and so on, ad infinitum. The (two-parameter) *generalized Cantor set*  $C^{(m,a)}$  is then defined as the intersection of the decreasing sequence of compact sets constructed in this way. It is easy to check that  $C^{(m,a)}$  is a perfect, uncountable compact subset of  $\mathbb{R}$ ; furthermore,  $C^{(m,a)}$  is also self-similar. For  $m = 2$ , the sets  $C^{(m,a)}$  are denoted by  $C^{(a)}$ . The classic ternary Cantor set is obtained as  $C^{(2,1/3)}$ . In order to avoid any possible confusion, we note that the generalized Cantor sets introduced here are different from the generalized Cantor strings introduced and studied in [Lap-vFr3, Chap. 10], as well as used in a key manner in [Lap-vFr3, Chap. 11].

We collect some of the basic properties of generalized Cantor sets in the following proposition.

**Proposition 4.2 (Generalized Cantor sets, [LapRaŽu1,2]).** *If  $A := C^{(m,a)}$  is the generalized Cantor set introduced in Definition 4.1, where  $m$  is an integer larger than 1, and  $a \in (0, 1/m)$ , then*

$$D := \dim_B A = D(\zeta_A) = \log_{1/a} m. \quad (34)$$

Furthermore, the tube formula associated with  $A$  is given by

$$|A_t| = t^{1-D} G(\log t^{-1}) \quad \text{for all } t \in (0, t_0), \quad (35)$$

where  $t_0$  is a suitable positive constant and  $G = G(\tau)$  is a continuous, positive and nonconstant periodic function, with minimal period  $T = \log(1/a)$ .

Moreover,  $A$  is Minkowski nondegenerate and Minkowski nonmeasurable; that is,  $0 < \mathcal{M}_*^D(A) < \mathcal{M}^{*D}(A) < \infty$ .<sup>17</sup>

Finally, the distance zeta function of  $A$  admits a meromorphic continuation to all of  $\mathbb{C}$  and the set of principal complex dimensions of  $A$  is given by

$$\dim_{PC} A = D + \frac{2\pi}{T} \mathbb{Z}. \quad (36)$$

Besides  $(\dim_{PC} A) \cup \{0\}$ , there are no other poles, and all of the poles of  $\zeta_A$  are simple. In particular,  $\mathcal{P}(\zeta_A) = (D + \frac{2\pi}{T} \mathbb{Z}) \cup \{0\}$ .

The definition of quasiperiodic sets is based on the following notion of quasiperiodic functions, which will be useful for our purposes.<sup>18</sup>

<sup>17</sup>The periodic function  $G = G(\tau)$ , as well as the values of  $\mathcal{M}_*^D(A)$  and  $\mathcal{M}^{*D}(A)$ , can be explicitly computed; see [LapRaŽu1, §3.1.1]

<sup>18</sup>We note that Definition 4.3, although rather close to the one provided in [Vin], is very different from the usual definition of Bohr-type quasiperiodic functions.

*Definition 4.3* ([LapRaŽu1,2]). We say that a function  $G = G(\tau) : \mathbb{R} \rightarrow \mathbb{R}$  is *transcendentally  $n$ -quasiperiodic*, with  $n \geq 2$ , if it is of the form  $G(\tau) = H(\tau, \dots, \tau)$ , where  $H : \mathbb{R}^n \rightarrow \mathbb{R}$  is a function that is nonconstant and  $T_k$ -periodic in its  $k$ -th component, for each  $k = 1, \dots, n$ , and the periods  $T_1, \dots, T_n$  are algebraically (and hence, rationally) independent.<sup>19</sup> The positive numbers  $T_i$  ( $i = 1, \dots, n$ ) are called the *quasiperiods* of  $G$ . If, instead, the set of quasiperiods  $\{T_1, \dots, T_n\}$  is rationally independent and algebraically dependent, we say that  $G$  is *algebraically  $n$ -quasiperiodic*.

*Definition 4.4* ([LapRaŽu1,2]). Given a bounded subset  $A$  of  $\mathbb{R}^N$ , we say that a function  $G : \mathbb{R} \rightarrow \mathbb{R}$  is *associated with the set  $A$*  (or *corresponds to  $A$* ) if it is nonnegative and  $A$  has the following tube formula:

$$|A_t| = t^{N-D}(G(\log(1/t)) + o(1)) \text{ as } t \rightarrow 0^+, \quad (37)$$

with  $0 < \liminf_{\tau \rightarrow +\infty} G(\tau) \leq \limsup_{\tau \rightarrow +\infty} G(\tau) < \infty$ . In addition, we say that  $A$  is a *transcendentally* (resp., *algebraically*)  *$n$ -quasiperiodic set* if the function  $G = G(\tau)$  is transcendentally (resp., algebraically)  $n$ -quasiperiodic. The smallest possible value of  $n$  is called the *order of quasiperiodicity* of  $A$ .

The following result, which has a variety of generalizations as will be briefly explained below, provides a construction of transcendentally 2-quasiperiodic fractal sets. Its proof is based on the classical Gel'fond–Schneider theorem (as described in [Gel]) from transcendental number theory.

**Theorem 4.5** ([LapRaŽu1,2]). *Let  $C^{(m_1, a_1)}$  and  $\dim_B C^{(m_2, a_2)}$  be two generalized Cantor sets such that their box dimensions coincide and are equal to  $D \in (0, 1)$ . Assume that  $I_1$  and  $I_2$  are two unit closed intervals of  $\mathbb{R}$ , with disjoint interiors, and define  $A_1 := (\min I_1) + C^{(m_1, a_1)} \subset I_1$  and  $A_2 := (\min I_2) + C^{(m_2, a_2)} \subset I_2$ . Let  $\{p_1, p_2, \dots, p_k\}$  be the set of all distinct prime factors of  $m_1$  and  $m_2$ , and write*

$$m_1 = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_k^{\alpha_k}, \quad m_2 = p_1^{\beta_1} p_2^{\beta_2} \dots p_k^{\beta_k},$$

where  $\alpha_i, \beta_i \in \mathbb{N} \cup \{0\}$  for  $i = 1, \dots, k$ . If the exponent vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$  of, respectively,  $m_1$  and  $m_2$ , defined by

$$\mathbf{e}_1 := (\alpha_1, \alpha_2, \dots, \alpha_k) \quad \text{and} \quad \mathbf{e}_2 := (\beta_1, \beta_2, \dots, \beta_k),$$

are linearly independent over the field of rational numbers, then the compact set  $A := A_1 \cup A_2 \subset \mathbb{R}$  is *transcendentally 2-quasiperiodic*.

Moreover, the distance zeta function  $\zeta_A$  can be meromorphically extended to the whole complex plane, and we have that  $D(\zeta_A) = D$ . The set  $\dim_{PC} A$  of principal complex dimensions of  $A$  is given by

$$\dim_{PC} A = D + \left( \frac{2\pi}{T_1} \mathbb{Z} \cup \frac{2\pi}{T_2} \mathbb{Z} \right). \quad (38)$$

<sup>19</sup>That is, linearly independent over the field of algebraic numbers.

Besides  $(\dim_{PC} A) \cup \{0\}$ , there are no other poles of the distance zeta function  $\zeta_A$  and they are all simple. In particular,

$$\mathcal{P}(\zeta_A) = \left( D + \left( \frac{2\pi}{T_1} \mathbb{Z} \cup \frac{2\pi}{T_2} \mathbb{Z} \right) \right) \cup \{0\}. \quad (39)$$

*Remark 4.6.* This result can be considerably extended by using Baker's theorem [Ba, Thm. 2.1] which, in turn, is a far-reaching extension of the aforementioned Gel'fond–Schneider's theorem. Indeed, for any fixed integer  $n \geq 2$ , using Baker's theorem and  $n$  generalized Cantor sets, an explicit construction of a class of transcendently  $n$ -quasiperiodic fractal sets is given in [LapRaŽu2] and [LapRaŽu1, §3.1]. In [LapRaŽu4] and [LapRaŽu1, Chap. 4], we even construct a set which is transcendently  $\infty$ -quasiperiodic; see §5.

## 5. Maximally hyperfractal $\infty$ -quasiperiodic sets

It is possible to construct a bounded subset  $A$  of the real line, such that the corresponding distance zeta function  $\zeta_A$  has for abscissa of (Lebesgue, i.e., absolute) convergence  $D(\zeta_A)$  any prescribed real number  $D \in (0, 1)$  and  $A$  is *maximally hyperfractal*; that is, any point on the critical line  $\{\operatorname{Re} s = D\}$  is a nonremovable singularity of the corresponding distance zeta function  $\zeta_A$ . In particular, there is no meromorphic continuation of  $\zeta_A$  to any open and connected neighborhood of the critical line (and, moreover, not even to any open and connected neighborhood of an arbitrary point on the critical line). Furthermore, it is possible to construct a maximally hyperfractal set which is  $\infty$ -transcendently quasiperiodic as well. A construction of such sets is described in detail in [LapRaŽu1, Chap. 4] or in [LapRaŽu4]. In the sequel, we provide a rough sketch of this construction.

The set  $A \subset \mathbb{R}$  which is a maximal hyperfractal and  $\infty$ -transcendently quasiperiodic set, can be constructed as the nonincreasing sequence

$$A = A_{\mathcal{L}} = \left\{ a_k := \sum_{j=k}^{\infty} \ell_j : k \in \mathbb{N} \right\} \quad (40)$$

of positive real numbers  $a_k$  converging to zero as  $k \rightarrow \infty$ , generated by a suitable bounded fractal string  $\mathcal{L} = (\ell_j)_{j \geq 1}$ . Roughly speaking, the fractal string  $\mathcal{L}$  is obtained as a (suitably defined) *union of an infinite sequence of bounded fractal strings*  $\mathcal{L}_k := (\ell_{kj})_{j \geq 1}$ , corresponding to generalized Cantor sets of the form  $c_k \cdot C^{(m_k, a_k)}$ , for  $k \in \mathbb{N}$ , with carefully chosen values of the parameters  $m_k$  and  $a_k$  appearing in Definition 4.1, and where  $(c_k)_{k \geq 1}$  is an appropriate summable sequence of positive real numbers.

More precisely, the union  $\mathcal{L} := \bigsqcup_{k=1}^{\infty} \mathcal{L}_k$  of the sequence of bounded fractal strings  $\mathcal{L}_k$  is defined as the set-theoretic union of the elements of the strings, but by definition, each of its elements has for multiplicity the sum of the corresponding multiplicities from all of the fractal strings  $\mathcal{L}_k$  to which belongs the element in question. Note that the multiplicity of an element of  $\mathcal{L}$  is well defined since this

element must belong to at most finitely many bounded fractal strings  $\mathcal{L}_k$ , which follows from the fact that the sequence  $c_k$  converges to 0 as  $k \rightarrow \infty$ . Moreover, we must assume that  $\sum_{k=1}^{\infty} c_k < \infty$ , so that the string  $\mathcal{L}$  be bounded (i.e.,  $\sum_{j=1}^{\infty} \ell_j < \infty$ ). We can also ensure that for each positive integer  $k$ , the corresponding upper box dimension of  $\mathcal{L}_k$  (that is, of the set  $A_{\mathcal{L}_k}$ ) be equal to a fixed value of  $D \in (0, 1)$ , prescribed in advance. (Note that the set  $A_{\mathcal{L}}$  is distinct from  $\cup_{k=1}^{\infty} A_{\mathcal{L}_k}$ .)

Recall that the oscillatory period of  $\mathcal{L}_k$  (in the sense of [Lap-vFr3]), which is defined by  $\mathbf{p}_k := \frac{2\pi}{\log(1/a_k)}$ , provides valuable information about the density of the set of principal complex dimensions of  $\mathcal{L}_k$  on the critical line  $\{\operatorname{Re} s = D\}$ . More precisely, by choosing the coefficient  $a_k \in (0, 1/m_k)$  so that  $a_k \rightarrow 0$  as  $k \rightarrow \infty$ , we see that for the set of principal complex dimensions of the generalized Cantor string  $\mathcal{L}_k$  (i.e., the set of the principal poles of  $\zeta_{\mathcal{L}_k}$ ),  $\dim_{PC} \mathcal{L}_k = \dim_{PC} C^{(m_k, a_k)} = D + \mathbf{p}_k \mathbb{Z}$ , becomes denser and denser on the critical line, as  $k \rightarrow \infty$ , since then the oscillatory period  $\mathbf{p}_k$  tends to zero. Therefore, the distance zeta function of the fractal string  $\mathcal{L} := \sqcup_{k=1}^{\infty} \mathcal{L}_k$  will have  $D + \left(\cup_{k=1}^{\infty} \mathbf{p}_k \mathbb{Z}\right)$  as a set of singularities, which is densely packed on the critical line  $\{\operatorname{Re} s = D\} = D + \mathbb{R}$ , since the set  $\cup_{k=1}^{\infty} \mathbf{p}_k \mathbb{Z}$  is clearly dense in  $\mathbb{R}$ . In conclusion, the whole critical line  $\{\operatorname{Re} s = D\}$  consists of nonremovable singularities of  $\zeta_{\mathcal{L}}$ ,<sup>20</sup> which by definition means that the fractal string  $\mathcal{L}$  is maximally hyperfractal. Hence, the corresponding set  $A := A_{\mathcal{L}}$  is also maximally hyperfractal.

Since the coefficients  $a_k$ , appearing in the definition of the generalized Cantor set (see Definition 4.1), have been chosen above so that  $a_k \rightarrow 0$  as  $k \rightarrow \infty$ , it is clear that  $m_k \rightarrow \infty$ , because  $D = \dim_B C^{(m_k, a_k)} = \frac{m_k}{\log(1/a_k)}$ , where  $D \in (0, 1)$  is given in advance and independent of  $k$ . This enables us to use our result mentioned in Remark 4.6, obtained by means of Baker's theorem from transcendental number theory [Ba], in order to ensure that the sequence of quasiperiods  $T_k := \log(1/a_k)$ ,  $k \in \mathbb{N}$ , is algebraically independent (that is, any finite subset of this set of quasiperiods is linearly independent over the field of algebraic real numbers).<sup>21</sup> According to Definition 4.3, this means that  $\mathcal{L}$  is  $\infty$ -transcendentally quasiperiodic, and so is the corresponding bounded subset  $A := A_{\mathcal{L}}$  of the real line.

As we see from the above rough description, the nature of a subset  $A := A_{\mathcal{L}}$  of the real line which is maximally hyperfractal and  $\infty$ -transcendentally quasiperiodic, is in general extremely complex, although it is, in fact, 'just' defined in terms of a nonincreasing sequence of positive real numbers converging to zero.

<sup>20</sup>In light of the discussion surrounding Eq. (4) above, the same is true if  $\zeta_{\mathcal{L}}$  is replaced by  $\zeta_{A_{\mathcal{L}}}$  or, more generally, by the *relative distance zeta function*  $\zeta_A(\cdot, \Omega)$  defined by  $\zeta_A(s, \Omega) := \int_{\Omega} d(x, A)^{s-N} dx$  (see §6 and [LapRaŽu4] or [LapRaŽu1, Chap. 4]), where  $A = \partial\Omega$  is the boundary of any geometric realization of  $\mathcal{L}$  by a bounded open subset  $\Omega$  of  $\mathbb{R}$ .

<sup>21</sup>The algebraic independence of the set of quasiperiods  $\{T_k : k \geq 1\}$ , with  $k \geq 1$ , can be deduced (using the aforementioned Baker's theorem, [Ba]) if we assume, in addition, that the sequence  $(\mathbf{e}_k)_{k \geq 1}$  (suitably redefined), corresponding to the sequence  $(m_k)_{k \geq 1}$ , is rationally independent.

In closing this discussion, we mention that this construction (as well as Theorem 4.5 and its generalization mentioned in Remark 4.6), extends to any  $N \geq 2$ , by letting  $B := A \times [0, 1]^{N-1} \subset \mathbb{R}^N$ ; see [LapRaŽu1,2,4].

## 6. Fractal zeta functions of relative fractal drums

In this section, we survey some of the definitions and results from [LapRaŽu1, Chap. 4]; see also [LapRaŽu4]. Let  $A$  be a (possibly unbounded) subset of  $\mathbb{R}^N$  and let  $\Omega$  be a (possibly unbounded) Borel subset of  $\mathbb{R}^N$  of finite  $N$ -dimensional Lebesgue measure. We say that the ordered pair  $(A, \Omega)$  is a *relative fractal drum* (or RFD, in short) if there exists a positive real number  $\delta$  such that  $\Omega \subseteq A_\delta$ . It is easy to see that for every  $\delta > 0$ , any bounded subset  $A$  can be identified with the relative fractal drum  $(A, A_\delta)$ . Furthermore, any bounded fractal string  $\mathcal{L} = (\ell_j)_{j=1}^\infty$  can be identified with the relative fractal drum  $(\cup_{j=1}^\infty \partial I_j, \cup_{j=1}^\infty I_j)$ , where  $(I_j)_{j=1}^\infty$  is a family of pairwise disjoint open intervals in  $\mathbb{R}$  such that  $|I_j|_1 = \ell_j$  for all  $j \geq 1$ .

Given a relative fractal drum  $(A, \Omega)$  in  $\mathbb{R}^N$  and for a fixed real number  $r$ , we define the *relative upper* and *relative lower  $r$ -dimensional Minkowski contents* of  $(A, \Omega)$ , respectively, by

$$\mathcal{M}^{*r}(A, \Omega) := \limsup_{t \rightarrow 0^+} \frac{|A_t \cap \Omega|}{t^{N-r}}, \quad \mathcal{M}_*^r(A, \Omega) := \liminf_{t \rightarrow 0^+} \frac{|A_t \cap \Omega|}{t^{N-r}}.$$

The *relative upper* and *relative lower box* (or *Minkowski*) *dimensions* of  $(A, \Omega)$  are then defined, respectively, by

$$\begin{aligned} \overline{\dim}_B(A, \Omega) &:= \inf\{r \in \mathbb{R} : \mathcal{M}^{*r}(A, \Omega) = 0\}, \\ \underline{\dim}_B(A, \Omega) &:= \inf\{r \in \mathbb{R} : \mathcal{M}_*^r(A, \Omega) = 0\}. \end{aligned} \tag{41}$$

It is easy to check that  $-\infty \leq \underline{\dim}_B(A, \Omega) \leq \overline{\dim}_B(A, \Omega) \leq N$ , and it is shown in [LapRaŽu1,4] that the relative box dimensions can indeed attain arbitrary negative values as well, including  $-\infty$  (an obvious example is when  $A_\delta \cap \Omega = \emptyset$  for some  $\delta > 0$ ). Intuitively, negative relative box dimensions correspond to the *property of flatness* of the RFD under consideration. If  $\dim_B(A, \Omega) = -\infty$ , then the RFD  $(A, \Omega)$  is said to be *infinitely flat*. A nontrivial example of an infinitely flat RFD  $(A, \Omega)$  in  $\mathbb{R}^2$  is given by  $A := \{(0, 0)\}$  and  $\Omega := \{(x, y) \in (0, 1)^2 : 0 < y < e^{-1/x}\}$ . Other examples of flat RFDs can be found in [LapRaŽu1,4].

If  $(A, \Omega)$  is such that  $\underline{\dim}_B(A, \Omega) = \overline{\dim}_B(A, \Omega)$ , then this common value is denoted by  $\dim_B(A, \Omega)$  and is called the *box* (or *Minkowski*) *dimension* of  $(A, \Omega)$ . Moreover, if  $(A, \Omega)$  is such that, for some  $D \in (-\infty, N]$ , we have  $0 < \mathcal{M}_*^D(A, \Omega) \leq \mathcal{M}^{*D}(A, \Omega) < \infty$  (in particular, then  $\dim_B(A, \Omega)$  exists and  $D = \dim_B(A, \Omega)$ ), we say that  $(A, \Omega)$  is *Minkowski nondegenerate*. If  $\mathcal{M}_*^D(A, \Omega) = \mathcal{M}^{*D}(A, \Omega)$ , then the common value is denoted by  $\mathcal{M}^D(A, \Omega)$  and called the *Minkowski content* of  $(A, \Omega)$ . Finally, assuming that  $(A, \Omega)$  is such that  $\mathcal{M}^D(A, \Omega)$  exists and  $0 < \mathcal{M}^D(A, \Omega) < \infty$ , we say that the RFD  $(A, \Omega)$  is *Minkowski measurable*.

To any given RFD  $(A, \Omega)$  in  $\mathbb{R}^N$ , we can associate the corresponding *relative distance zeta function* and the *relative tube zeta function* defined, respectively, by

$$\zeta_A(s, \Omega) := \int_{\Omega} d(x, A)^{s-N} dx, \quad \tilde{\zeta}_A(s, \Omega) := \int_0^{\delta} t^{s-N-1} |A_t \cap \Omega| dt, \quad (42)$$

for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large, where  $\delta$  is a fixed positive real number. They are a valuable theoretical and technical new tool in the study of fractals.

The basic result dealing with relative distance zeta functions, analogous to Theorem 2.4 of §2, is provided by the following theorem.

**Theorem 6.1** ([LapRaŽu1,4]). *Let  $(A, \Omega)$  be an arbitrary RFD. Then:*

(a) *The distance zeta function  $\zeta_A(\cdot, \Omega)$  is holomorphic on  $\{\operatorname{Re} s > \overline{\dim}_B(A, \Omega)\}$ . Moreover,  $\Pi(\zeta_A(\cdot, \Omega)) = \{\operatorname{Re} s > \overline{\dim}_B(A, \Omega)\}$ ; that is,*

$$D(\zeta_A(\cdot, \Omega)) = \overline{\dim}_B(A, \Omega). \quad (43)$$

(b) *If the box (or Minkowski) dimension  $D := \dim_B(A, \Omega)$  exists,  $D < N$ , and  $\mathcal{M}_*^D(A, \Omega) > 0$ , then  $\zeta_A(s, \Omega) \rightarrow +\infty$  as  $s \in \mathbb{R}$  converges to  $D$  from the right. In particular,  $\mathcal{H}(\zeta_A(\cdot, \Omega)) = \Pi(\zeta_A(\cdot, \Omega)) = \{\operatorname{Re} s > \dim_B(A, \Omega)\}$ ; that is,*

$$D_{\text{hol}}(\zeta_A(\cdot, \Omega)) = D(\zeta_A(\cdot, \Omega)) = \dim_B(A, \Omega). \quad (44)$$

An entirely analogous result holds for the tube zeta function  $\tilde{\zeta}_A(\cdot, \Omega)$ , except for the fact that the hypothesis  $D < N$  is no longer needed in the counterpart of part (b) of Theorem 6.1.

A very useful property of relative distance zeta functions is the following *scaling property*: for any RFD  $(A, \Omega)$  and for any positive real number  $\lambda$ , we have

$$\zeta_{\lambda A}(s, \lambda\Omega) = \lambda^s \zeta_A(s, \Omega). \quad (45)$$

We refer the interested reader to [LapRaŽu1, Chap. 4] and [LapRaŽu4–7] for many other related results, examples and comments. We mention, in particular, that ‘fractal drums’ (that is, ‘drums with fractal boundary’, in the sense of [Lap1–3], for example)<sup>22</sup> correspond to RFDs of the form  $(\partial\Omega, \Omega)$ , where  $\Omega$  is a nonempty bounded open subset of  $\mathbb{R}^N$ , and that the results discussed in §5 above are applied in a crucial way in order to show the optimality of certain inequalities pertaining to the meromorphic continuations of the spectral zeta functions of fractal drums (viewed as RFDs); see [LapRaŽu1, §4.3] and [LapRaŽu7].

## 7. Fractal tube formulas and a Minkowski measurability criterion

In this section, we briefly explain how under suitable growth conditions on the relative distance (or tube) zeta function (see a variant of the languidity (resp., of the strong languidity) condition of [Lap-vFr3, §5.3] given in [LapRaŽu5,6]), it is possible to recover a pointwise or distributional fractal tube formula for a relative fractal drum  $(A, \Omega)$  in  $\mathbb{R}^N$ , expressed as a sum of residues over its visible complex

<sup>22</sup>See also [Lap-vFr3, §12.5], [Lap6] and [LapRaŽu7] for many other references on fractal drums.

dimensions. These fractal tube formulas, along with a Tauberian theorem due to Wiener and Pitt (which generalizes Ikehara’s Tauberian theorem, see [Kor, Pos]) make it possible to derive a Minkowski measurability criterion for a large class of relative fractal drums (and compact subsets) of  $\mathbb{R}^N$ . These results generalize to higher dimensions the corresponding ones obtained for fractal strings (that is, when  $N = 1$ ) in [Lap-vFr3, §8.1 and §8.3].

The results of this section are announced in [LapRaŽu5] and fully proved in [LapRaŽu6]. (See also [LapRaŽu1, Chap. 5].) Furthermore, we refer the interested reader to [LapRaŽu1,6] and [Lap-vFr3, §8.2 and §13.1] for additional references on tube formulas in various settings, including [DeKÖÜ, Fed1, Gra, HuLaWei, Schn, Zäh, Wey, LapPe1–2, LapPeWi1–2, LapLu, LapLu-vFr1–2]. (See also [Lap-vFr3, §13.1, §13.2 and §13.4].)

In order to be able to state the fractal tube formulas, we introduce the following notions, adapted from [Lap-vFr3] to the present much more general context. The *screen*  $S$  is the graph of a bounded, real-valued, Lipschitz continuous function  $S(\tau)$ , with the horizontal and vertical axes interchanged:  $S := \{S(\tau) + \tau : \tau \in \mathbb{R}\}$  and we let  $\sup S := \sup_{\tau \in \mathbb{R}} S(\tau) \in \mathbb{R}$ . Given a relative fractal drum  $(A, \Omega)$  of  $\mathbb{R}^N$ , we always assume that the screen  $S$  lies to the left of the critical line  $\{\operatorname{Re} s = \overline{\dim}_B(A, \Omega)\}$ , i.e., that  $\sup S \leq \overline{\dim}_B(A, \Omega)$ . Furthermore, the *window*  $W$  is defined as  $W := \{s \in \mathbb{C} : \operatorname{Re} s \geq S(\operatorname{Im} s)\}$ . The relative fractal drum  $(A, \Omega)$  is said to be *admissible* if its tube (or distance) zeta function can be meromorphically extended to an open connected neighborhood of some window  $W$ .

Assume now that  $(A, \Omega)$  is an admissible relative fractal drum of  $\mathbb{R}^N$  for some screen  $S$  such that its distance zeta function satisfies appropriate growth conditions (see [LapRaŽu5,6] for details).<sup>23</sup> Then its relative tube function satisfies the following identity, for all positive real numbers  $t$  sufficiently small:<sup>24</sup>

$$|A_t \cap \Omega| = \sum_{\omega \in \mathcal{P}(\zeta_A(\cdot, \Omega), W)} \operatorname{res} \left( \frac{t^{N-s}}{N-s} \zeta_A(s, \Omega), \omega \right) + \mathcal{R}(t). \quad (46)$$

The above *fractal tube formula* is interpreted pointwise or distributionally, depending on the growth properties of  $\zeta_A(\cdot, \Omega)$  and then,  $\mathcal{R}(t)$  is a pointwise or distributional<sup>25</sup> asymptotic error term of order at most  $O(t^{N-\sup S})$  as  $t \rightarrow 0^+$ . Moreover, if  $S$  lies strictly to the left of the vertical line  $\{\operatorname{Re} s = D\}$  (that is, if  $\sup S < D$ ), then  $\mathcal{R}(t)$  is  $o(t^{N-\sup S})$ , pointwise or distributionally, as  $t \rightarrow 0^+$ . In the case when  $\zeta_A(\cdot, \Omega)$  satisfies stronger growth assumptions (i.e., the analog of the “strong languidity condition” of [Lap-vFr3, Def. 5.3]), we obtain a tube formula without an error term (i.e.,  $\mathcal{R}(t) \equiv 0$ ) and with  $W = \mathbb{C}$ . Following [Lap-vFr3], the resulting formula is then called an *exact* fractal tube formula.

<sup>23</sup>Roughly speaking,  $\zeta_{(A, \Omega)} := \zeta_A(\cdot, \Omega)$  is assumed to grow at most polynomially along the vertical direction of the screen and along suitable horizontal directions (avoiding the poles of  $\zeta_{(A, \Omega)}$ ); see [Lap-vFr3, Def. 5.2] for the so-called “languidity condition”.

<sup>24</sup>The ranges within which the formulas are valid are fully specified in [LapRaŽu5,6].

<sup>25</sup>For the precise definition of distributional asymptotics, see [Lap-vFr3, §5.4.2], [LapRaŽu5–6] and the relevant references therein.

The tube formula (46) can also be expressed in terms of the relative tube zeta function when analogous growth conditions are imposed on  $\tilde{\zeta}_A(\cdot, \Omega)$ :<sup>26</sup>

$$|A_t \cap \Omega| = \sum_{\omega \in \mathcal{P}(\tilde{\zeta}_A(\cdot, \Omega), W)} \operatorname{res} \left( t^{N-s} \tilde{\zeta}_A(s, \Omega), \omega \right) + \mathcal{R}(t). \quad (47)$$

In fact, the key observation for deriving the above formula is the fact that

$$\tilde{\zeta}_A(s, \Omega) = \int_0^{+\infty} t^{s-N-1} \chi_{(0,\delta)}(t) |A_t \cap \Omega| dt = \{\mathfrak{M}f\}(s), \quad (48)$$

where  $\chi_E$  is the characteristic function of the set  $E$ ,  $\{\mathfrak{M}\psi\}(s) := \int_0^{+\infty} t^{s-1} \psi(t) dt$  is the Mellin transform of the function  $\psi$ , and  $f(t) := t^{-N} \chi_{(0,\delta)}(t) |A_t \cap \Omega|$ . One then applies the inverse Mellin transform (see [Titch]) to recover the relative tube function  $t \mapsto |A_t \cap \Omega|$  and proceeds in a similar manner as in [Lap-vFr3, Chap. 5] for the case of fractal strings.

As an application, the following result generalizes the Minkowski measurability criterion given in [Lap-vFr3, Thm. 8.15] for fractal strings to the present case of relative fractal drums.

**Theorem 7.1 (Minkowski measurability criterion, [LapRaŽu5,6]).** *Let  $(A, \Omega)$  be an admissible relative fractal drum of  $\mathbb{R}^N$  such that  $D := \dim_B A$  exists and  $D < N$ . Furthermore, assume that its relative distance (or tube) zeta function satisfies appropriate growth conditions<sup>27</sup> for a screen passing between the critical line  $\{\operatorname{Re} s = D\}$  and all the complex dimensions of  $A$  with real part strictly less than  $D$ . Then, the following statements are equivalent:*

(a)  *$A$  is Minkowski measurable.*

(b)  *$D$  is the only pole of the distance zeta function  $\zeta_A$  located on the critical line  $\{\operatorname{Re} s = D\}$ , and it is simple.*

There exist relative fractal drums which do not satisfy the hypothesis of Theorem 7.1 concerning the screen; see [Lap-vFr3, Exple. 5.32]. We point out that the fractal tube formula (46) can be used to recover (or obtain for the first time) the (relative) fractal tube formulas for a variety of well-known (and not necessarily self-similar) fractal sets, as is illustrated by the following examples.

*Example 7.2.* Recall from Proposition 3.5 that the distance zeta function of the Sierpiński carpet  $A$  is given for all  $s \in \mathbb{C}$  by

$$\zeta_A(s) = \frac{8}{2^s s (s-1) (3^s - 8)} + 2\pi \frac{\delta^s}{s} + 4 \frac{\delta^{s-1}}{s-1},$$

<sup>26</sup>Note that in light of the functional equation (13), assuming growth conditions for  $\zeta_A$  is essentially equivalent to assuming them for  $\tilde{\zeta}_A$  (and vice versa).

<sup>27</sup>See [LapRaŽu5,6] for details about these growth conditions.

for  $\delta > 1/6$ , and is meromorphic on all of  $\mathbb{C}$ . It is easy to check that  $\zeta_A$  satisfies growth conditions which are good enough for (46) to hold pointwise without an error term and for all  $t \in (0, 1/2)$ :

$$|A_t| = \sum_{\omega \in \mathcal{P}(\zeta_A, \mathbb{C})} \operatorname{res} \left( \frac{t^{2-s}}{2-s} \zeta_A(s), \omega \right). \quad (49)$$

Now, also recall from Proposition 3.5 that  $\mathcal{P}(\zeta_A, \mathbb{C}) = \{0, 1\} \cup \{s_k : k \in \mathbb{Z}\}$ , where  $s_k = \log_3 8 + \frac{2\pi}{\log 3} k$  for all  $k \in \mathbb{Z}$ . Furthermore,  $\operatorname{res}(\zeta_A, 0) = 2\pi + 8/7$ ,  $\operatorname{res}(\zeta_A, 1) = 16/5$  and the residues at  $s_k$  are given in (31); so that (49) becomes the following exact, pointwise fractal tube formula, valid for all  $t \in (0, 1/2)$ :

$$|A_t| = \frac{t^{2-\log_3 8}}{\log 3} \sum_{k=-\infty}^{+\infty} \frac{2^{-s_k} t^{-\frac{2\pi}{\log 3} k}}{s_k(s_k-1)(2-s_k)} + \frac{16}{5} t + \left(2\pi + \frac{8}{7}\right) t^2. \quad (50)$$

The above example can be generalized to an  $N$ -dimensional analog of the Sierpiński carpet (see [LapRaŽu1,6]). We next establish the special case of this assertion for the relative 3-dimensional Sierpiński carpet.

*Example 7.3.* Let  $A$  be the three-dimensional analog of the Sierpiński carpet and  $\Omega$  the closed unit cube in  $\mathbb{R}^3$ . More precisely, we construct  $A$  by dividing  $\Omega$  into 27 congruent cubes and remove the open middle cube, then we iterate this step with each of the 26 remaining smaller closed cubes; and so on, ad infinitum. By choosing  $\delta > 1/6$ , we deduce that  $\zeta_A$  is meromorphic on  $\mathbb{C}$  and given for all  $s \in \mathbb{C}$  by (see [LapRaŽu1,6])

$$\zeta_A(s, \Omega) = \frac{48 \cdot 2^{-s}}{s(s-1)(s-2)(3^s-26)}. \quad (51)$$

In particular,  $\mathcal{P}(\zeta_A(\cdot, \Omega), \mathbb{C}) = \{0, 1, 2\} \cup (\log_3 26 + \mathbf{p} \mathbb{Z})$ , where  $\mathbf{p} := 2\pi / \log 3$ . Furthermore, we have that

$$\operatorname{res}(\zeta_A(\cdot, \Omega), 0) = -\frac{24}{25}, \quad \operatorname{res}(\zeta_A(\cdot, \Omega), 1) = \frac{24}{23}, \quad \operatorname{res}(\zeta_A(\cdot, \Omega), 2) = -\frac{6}{17}$$

and, by letting  $\omega_k := \log_3 26 + \mathbf{p}k$  for all  $k \in \mathbb{Z}$ ,

$$\operatorname{res}(\zeta_A(\cdot, \Omega), \omega_k) = \frac{24 \cdot 2^{-\omega_k}}{13 \cdot \omega_k(\omega_k-1)(\omega_k-2) \log 3}.$$

Again, the relative distance zeta function  $\zeta_A(\cdot, \Omega)$  satisfies sufficiently good growth conditions, which enables us to obtain the following exact pointwise relative tube formula, valid for all  $t \in (0, 1/2)$ :

$$|A_t \cap \Omega| = \frac{24 t^{3-\log_3 26}}{13 \log 3} \sum_{k=-\infty}^{+\infty} \frac{2^{-\omega_k} t^{-\mathbf{p}k}}{(3-\omega_k)(\omega_k-1)(\omega_k-2)\omega_k} - \frac{6}{17} t + \frac{12}{23} t^2 - \frac{8}{25} t^3.$$

In particular, we conclude that  $\dim_B(A, \Omega) = \log_3 26$  and, by Theorem 7.1, that, as expected,  $(A, \Omega)$  is not Minkowski measurable.

One can similarly recover the well-known fractal tube formula for the Sierpiński gasket obtained in [LapPe2] (and also, more recently, by a somewhat different method, in [DeKÖÜ]), as well as a tube formula for its  $N$ -dimensional analog described in [LapRaŽu1, Chap. 5].<sup>28</sup> We also point out that, in light of the functional equation (4), the above fractal tube formulas (46) and (47) generalize the corresponding ones obtained for fractal strings (i.e., when  $N = 1$ ) in [Lap-vFr3, §8.1]. Furthermore, these tube formulas can also be applied to a variety of fractal sets that are not self-similar, including ‘fractal nests’ and ‘geometric chirps’ (see [LapRaŽu1, Chaps. 3 and 4] for the definitions of these notions and [LapRaŽu6] along with [LapRaŽu1, Chap. 5] for the actual fractal tube formulas).

We conclude this section by briefly explaining how these results can also be applied in order to recover (and extend) the tube formulas for *self-similar sprays* generated by a suitable bounded open set  $G \subset \mathbb{R}^N$ . (See [LapPe2, LapPeWi1].) A self-similar spray is a collection  $(G_k)_{k \in \mathbb{N}}$  of pairwise disjoint sets  $G_k \subset \mathbb{R}^N$ , with  $G_0 := G$  and such that  $G_k$  is a scaled copy of  $G$  by some factor  $\lambda_k > 0$ . The sequence  $(\lambda_k)_{k \in \mathbb{N}}$  is called the *scaling sequence* associated with the spray and is obtained from a ‘ratio list’  $\{r_1, r_2, \dots, r_J\}$ , with  $0 < r_j < 1$  for each  $j \in \{1, 2, \dots, J\}$ , by building all possible words based on the ratios  $r_j$ . Let now  $(A, \Omega)$  be the relative fractal drum such that  $A := \partial(\cup_{k \in \mathbb{N}} G_k)$  and  $\Omega := \cup_{k \in \mathbb{N}} G_k$ , with  $\dim_B(\partial G, G) < N$ . Then, it is clear that its relative distance zeta function  $\zeta_A(\cdot, \Omega)$  satisfies the following functional equation, for all  $s \in \mathbb{C}$  with  $\operatorname{Re} s$  sufficiently large:

$$\zeta_A(s, \Omega) = \zeta_{\partial G}(s, G) + \zeta_{r_1 A}(s, r_1 \Omega) + \dots + \zeta_{r_J A}(s, r_J \Omega), \quad (52)$$

where  $(r_j A, r_j \Omega)$  denotes the relative fractal drum  $(A, \Omega)$  scaled by the factor  $r_j$ . Furthermore, by using the scaling property (45) of the relative distance zeta function, the above equation becomes

$$\zeta_A(s, \Omega) = \zeta_{\partial G}(s, G) + r_1^s \zeta_A(s, \Omega) + \dots + r_J^s \zeta_A(s, \Omega), \quad (53)$$

which yields that

$$\zeta_A(s, \Omega) = \frac{\zeta_{\partial G}(s, G)}{1 - \sum_{j=1}^J r_j^s}. \quad (54)$$

It is now enough to assume that the relative distance zeta function  $\zeta_{\partial G}(s, G)$  of the generating relative fractal drum  $(\partial G, G)$  satisfies suitable growth conditions in order to obtain the following formula for the ‘inner’ volume of  $A_t = (\partial \Omega)_t$  relative to  $\Omega := \cup_{k \in \mathbb{N}} G_k$ , for all positive  $t$  sufficiently small:

$$|A_t \cap \Omega| = \sum_{\omega \in \mathfrak{D}(W) \cup \mathcal{P}(\zeta_{\partial G}(\cdot, G), W)} \operatorname{res} \left( \frac{t^{N-s} \zeta_{\partial G}(s, G)}{(N-s) \left(1 - \sum_{j=1}^J r_j^s\right)}, \omega \right) + \mathcal{R}(t), \quad (55)$$

where  $\mathfrak{D}(W)$  denotes the set of all visible complex solutions of  $\sum_{j=1}^J r_j^s = 1$  (in  $W$ ) and  $W$  is the window defined earlier. It is easy to check that (at least) in the case of

<sup>28</sup>We can also recover and extend the significantly more general fractal tube formulas obtained (for fractal sprays and self-similar tilings) in [LapPeWi1] and used, in particular, in [LapPeWi2].

monophase or pluriphase generators (in the sense of [LapPe2] and [LapPeWi1,2]), these growth conditions are satisfied, so that one obtains exactly the same distributional or pointwise fractal tube formulas as in [LapPe2] or [LapPeWi1], respectively, after having calculated the distance zeta function  $\zeta_{\partial G}(\cdot, G)$  of the generator. Moreover, if  $\zeta_{\partial G}(\cdot, G)$  is strongly languid, we can let  $\mathcal{R}(t) \equiv 0$  and  $W = \mathbb{C}$  in (55) and therefore obtain *exact* fractal tube formulas.

We conclude this survey by pointing out that a broad variety of open problems and suggestions for directions for future research in this area are proposed in [LapRaŽu1, Chap. 6].

## References

- [Ba] A. Baker, *Transcendental Number Theory*, Cambridge Univ. Press, Cambridge, 1975.
- [BesTay] A. S. Besicovitch and S. J. Taylor, On the complementary intervals of a linear closed set of zero Lebesgue measure, *J. London Math. Soc.* **29** (1954), 449–459.
- [Bou] G. Bouligand, Ensembles impropres et nombre dimensionnel, *Bull. Sci. Math.* (2) **52** (1928), 320–344 and 361–376.
- [DeKÖÜ] A. Deniz, Ş. Koçak, Y. Özdemir and A. E. Üreyen, Tube volumes via functional equations, *J. Geom.* **105** (2014), 1–10.
- [ElLapMacRo] K. E. Ellis, M. L. Lapidus, M. C. Mackenzie and J. A. Rock, Partition zeta functions, multifractal spectra, and tapestries of complex dimensions, in: *Benoît Mandelbrot: A Life in Many Dimensions* (M. Frame and N. Cohen, eds.), The Mandelbrot Memorial Volume, World Scientific, Singapore, 2015, pp. 267–322. (Also: e-print, [arXiv:1007.1467v2](https://arxiv.org/abs/1007.1467v2) [math-ph], 2011; IHES preprint, IHES/M/12/15, 2012.)
- [Fal1] K. J. Falconer, On the Minkowski measurability of fractals, *Proc. Amer. Math. Soc.* **123** (1995), 1115–1124.
- [Fal2] K. J. Falconer, *Fractal Geometry: Mathematical Foundations and Applications*, third edition, John Wiley and Sons, Chichester, 2014.
- [Fed1] H. Federer, Curvature measures, *Trans. Amer. Math. Soc.* **93** (1959), 418–491.
- [Fed2] H. Federer, *Geometric Measure Theory*, Springer-Verlag, New York, 1969.
- [Gel] A. O. Gel'fond, *Transcendental and Algebraic Numbers*, Dover Phoenix editions, Dover Publications, New York, 1960.
- [Gra] A. Gray, *Tubes*, second edition, Progress in Math., vol. 221, Birkhäuser, Boston, 2004.
- [HamLap] B. M. Hambly and M. L. Lapidus, Random fractal strings: their zeta functions, complex dimensions and spectral asymptotics, *Trans. Amer. Math. Soc.* No. 1, **358** (2006), 285–314.
- [HarPol] R. Harvey and J. Polking, Removable singularities of solutions of linear partial differential equations, *Acta Math.* **125** (1970), 39–56.

- [HeLap] C. Q. He and M. L. Lapidus, *Generalized Minkowski Content, Spectrum of Fractal Drums, Fractal Strings and the Riemann Zeta-Function*, *Memoirs Amer. Math. Soc.* No. 608, **127** (1997), 1–97.
- [HerLap1] H. Herichi and M. L. Lapidus, *Quantized Number Theory, Fractal Strings and the Riemann Hypothesis: From Spectral Operators to Phase Transitions and Universality*, research monograph, World Scientific Publishers, Singapore, 2016, to appear, approx. 220 pages.
- [HerLap2] H. Herichi and M. L. Lapidus, Fractal complex dimensions, Riemann hypothesis and invertibility of the spectral operator, in: *Fractal Geometry and Dynamical Systems in Pure and Applied Mathematics I: Fractals in Pure Mathematics* (D. Carfi, M. L. Lapidus, E. P. J. Pearse and M. van Frankenhuijsen, eds.), Contemporary Mathematics, vol. **600**, Amer. Math. Soc., Providence, R. I., 2013, pp. 51–89. (Also: e-print, [arXiv:1210.0882v3](https://arxiv.org/abs/1210.0882v3) [math.FA], 2013; IHES preprint, IHES/M/12/25, 2012.)
- [HorŽu] L. Horvat and D. Žubrinić, Maximally singular Sobolev functions, *J. Math. Anal. Appl.* No. 2, **304** (2005), 531–541.
- [HuLaWei] D. Hug, G. Last and W. Weil, A local Steiner-type formula for general closed sets and applications, *Math. Z.* **246** (2004), 237–272.
- [Kor] J. Korevaar, *Tauberian Theory: A Century of Developments*, Springer-Verlag, Heidelberg, 2004.
- [LalLap1] N. Lal and M. L. Lapidus, Hyperfunctions and spectral zeta functions of Laplacians on self-similar fractals, *J. Phys. A: Math. Theor.* **45** (2012), 365205, 14 pp. (Also: e-print, [arXiv:1202.4126v2](https://arxiv.org/abs/1202.4126v2) [math-ph], 2012; IHES preprint, IHES/M/12/14, 2012.)
- [LalLap2] N. Lal and M. L. Lapidus, The decimation method for Laplacians on fractals: Spectra and complex dynamics, in: *Fractal Geometry and Dynamical Systems in Pure and Applied Mathematics II: Fractals in Applied Mathematics* (D. Carfi, M. L. Lapidus, E. P. J. Pearse and M. van Frankenhuijsen, eds.), Contemporary Mathematics, vol. **601**, Amer. Math. Soc., Providence, R. I., 2013, pp. 227–249. (Also: e-print, [arXiv:1302.4007v2](https://arxiv.org/abs/1302.4007v2) [math-ph], 2014; IHES preprint, IHES/M/12/31, 2012.)
- [Lap1] M. L. Lapidus, Fractal drum, inverse spectral problems for elliptic operators and a partial resolution of the Weyl–Berry conjecture, *Trans. Amer. Math. Soc.* **325** (1991), 465–529.
- [Lap2] M. L. Lapidus, Spectral and fractal geometry: From the Weyl–Berry conjecture for the vibrations of fractal drums to the Riemann zeta-function, in: *Differential Equations and Mathematical Physics* (C. Bennewitz, ed.), Proc. Fourth UAB Internat. Conf. (Birmingham, March 1990), Academic Press, New York, 1992, pp. 151–182.
- [Lap3] M. L. Lapidus, Vibrations of fractal drums, the Riemann hypothesis, waves in fractal media, and the Weyl–Berry conjecture, in: *Ordinary and Partial Differential Equations* (B. D. Sleeman and R. J. Jarvis, eds.), vol. IV, Proc. Twelfth Internat. Conf. (Dundee, Scotland, UK, June 1992), Pitman Research Notes in Math. Series, vol. **289**, Longman Scientific and Technical, London, 1993, pp. 126–209.

- [Lap4] M. L. Lapidus, *In Search of the Riemann Zeros: Strings, Fractal Membranes and Noncommutative Spacetimes*, Amer. Math. Soc., Providence, R. I., 2008.
- [Lap5] M. L. Lapidus, Towards quantized number theory: Spectral operators and an asymmetric criterion for the Riemann hypothesis, preprint, 2014. (For inclusion in the special issue of the *Philosophical Transactions of the Royal Society*, Ser. A, titled “*Geometric Concepts in the Foundations of Physics*”, 2015.) (Also: e-print, [arXiv:1501.05362v1 \[math-ph\]](#), 2015.)
- [Lap6] M. L. Lapidus, The sound of fractal strings and the Riemann hypothesis, in: *Analytic Number Theory: In Honor of Helmut Maier’s 60th Birthday* (C. M. Pomerance and T. Rassias, eds.), Springer, Berlin, Heidelberg and New York, in press, 2015. (Also: e-print, [arXiv:1505.01548v1 \[math-ph\]](#), 2015.)
- [LapLéRo] M. L. Lapidus, J. Lévy-Véhel and J. A. Rock, Fractal strings and multifractal zeta functions, *Lett. Math. Phys.* No. 1, **88** (2009), 101–129 (special issue dedicated to the memory of Moshe Flato). (Springer Open Access: DOI 10.1007/s1105-009-0302-y.) (Also: e-print, [arXiv:math-ph/0610015v3](#), 2009.)
- [LapLu] M. L. Lapidus and H. Lu, The geometry of  $p$ -adic fractal strings: A comparative survey, in: *Advances in Non-Archimedean Analysis*, Proc. 11th Internat. Conf. on “*p-Adic Functional Analysis*” (Clermont-Ferrand, France, July 2010), (J. Araujo, B. Diarra and A. Escassut, eds.), Contemporary Mathematics, vol. **551**, Amer. Math. Soc., Providence, R. I., 2011, pp. 163–206. (Also: e-print, [arXiv:1105.2966v1 \[math.MG\]](#), 2011.)
- [LapLu-vFr1] M. L. Lapidus, H. Lu and M. van Frankenhuysen, Minkowski measurability and exact fractal tube formulas for  $p$ -adic self-similar strings, in: *Fractal Geometry and Dynamical Systems in Pure and Applied Mathematics I: Fractals in Pure Mathematics* (D. Carfi, M. L. Lapidus, E. P. J. Pearse and M. van Frankenhuysen, eds.), Contemporary Mathematics, vol. **600**, Amer. Math. Soc., Providence, R. I., 2013, pp. 161–184. (Also: e-print, [arXiv:1209.6440v1 \[math.MG\]](#), 2012; IHES preprint; IHES/M/12/23, 2013.)
- [LapLu-vFr2] M. L. Lapidus, H. Lu and M. van Frankenhuysen, Minkowski dimension and explicit tube formulas for  $p$ -adic fractal strings, preprint, 2015.
- [LapMa1] M. L. Lapidus and H. Maier, Hypothèse de Riemann, cordes fractales vibrantes et conjecture de Weyl–Berry modifiée, *C. R. Acad. Sci. Paris Sér. I Math.* **313** (1991), 19–24.
- [LapMa2] M. L. Lapidus and H. Maier, The Riemann hypothesis and inverse spectral problems for fractal strings, *J. London Math. Soc.* (2) **52** (1995), 15–34.
- [LapPe1] M. L. Lapidus and E. P. J. Pearse, A tube formula for the Koch snowflake curve, with applications to complex dimensions, *J. London Math. Soc.* No. 2, **74** (2006), 397–414. (Also: e-print, [arXiv:math-ph/0412029v2](#), 2005.)
- [LapPe2] M. L. Lapidus and E. P. J. Pearse, Tube formulas and complex dimensions of self-similar tilings, *Acta Applicandae Mathematicae* No. 1, **112** (2010), 91–137. (Springer Open Access: DOI 10.1007/S10440-010-9562-x.) (Also: e-print, [arXiv:math.DS/0605527v5](#), 2010; IHES preprint, IHES/M/08/27, 2008.)

- [LapPeWi1] M. L. Lapidus, E. P. J. Pearse and S. Winter, Pointwise tube formulas for fractal sprays and self-similar tilings with arbitrary generators, *Adv. in Math.* **227** (2011), 1349–1398. (Also: e-print, [arXiv:1006.3807v3 \[math.MG\]](#), 2011.)
- [LapPeWi2] M. L. Lapidus, E. P. J. Pearse and S. Winter, Minkowski measurability results for self-similar tilings and fractals with monophasic generators, in: *Fractal Geometry and Dynamical Systems in Pure and Applied Mathematics I: Fractals in Pure Mathematics* (D. Carfi, M. L. Lapidus, E. P. J. Pearse and M. van Frankenhuysen, eds.), *Contemporary Mathematics*, vol. **600**, Amer. Math. Soc., Providence, R. I., 2013, pp. 185–203. (Also: e-print, [arXiv:1104.1641v3 \[math.MG\]](#), 2012; IHES preprint, IHES/M/12/33, 2012.)
- [LapPo1] M. L. Lapidus and C. Pomerance, Fonction zêta de Riemann et conjecture de Weyl–Berry pour les tambours fractals, *C. R. Acad. Sci. Paris Sér. I Math.* **310** (1990), 343–348.
- [LapPo2] M. L. Lapidus and C. Pomerance, The Riemann zeta-function and the one-dimensional Weyl–Berry conjecture for fractal drums, *Proc. London Math. Soc.* (3) **66** (1993), No. 1, 41–69.
- [LapPo3] M. L. Lapidus and C. Pomerance, Counterexamples to the modified Weyl–Berry conjecture on fractal drums, *Math. Proc. Cambridge Philos. Soc.* **119** (1996), 167–178.
- [LapRaŽu1] M. L. Lapidus, G. Radunović and D. Žubrinić, *Fractal Zeta Functions and Fractal Drums: Higher-Dimensional Theory of Complex Dimensions*, research monograph, Springer, New York, 2016, to appear, approx. 560 pages.
- [LapRaŽu2] M. L. Lapidus, G. Radunović and D. Žubrinić, Distance and tube zeta functions of fractals and arbitrary compact sets, preprint, 2015. (Also: e-print [arXiv: 1506.3525 \[math-ph\]](#), 2015.)
- [LapRaŽu3] M. L. Lapidus, G. Radunović and D. Žubrinić, Complex dimensions of fractals and meromorphic extensions of fractal zeta functions, preprint, 2015.
- [LapRaŽu4] M. L. Lapidus, G. Radunović and D. Žubrinić, Zeta functions and complex dimensions of relative fractal drums: Theory, examples and applications, in preparation, 2015.
- [LapRaŽu5] M. L. Lapidus, G. Radunović and D. Žubrinić, Fractal tube formulas and a Minkowski measurability criterion for compact subsets of Euclidean spaces, preprint, 2014. (Also: e-print, [arXiv:1411.5733.v2 \[math-ph\]](#), 2014.)
- [LapRaŽu6] M. L. Lapidus, G. Radunović and D. Žubrinić, Fractal tube formulas for compact sets and relative fractal drums, with application to a Minkowski measurability criterion, preprint, 2015.
- [LapRaŽu7] M. L. Lapidus, G. Radunović and D. Žubrinić, Fractal zeta functions and complex dimensions of relative fractal drums, *J. Fixed Point Theory and Appl.* No. 2, **15** (2014), 321–378. Festschrift issue in honor of Haim Brezis’ 70th birthday. (DOI 10.1007/s11784-014-0207-y.) (Also: e-print, [arXiv:1407.8094v3 \[math-ph\]](#), 2014.)

- [LapRo] M. L. Lapidus and J. A. Rock, Towards zeta functions and complex dimensions of multifractals, *Complex Variables and Elliptic Equations* No. 6, **54** (2009), 545–560. (Also: e-print, [arXiv:math-ph/0810.0789](https://arxiv.org/abs/math-ph/0810.0789), 2008.)
- [LapRoŽu] M. L. Lapidus, J. A. Rock and D. Žubričić, Box-counting fractal strings, zeta functions, and equivalent forms of Minkowski dimension, in: *Fractal Geometry and Dynamical Systems in Pure and Applied Mathematics I: Fractals in Pure Mathematics* (D. Carfi, M. L. Lapidus, E. P. J. Pearse and M. van Frankenhuijsen, eds.), Contemporary Mathematics, vol. **600**, Amer. Math. Soc., Providence, R. I., 2013, pp. 239–271. (Also: e-print, [arXiv:1207.6681v2 \[math-ph\]](https://arxiv.org/abs/1207.6681v2), 2013; IHES preprint, IHES/M/12/22, 2012.)
- [Lap-vFr1] M. L. Lapidus and M. van Frankenhuijsen, *Fractal Geometry and Number Theory: Complex Dimensions of Fractal Strings and Zeros of Zeta Functions*, Birkhäuser, Boston, 2000.
- [Lap-vFr2] M. L. Lapidus and M. van Frankenhuijsen, *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and Spectra of Fractal Strings*, Springer Monographs in Mathematics, Springer, New York, 2006.
- [Lap-vFr3] M. L. Lapidus and M. van Frankenhuijsen, *Fractal Geometry, Complex Dimensions and Zeta Functions: Geometry and Spectra of Fractal Strings*, second revised and enlarged edition (of the 2006 edn., [Lap-vFr2]), Springer Monographs in Mathematics, Springer, New York, 2013.
- [LéMen] J. Lévy-Véhel and F. Mendivil, Multifractal and higher-dimensional zeta functions, *Nonlinearity* No. 1, **24** (2011), 259–276.
- [Mat] P. Mattila, *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*, Cambridge Univ. Press, Cambridge, 1995.
- [Pos] A. G. Postnikov, *Tauberian Theory and its Applications*, Proc. Steklov Institute of Mathematics, vol. **144**, 1979 (English transl., issue 2, 1980), Amer. Math. Soc., Providence, R.I., 1980.
- [RaWi] J. Rataj and S. Winter, Characterization of Minkowski measurability in terms of surface area, *J. Math. Anal. Appl.* **400** (2013), 120–132. (Also: e-print, [arXiv:1111.1825v2 \[math.CA\]](https://arxiv.org/abs/1111.1825v2), 2012.)
- [Schn] R. Schneider, *Convex Bodies: The Brunn–Minkowski Theory*, Cambridge Univ. Press, Cambridge, 2003.
- [Ser] J.-P. Serre, *A Course in Arithmetic*, English transl., Springer-Verlag, Berlin, 1973.
- [Sta] L. L. Stachó, On the volume function of parallel sets, *Acta Sci. Math.* **38** (1976), 365–374.
- [Tep1] A. Teplyaev, Spectral zeta functions of symmetric fractals, in: *Progress in Probability*, vol. **57**, Birkhäuser-Verlag, Basel, 2004, pp. 245–262.
- [Tep2] A. Teplyaev, Spectral zeta functions of fractals and the complex dynamics of polynomials, *Trans. Amer. Math. Soc.* **359** (2007), 4339–4358. (Also: e-print, [arXiv:math.SP/0505546](https://arxiv.org/abs/math.SP/0505546), 2005.)
- [Titch] E. C. Titchmarsh, *Introduction to the Theory of Fourier Integrals*, second edition, Oxford Univ. Press, Oxford, 1948.
- [Tri] C. Tricot, *Curves and Fractal Dimension*, Springer-Verlag, Berlin, 1995.

- [Vin] I. M. Vinogradov (ed.), *Matematicheskaya entsiklopediya*, Sov. Entsiklopediya, Moscow, 1979, vol. 2, pp. 818–819. (Also: online, English transl., Quasi-periodic function. Yu. V. Komlenko and E. L. Tonkov (authors), *Encyclopedia of Mathematics*, public wiki monitored by an editorial board under the management of the European Mathematical Society, URL: [www.encyclopediaofmath.org](http://www.encyclopediaofmath.org))
- [Wey] H. Weyl, On the volume of tubes, *Amer. J. Math.* **61** (1939), 461–472.
- [Zäh] M. Zähle, Curvatures and currents for unions of sets with positive reach, *Geom. Dedicata* **23** (1987), 155–171.
- [Žu1] D. Žubrinić, Analysis of Minkowski contents of fractal sets and applications, *Real Anal. Exchange* No. 2, **31** (2005/2006), 315–354.
- [Žu2] D. Žubrinić, Hausdorff dimension of singular sets of Sobolev functions and applications, in: *More Progress in Analysis*, Proc. of the 5th Internat. ISAAC Congress (H. G. W. Begher and F. Nicolosi, eds.), World Scientific, 2009, pp. 793–802.
- [ŽuŽup] D. Žubrinić and V. Županović, Fractal analysis of spiral trajectories of some planar vector fields, *Bulletin des Sciences Mathématiques* No. 6, **129** (2005), 457–485.

Michel L. Lapidus  
University of California, Riverside  
Department of Mathematics  
California 92521-0135  
USA  
e-mail: [lapidus@math.ucr.edu](mailto:lapidus@math.ucr.edu)

Goran Radunović  
University of Zagreb, Faculty of Electrical Engineering and Computing  
Department of Applied Mathematics  
Unska 3, 10000 Zagreb  
Croatia  
e-mail: [goran.radunovic@fer.hr](mailto:goran.radunovic@fer.hr)

Darko Žubrinić  
University of Zagreb, Faculty of Electrical Engineering and Computing  
Department of Applied Mathematics  
Unska 3, 10000 Zagreb  
Croatia  
e-mail: [darko.zubrnic@fer.hr](mailto:darko.zubrnic@fer.hr)