

Reduction Theorems for Characteristic Functors on Finite p -Groups and Applications to p -Nilpotence Criteria

Paul Lescot

November 21st, 2006

Abstract

We formalize various properties of characteristic functors on p -groups, and discuss relationships between them. Applications to the Thompson subgroup and certain of its analogues are then given.

1 Introduction

In a now classical paper ([8]), John Thompson introduced, for p a prime number and S a p -group, the subgroup $J_R(S)$ (there denoted by $J(S)$) generated by the abelian subgroups of S of maximal rank :

$$J_R(S) \equiv_{def} \langle A \in ab(S) \mid m(A) = \max_{B \in ab(S)} m(B) \rangle, \quad (1.1)$$

where $ab(S)$ denotes the set of all abelian subgroups of S , and, for C an abelian group, $m(C)$ denotes the minimal cardinality of a generating system of C .

Later on, in [2], Glauberman modified that definition to :

$$J(S) \equiv_{def} \langle A \in ab(S) \mid |A| = \max_{B \in ab(S)} |B| \rangle. \quad (1.2)$$

Thompson had formulated a p -nilpotence criterion using J_R ; this work was later built upon by Glauberman ([2]) with his ZJ -Theorem, and by Thompson himself ([9]). For the prime $p = 2$, it is often more convenient to work with the subgroup $J_e(S)$, defined using *elementary* abelian subgroups instead of abelian ones :

$$J_e(S) \equiv_{def} \langle A \in ab_e(S) \mid |A| = \max_{B \in ab_e(S)} |B| \rangle \quad (1.3)$$

where $ab_e(S)$ denotes the set of *elementary* abelian subgroups of S .

The functors J_e , J_R and J are *excellently abelian-generated characteristic p -functors* in the sense of §3 below. In §4, we shall establish various reduction

results concerning such objects ; most notably, in certain cases, the normality of $W(S)$ in G (for $S \in \text{Syl}_p(G)$ and W a characteristic p -functor) can be inferred from the (apparently much weaker) property of *control of p -nilpotence* by W (see Theorem 4.1(2)). In the fifth paragraph, we shall specialize our results to the prime $p = 2$ and the functors J_e and $\hat{J}$ (for the definition of the last one of which see [3]), and shall henceforth refine, in a very particular case, Thompson's Factorization Theorem ([9], Theorem 1(c)), thus recovering the results of [6].

In the course of the proof some reduction lemmas of independent interest, concerning normality of p -subgroups, and control of p -nilpotence, will be established.

Our notations are standard : for G a (finite) group and p a prime number, $O_p(G)$ will denote the largest normal p -subgroup of G , $O_{p'}(G)$ the largest normal subgroup of G with order prime to p , and $Z(G)$ the center of G . We set $o(G) = |G|$, $r_e(G) = m(G)$ if G is an elementary abelian p -group for some prime p , and $r_e(G) = 0$ else; for $(x, y) \in G^2$:

$$y^x := x^{-1}yx ,$$

and, for $A \subseteq G$ and $x \in G$:

$$A^x := \{y^x | y \in A\}.$$

As usual, by a slight abuse of language, G will be said to have p -length one if $G = O_{p', p, p'}(G)$. By a *class* of groups, we shall mean a family of groups containing every subgroup and every homomorphic image of each of its elements. $\mathcal{Ab}$ will denote the class of finite abelian groups, $\mathcal{Solv}$ the class of finite solvable groups, and, for p a prime, $\mathcal{Ab}_p$ the class of finite abelian p -groups. For H a finite group, $\mathcal{C}'(H)$ will denote the class of finite groups, no section of which is isomorphic to H . For p a prime and $n \in \mathbf{N}$, $\mathcal{C}_p^n$ will denote the class of finite groups, one (*i.e.* all) of whose Sylow p -subgroups has (resp. have) nilpotency class at most n . By $ab(G)$ we shall denote the set of abelian subgroups of a group G . Finally, Σ_n will denote the symmetric group of degree n .

I am deeply grateful to the organizers of the conference "Finite Groups 2003" (Gainesville, March 6th-13th, 2003) for giving me the opportunity to present for the first time the main results of this paper.

2 A preliminary Lemma

The following result was first stated by Hayashi ([5], Lemma 3.9, p.101), though with an incomplete proof ; our own attempt at a proof ([6], Lemme) was not conclusive either (the sentence " Q , agissant sans point fixe sur le 2-groupe abélien élémentaire X , est donc cyclique" is ambiguous, as in order to thus establish the cyclicity of Q , we need to know that **each nonidentity element of Q** acts on X without fixed point, which is not obvious). Here, we shall take the opportunity to clarify the matter once and for all ; during the course of the proof, we shall feel free to use some ideas from [5] and [6].

Lemma 2.1 *Let G be a (solvable) $\{2, 3\}$ -group ; then the following statements are equivalent :*

- (1) G is Σ_4 -free , and :
- (2) $G = O_{3,2,3}(G)$.

Remark 2.2 According to Burnside's $p^a q^b$ -Theorem, the solvability hypothesis is redundant.

Proof. The implication (2) $\implies$ (1) is obvious, as the condition $G = O_{3,2,3}(G)$ is inherited by all sections of G , and $\Sigma_4 \neq O_{3,2,3}(\Sigma_4)$.

Let G denote a minimal counterexample to the statement that (1) $\implies$ (2) ; it is clear that $O_3(G) = 1$, that G possesses a unique minimal non-trivial normal subgroup X , that X is a 2-group, and that $N_0 = O_{2,3}(G) \subset G$ is the unique maximal normal subgroup of G . It follows (as $O_3\left(\frac{G}{N_0}\right) = 1$) that $\frac{G}{N_0}$ has order 2; therefore one has $O^3(G) \not\subseteq N_0$, whence $G = O^3(G)$, thus

$$O^3\left(\frac{G}{X}\right) = \frac{O^3(G)X}{X} = \frac{G}{X} .$$

But, by the minimality of G , one may write

$$O_{3,2,3}\left(\frac{G}{X}\right) = \frac{G}{X} ,$$

whence

$$\frac{G}{X} = O_{3,2}\left(\frac{G}{X}\right) .$$

Take now $Q \in \text{Syl}_3(G)$; we have just established that $QX \triangleleft G$, and the Frattini argument yields :

$$G = XN_G(Q) .$$

Let $L =_{\text{def}} N_G(Q)$; then $L \neq G$ and $G = LX$. Let us assume $L \subseteq H \subset G$; then

$$\begin{aligned} H &= H \cap G \\ &= H \cap LX \\ &= L(H \cap X) ; \end{aligned}$$

but $H \cap X \triangleleft \langle H, X \rangle = G$, whence $H \cap X = 1$ or $H \cap X = X$. In the second case, $H = LX = G$, a contradiction ; therefore $H \cap X = 1$, and $H = L(H \cap X) = L$: L is a maximal subgroup of G . Taking now $H = L$ in the above argument yields :

$$L \cap X = 1 .$$

Let $C = C_L(X)$; then $C \triangleleft LX = G$, and $X \not\subseteq C$ (else one would have $G = LX = L$, a contradiction), therefore $C = 1$. As $X \triangleleft G$, $X \subseteq O_2(G)$, whence $X \triangleleft O_2(G)$ and $Y = X \cap Z(O_2(G)) \neq 1$; but $Y \triangleleft G$, therefore $Y = X$, *i.e.* $X \subseteq Z(O_2(G))$. It follows that

$$\begin{aligned} O_2(G) &\subseteq C_G(X) \\ &= C_G(X) \cap XL \\ &= XC_L(X) \\ &= X. \end{aligned}$$

Therefore $X = O_2(G)$. Let us set $\bar{G} = \frac{G}{X}$; then $O_2(\bar{G}) = \frac{O_2(G)}{X} = 1$, and (as $\bar{G}$ is solvable)

$$C_{\bar{G}}(O_3(\bar{G})) \subseteq O_3(\bar{G}). (*)$$

Let now $\bar{t} = tX$ denote an element of order 2 in $\bar{G} = \frac{G}{X}$; according to (*), $\bar{t}$ does not centralize $O_3(\bar{G})$, therefore some $\bar{y} \in O_3(\bar{G})$ is not centralized by $\bar{t}$, thus $\bar{z} =_{def} [\bar{t}, \bar{y}] \neq 1$, $\bar{z} \in O_3(\bar{G})$, and

$$\begin{aligned} \bar{z}^{\bar{t}} &= \bar{t}^{-1} \bar{z} \bar{t} \\ &= \bar{t}^{-1} \bar{t}^{-1} \bar{y}^{-1} \bar{t} \bar{y} \bar{t} \\ &= \bar{y}^{-1} \bar{t} \bar{y} \bar{t} \\ &= (\bar{t}^{-1} \bar{y}^{-1} \bar{t} \bar{y})^{-1} \\ &= \bar{z}^{-1}. \end{aligned}$$

Let $\omega(\bar{z}) = 3^m (m \geq 1)$, and $\bar{v} =_{def} \bar{z}^{3^{m-1}}$; then $\omega(\bar{v}) = 3$ and $\bar{v}^{\bar{t}} = \bar{v}^{-1}$, whence $\langle \bar{t}, \bar{v} \rangle \simeq \Sigma_3$. Set now $V = X \langle t, t^v \rangle$; then $\frac{V}{X} = \langle \bar{t}, \bar{v} \rangle \simeq \Sigma_3$, and $O_3(V) \subseteq C_G(O_2(V)) \subseteq C_G(X) \subseteq X$, whence $O_3(V) = 1$. If $V \neq G$, then (by induction) $V = O_{3,2,3}(V)$, whence $V = O_{2,3}(V)$, $t \in O_2(V)$, $\langle t, t^v \rangle \subseteq O_2(V)$, V is a 2-group, and hence also is $\bar{V}$, a contradiction. Therefore $V = G$ and $L \simeq \frac{G}{X} = \bar{V} \simeq \Sigma_3$. It follows that $G = LX = L \rtimes X$, X (as a minimal normal subgroup of G) being a nontrivial irreducible $\mathbf{F}_2 L \simeq \mathbf{F}_2 \Sigma_3$ -module. But then X has to be isomorphic to the canonical module $\mathbf{F}_2^2$ for $\Sigma_3 \simeq SL_2(\mathbf{F}_2)$, and one obtains $G \simeq \Sigma_3 \rtimes \mathbf{F}_2^2 \simeq \Sigma_4$, a contradiction. $\square$

3 Characteristic p -functors : generalities

For p a prime number, $\mathcal{G}_p$ will denote the category of finite p -groups (morphisms in $\mathcal{G}_p$ being the group isomorphisms in the usual sense).

Definition 3.1 ([2],p.1116) *By a characteristic p -functor we shall mean a functor $K : \mathcal{G}_p \rightarrow \mathcal{G}_p$ such that, for each $P \in \mathcal{G}_p$, $K(P) \subseteq P$ and $K(P) \neq 1$ if $P \neq 1$.*

Clearly, whenever K_1 and K_2 are characteristic p -functors, $K_1 \circ K_2$ (simply denoted by $K_1 K_2$), defined by :

$$(K_1 \circ K_2)(P) \equiv_{def} K_1(K_2(P))$$

is one. Examples of characteristic p -functors include J_R , J , $\hat{J}$, J_e , Z , and Ω_n ($n \in \mathbf{N}$), the last one defined by :

$$\Omega_n(P) \equiv_{def} \{x \in P \mid x^{p^n} = 1\}.$$

A general class of characteristic p -functors is obtained *via* :

Definition 3.2 *Let φ denote a mapping from $\mathcal{A}b_p$ to $\mathbf{N}$, invariant under isomorphisms, and such that*

$$A \neq 1 \implies \varphi(A) \geq 1 ;$$

then, for P a p -group, let

$$K_\varphi(P) \equiv_{def} \{A \text{ abelian subgroup of } P \mid \varphi(A) = \max_{B \subseteq A; B \text{ abelian}} \varphi(B) > 0\}.$$

It is easily seen that K_φ is a characteristic p -functor ; such characteristic p -functors will be termed *excellently abelian generated* . Clearly, J , J_R and J_e are such ; in fact , $J = K_o$, $J_R = K_m$ and $J_e = K_{r_e}$.

Definition 3.3 *The characteristic p -functor W is termed excellent if, whenever G is a finite group, $P \in \text{Syl}_p(G)$, $x \in G$, and $W(P) \subseteq Q \subseteq P^x$, then $W(P) = W(Q) = W(P^x)(= W(P)^x)$. In particular, $W(P)$ is weakly closed in P , and characteristic in any p -subgroup of G that contains it.*

Lemma 3.4 *Any excellently abelian generated characteristic p -functor is excellent.*

Proof. For S a p -group, let

$$r_\varphi(S) =_{def} \max_{A \in \mathcal{A}b(S)} \varphi(A) .$$

Let us assume that $K_\varphi(P) \subseteq Q \subseteq P^x$, and let $A_0 \in \mathcal{A}b(P)$ such that

$$\varphi(A_0) = \max_{A \in \mathcal{A}b(P)} \varphi(A) = r_\varphi(P) .$$

Obviously,

$$\begin{aligned}
r_\varphi(Q) &\leq r_\varphi(P^x) \\
&= \max_{A \in ab(P^x)} \varphi(A) \\
&= \max_{C \in ab(P)} \varphi(C^x) \\
&= \max_{C \in ab(P)} \varphi(C) \\
&\quad (\text{as } \varphi \text{ is invariant under isomorphisms}) \\
&= r_\varphi(P) \\
&= \varphi(A_0) \\
&\leq r_\varphi(Q) \text{ (as } A_0 \subseteq K_\varphi(P) \subseteq Q).
\end{aligned}$$

Therefore $r_\varphi(P) = r_\varphi(Q)$, whence

$$\begin{aligned}
K_\varphi(Q) &= \langle A \in ab(Q) \mid \varphi(A) = r_\varphi(Q) \rangle \\
&= \langle A \in ab(Q) \mid \varphi(A) = r_\varphi(P) \rangle \\
&= \langle A \in ab(P) \mid \varphi(A) = r_\varphi(P) \rangle \\
&= K_\varphi(P)
\end{aligned}$$

(because $A \in ab(P)$ and $\varphi(A) = r_\varphi(P)$ yield $A \subseteq K_\varphi(P) \subseteq Q$)

Incidentally we have shown that $r_\varphi(Q) = r_\varphi(P^x)$, whence $K_\varphi(Q) \subseteq K_\varphi(P^x)$ and $K_\varphi(P) = K_\varphi(Q) \subseteq K_\varphi(P^x) = (K_\varphi(P))^x$, and equality all along follows. $\square$

4 A Reduction Theorem

Let p , W and $\mathcal{C}$ denote respectively a prime number, a characteristic p -functor, and a class of groups ; the following properties of the triple $(W, \mathcal{C}, p)$ will be considered (S denoting a Sylow p -subgroup of the group G) :

(P1) For each $G \in \mathcal{C}$, one has

$$G = N_G(W(S))O_{p'}(G) .$$

(P2) For each p -solvable $G \in \mathcal{C}$, one has

$$G = N_G(W(S))O_{p'}(G) .$$

(P3) For each solvable $G \in \mathcal{C}$, one has

$$G = N_G(W(S))O_{p'}(G) .$$

(P4) For each solvable $G \in \mathcal{C}$, all of whose Sylow q -subgroups for all primes $q \neq p$ are abelian, one has

$$G = N_G(W(S))O_{p'}(G) .$$

- (P5) W controls p -length 1 in $\mathcal{C}$, i.e. for each p -solvable $G \in \mathcal{C}$, if $N_G(W(S))$ has p -length one then G has p -length one.
- (P6) W controls p -nilpotence in $\mathcal{C}$, i.e. for each $G \in \mathcal{C}$, if $N_G(W(S))$ is p -nilpotent then G is p -nilpotent.

Stellmacher's result ([7]) asserts the existence of a (non-explicit) characteristic 2-functor W such that (P1)(and hence (P2),..., (P6)) hold for $(W, \mathcal{C}'(\Sigma_4), 2)$.

Theorem 4.1 (1) One has $(P1) \Rightarrow (P2) \Rightarrow (P3) \Rightarrow (P4) \Rightarrow (P6)$, and $(P3) \Rightarrow (P5) \Rightarrow (P6)$.

(2) If $p = 2$, $W(S) \subseteq \Omega_1(S)$ for all S , and either

(i) $\mathcal{C} \subseteq \mathcal{C}_{2,2}$ and W is excellent ,

or

(ii) W is excellently abelian generated,

then $(P6) \Rightarrow (P2)$, and hence properties $(P2), \dots, (P6)$ are equivalent.

Proof. (1) The implications $(P1) \Rightarrow (P2) \Rightarrow (P3) \Rightarrow (P4)$ are trivial.

In order to establish that $(P3) \Rightarrow (P5)$, let us assume (P3), let G denote a counterexample to (P5) with minimal order. We shall use arguments similar to Bauman's in [1], pp.388–389. If $O_{p'}(G) \neq 1$, let $\bar{G} =_{def} \frac{G}{O_{p'}(G)}$; then one has :

$$\begin{aligned} N_{\bar{G}}(W(\bar{S})) &= N_{\bar{G}}\left(\frac{W(S)O_{p'}(G)}{O_{p'}(G)}\right) \\ &= \frac{N_G(W(S))O_{p'}(G)}{O_{p'}(G)} \\ &\quad \text{(by the Frattini argument)} \\ &\simeq \frac{N_G(W(S))}{N_G(W(S)) \cap O_{p'}(G)} . \end{aligned}$$

Therefore $N_{\bar{G}}(W(\bar{S}))$ has p -length one, whence, by induction (as $\bar{G} \in \mathcal{C}$ and $\bar{G}$ is p -solvable), $\bar{G}$ has p -length one, hence so has G , a contradiction. Thus $O_{p'}(G) = 1$, whence (as G is p -solvable) $C_G(O_p(G)) \subseteq O_p(G)$; in particular, $O_p(G) \neq \{1\}$. Let now $\bar{G} = \frac{G}{O_p(G)}$, and let $\bar{H} = N_{\bar{G}}(W(\bar{S}))$; if $H = G$, then $W(\bar{S}) \triangleleft \bar{G}$, thus $W(\bar{S}) \subseteq O_p(\bar{G}) = 1$, $W(\bar{S}) = 1$, $\bar{S} = 1$, $S = O_p(G)$, $W(S) = W(O_p(G)) \triangleleft G$, and $G = N_G(W(S))$ has p -length one, a contradiction. Therefore $H \subset G$; as $N_H(W(S)) \subseteq N_G(W(S))$ has p -length one, so has H by induction, hence so has $\bar{H}$, hence so has $\bar{G}$, again by induction ($\bar{G}$ and H both belonging to $\mathcal{C}$). Let $\bar{K} = O_{p'}(\bar{G})$; it appears that $\bar{S}\bar{K} \triangleleft \bar{G}$, hence $SK \triangleleft G$; if $SK \neq G$, one finds by induction that SK has p -length 1; but $SK \triangleleft G$, whence $O_{p'}(SK) \triangleleft G$ and $O_{p'}(SK) \subseteq O_{p'}(G) = 1$. Therefore

$S \triangleleft SK$, whence $S = O_p(SK) \triangleleft G$, and again $W(S) \triangleleft G$ and $G = N_G(W(S))$, a contradiction. Therefore $G = SK$, and $\bar{G} = \bar{S}\bar{K}$.

For $q \in \pi(\bar{K})$, let $\bar{Q}$ denote a Sylow q -subgroup of $\bar{K}$; the total number of Sylow q -subgroups of $\bar{K}$ is $|\bar{K} : N_{\bar{K}}(\bar{Q})| \neq 0[p]$, therefore one of them, $\bar{K}_q$, is $\bar{S}$ -invariant. If, for each $q \in \pi(\bar{K})$, one has $SK_q \neq G$, then, by induction, SK_q has p -length one; but $O_{p'}(SK_q) \subseteq C_G(O_p(SK_q)) \subseteq C_G(O_p(G)) \subseteq O_p(G)$, thus $O_{p'}(SK_q) = 1$ and $S \triangleleft SK_q$, thus $K_q \subseteq N_G(S)$, hence

$$\bar{K} = \langle \bar{K}_q | q \in \pi(\bar{K}) \rangle \subseteq \overline{N_G(S)}$$

and $S \triangleleft SK = G$, a contradiction. Thus for some prime q one has $G = SK_q$, and it appears that G is solvable (in fact, a solvable $\{p, q\}$ -group for some prime q). But now (P3) yields that $G = N_G(W(S))$, whence G has p -length one, a contradiction (in this proof, due to the hypotheses on $\mathcal{C}$, all the groups that appear *belong to* $\mathcal{C}$; such will be the case in all subsequent similar reasonings).

Assuming (P4), let G denote a counterexample to (P6), with minimal order; then Thompson's arguments ([8], pp.43–44) yield that $O_{p'}(G) = 1$, $O_p(G) \neq 1$ and G is a $\{p, q\}$ -group with (elementary) abelian Sylow subgroups for some prime $q \neq p$. But then (P4) yields that $G = N_G(W(S))$, whence G has p -length one, a contradiction. Therefore (P4) $\Rightarrow$ (P6) is established.

In order to establish that (P5) $\Rightarrow$ (P6), the same argument works; here, we only need Thompson's reduction up to an earlier point, *viz.* $O_{p'}(G) = 1$ and G p -solvable.

(2) Let us assume all the conditions in (2), and let G denote a minimum counterexample to (P6) $\Rightarrow$ (P2); it is clear, as usual, that $O_{2'}(G) = 1$, and then (by the same reasoning as in (1)) that $O_{2'}(H) = 1$ for any subgroup H of G containing S , and therefore that $M \equiv N_G(W(S))$ is the unique maximal subgroup of G containing S . Let $\bar{G} = \frac{G}{O_2(G)}$; then $\bar{G}$ is 2-solvable, and $\bar{M}$ is the unique maximal subgroup of G containing $\bar{S}$. By induction, one has

$$\begin{aligned} \bar{G} &= N_{\bar{G}}(W(\bar{S}))O_{2'}(\bar{G}) \\ &= N_{\bar{G}}(W(\bar{S}))(\bar{S}O_{2'}(\bar{G})); \end{aligned}$$

the two factors on the right-hand side of this equality contain $\bar{S}$, whence at least one is not contained in $\bar{M}$, *i.e.* either $N_{\bar{G}}(W(\bar{S})) = \bar{G}$ or $\bar{G} = \bar{S}O_{2'}(\bar{G})$. The first possibility leads to a contradiction as in the proof that (P3) $\Rightarrow$ (P5); therefore $\bar{G} = \bar{S}O_{2'}(\bar{G})$, *i.e.* G has 2-length one.

As $\bar{S}$ is contained into a unique maximal subgroup of $\bar{G}$ ($\bar{M}$), $O_{2'}(\bar{G})$ possesses a unique maximal $\bar{S}$ -invariant proper subgroup: $O_{2'}(\bar{G}) \cap \bar{M}$. It follows, first, that $O_{2'}(\bar{G})$ is a q -group for some prime $q \neq 2$: $O_{2'}(\bar{G}) = \bar{Q}$ ($Q \in \text{Syl}_q(G)$), and therefore $G = SQ$ is a solvable $\{2, q\}$ -group, and secondly that $\bar{S}$ acts irreducibly on $\frac{\bar{Q}}{\Phi(\bar{Q})}$; in particular, $Z(\bar{S})$ is cyclic.

Let $N \equiv_{def} W(S)^G > \triangleleft G$; then $O_{2'}(N) = 1$, and $S \cap N \in \text{Syl}_2(N)$. If $N < G$, the minimality of G yields:

$$\begin{aligned}
N &= N_N(W(S \cap N))O_{2'}(N) \\
&= N_N(W(S \cap N)) .
\end{aligned}$$

But $W(S) \subseteq S \cap N \subseteq S$, whence $W(S) = W(S \cap N)$, as W is excellent (in case (i) by assumption, and in case (ii) by Lemma 3.4). The Frattini argument now yields that :

$$\begin{aligned}
G &= NN_G(S \cap N) \\
&\subseteq NN_G(W(S \cap N)) \\
&\subseteq N_G(W(S \cap N)) \\
&\subseteq G ,
\end{aligned}$$

whence $G = N_G(W(S \cap N)) = N_G(W(S))$ is 2-nilpotent, a contradiction. Therefore $N = G$, *i.e.* $G = \langle W(S)^G \rangle$; thence

$$\begin{aligned}
\bar{G} &= \langle \overline{W(S)}^{\bar{G}} \rangle \\
&= \langle \overline{W(S)}^{\bar{S}\bar{Q}} \rangle \\
&= \langle \overline{W(S)}^{\bar{Q}} \rangle \\
&\subseteq \overline{W(S)}\bar{Q} \text{ (as } \bar{Q} \triangleleft \bar{G} \text{)} ,
\end{aligned}$$

and $\bar{S} = \bar{S} \cap \overline{W(S)}\bar{Q} = \overline{W(S)}(\bar{S} \cap \bar{Q}) = \overline{W(S)}$, *i.e.* $S = W(S)O_2(G)$.

In case (ii), let $W = K_\varphi$; then $W(S) \not\subseteq O_2(G)$ (else one would have $S = W(S)O_2(G) = O_2(G) \triangleleft G$), whence there is an abelian subgroup A of S with $\varphi(A) = r_\varphi(P)$ and $A \not\subseteq O_2(G)$. Let $N = \langle A^G \rangle \triangleleft G$; if $N \neq G$, then, by induction, it follows as above that $W(S \cap N) \triangleleft N$ whence $W(S \cap N) \subseteq O_2(N) \subseteq O_2(G)$. But

$$\varphi(A) \leq r_\varphi(S \cap N) \leq r_\varphi(S) = \varphi(A)$$

whence $\varphi(A) = r_\varphi(S \cap N)$ and $A \subseteq K_\varphi(S \cap N) = W(S \cap N) \subseteq O_2(N) \subseteq O_2(G)$, a contradiction. Therefore $G = \langle A^G \rangle$, whence

$$\begin{aligned}
\bar{G} &= \langle \bar{A}^{\bar{G}} \rangle \\
&= \langle \bar{A}^{\bar{S}\bar{Q}} \rangle \\
&= \langle \bar{A}^{\bar{S}} \rangle \bar{Q} \text{ (as } \bar{Q} \triangleleft \bar{G} \text{)} ;
\end{aligned}$$

therefore

$$\begin{aligned}
\bar{S} &= \bar{S} \cap \bar{G} \\
&= \bar{S} \cap \langle \bar{A}^{\bar{S}} \rangle \bar{Q} \\
&= \langle \bar{A}^{\bar{S}} \rangle (\bar{S} \cap \bar{Q}) \\
&= \langle \bar{A}^{\bar{S}} \rangle .
\end{aligned}$$

By a well-known property of p -groups, it follows that $\bar{S} = \bar{A}$; in particular, $\bar{S}$ is abelian.

In case (i), $\mathcal{C} \subseteq \mathcal{C}_{2,2}$, *i.e.* $cl(S) \leq 2$, whence

$$\begin{aligned} [S, S] &\subseteq Z(S) \\ &\subseteq C_G(O_2(G)) \\ &\subseteq O_2(G) \end{aligned}$$

(by the solvability of G and the Hall-Higman Lemma), whence, again, $\bar{S}$ is abelian. Therefore, $\bar{S}$ is abelian in both cases, (i) and (ii). Now, from the fact that $Z(\bar{S})$ is cyclic, follows that $\bar{S}$ itself is. But $\bar{S} = \overline{W(S)} \subseteq \Omega_1(\bar{S}) \subseteq \Omega_1(\bar{S})$ (by the hypothesis); therefore $\bar{S}$ has order 2.

Now, as $\bar{S}$ acts irreducibly on the $\mathbf{F}_q$ -module $M = \frac{\bar{Q}}{\Phi(\bar{Q})}$, the nontrivial element $\bar{t}$ of $\bar{S}$ either centralizes each element of M , or inverts each element of M ; now, irreducibility forces $|M| = q$, *i.e.* $\frac{\bar{Q}}{\Phi(\bar{Q})} = M$ is cyclic ; but then so are $\bar{Q}$, and $Q \simeq \bar{Q}$.

Let now $\bar{H} = \bar{S}\Phi(\bar{Q})$; then $H < G$ (in fact, $|G : H| = q$), and $S \subseteq H$. Therefore H is contained in $M = N_G(W(S))$, whence

$$\begin{aligned} [\bar{S}, \Phi(\bar{Q})] &= [\overline{W(S)}, \Phi(\bar{Q})] \\ &\subseteq [\overline{W(S)}, \bar{H}] \cap \Phi(\bar{Q}) \\ &\subseteq [\overline{W(S)}, \bar{M}] \cap \Phi(\bar{Q}) \\ &\subseteq \overline{W(S)} \cap \Phi(\bar{Q}) \\ &= 1, \end{aligned}$$

i.e. $\bar{S}$ centralizes $\Phi(\bar{Q})$. If $|\bar{Q}| \geq q^2$, then $\Omega_1(\bar{Q}) \subseteq \Phi(\bar{Q})$, whence $\bar{S}$ centralizes $\Omega_1(\bar{Q})$, and therefore $\bar{S}$ centralizes $\bar{Q}$, a contradiction. Thus $|\bar{Q}| = q$, and $\bar{G} = \bar{S}\bar{Q}$ is dihedral of order $2q$; it follows that $\bar{S}$ is a maximal subgroup of $\bar{G}$, *i.e.* S is a maximal subgroup of G . Therefore $S = M = N_G(W(S))$, and $N_G(W(S))$ is 2-nilpotent ; but now (P6) yields that G itself is 2-nilpotent, a contradiction. $\square$

5 Of J_e and $\hat{J}$

By a well-known variation([4],Theorem 1(c), and Remarks p.372) on Thompson's factorization([9]), any solvable Σ_3 -free finite group G with Sylow 2-subgroup S satisfies :

$$G = N_G(J_e(S))C_G(Z(S))O_{2'}(G). \quad (5.1)$$

In [3] Glauberman introduced a new characteristic functor $\hat{J}$ having the property that, for each 2-group S , one has :

$$J_e(S) \subseteq \hat{J}(S) \subseteq S. \quad (5.2)$$

For this functor he was able to prove ([3], Theorem 7.4, p.48) that, for any 2-constrained Σ_4 -free finite group G and each $S \in \text{Syl}_2(G)$, one had :

$$G = N_G(\hat{J}(S))C_G(Z(S))O_{2'}(G) \quad (5.3).$$

By (5.2) one finds $J_e(S) = J_e(\hat{J}(S))\text{char}\hat{J}(S)$ whence

$$N_G(\hat{J}(S)) \subseteq N_G(J_e(S)) ;$$

(5.3) is therefore stronger than (5.1).

In the particular case that S has nilpotence class at most two, we can state

Theorem 5.1 *Let G be a 2-constrained, Σ_4 -free finite group with Sylow 2-subgroup S of nilpotence class at most two ; then one has :*

$$G = N_G(\hat{J}(S))O_{2'}(G).$$

By the above remark follows

Corollary 5.2 *In the situation of the Theorem,*

$$G = N_G(J_e(S))O_{2'}(G).$$

Thus one can assert

Corollary 5.3 *Let G be a finite solvable Σ_4 -free group with Sylow 2-subgroup S of class at most two ; then :*

$$G = N_G(J_e(S))O_{2'}(G).$$

In other words, $(J_e, \mathcal{C}'(\Sigma_4) \cap \text{Solv}, 2)$ satisfies (P1), and hence (P2), ..., (P6).

This Corollary was first proved by the author in [6].

Proof. of Theorem 5.1. Let G be a counterexample of minimal order.

(1) $\mathbf{O}_{2'}(\mathbf{G}) = 1$.

If not, $\bar{G} = \frac{G}{\mathbf{O}_{2'}(G)}$ is of smaller order than G and satisfies the hypothesis, whence

$$\bar{G} = N_{\bar{G}}(\hat{J}(\bar{S}))O_{2'}(\bar{G}) = N_{\bar{G}}(\hat{J}(\bar{S})).$$

But the canonical map $S \rightarrow \frac{SO_{2'}(G)}{\mathbf{O}_{2'}(G)} = \bar{S}$ is an isomorphism, whence

$$\hat{J}(\bar{S}) = \frac{\hat{J}(S)O_{2'}(G)}{\mathbf{O}_{2'}(G)} \text{ and}$$

$$N_{\bar{G}}(\hat{J}(\bar{S})) = \frac{N_G(\hat{J}(S)O_{2'}(G))}{\mathbf{O}_{2'}(G)} = \frac{N_G(\hat{J}(S))O_{2'}(G)}{\mathbf{O}_{2'}(G)},$$

by the Frattini argument. Thus we get $G = N_G(\hat{J}(S))O_{2'}(G)$, a contradiction.

- (2) $\mathbf{C}_G(\mathbf{O}_2(\mathbf{G})) \subseteq \mathbf{O}_2(\mathbf{G})$.

Obvious, because G is 2-constrained and $O_{2'}(G) = 1$.

- (3) $\mathbf{M} = \mathbf{N}_G(\hat{\mathbf{J}}(\mathbf{S}))$ is the unique maximal subgroup of $\mathbf{G}$ that contains $\mathbf{S}$.

By hypothesis $M \subset G$. Let H be a proper subgroup of G containing S ; one has $O_2(G) \subseteq S \subseteq H$, whence (as in the proof of Theorem 4.1(1))

$$O_2(G) \subseteq O_2(H)$$

and :

$$\begin{aligned} C_H(O_2(H)) &= H \cap C_G(O_2(H)) \\ &\subseteq H \cap C_G(O_2(G)) \\ &\subseteq H \cap O_2(G) \text{ (by (2))} \\ &\subseteq O_2(H). \end{aligned}$$

Therefore $O_{2'}(H) = 1$ and H is 2-constrained with Sylow 2-subgroup S ; the minimality of G now yields :

$$\begin{aligned} H &= N_H(\hat{J}(S))O_{2'}(H) = N_H(\hat{J}(S)) \\ &\subseteq N_G(\hat{J}(S)) = M. \end{aligned}$$

Thus M is a proper subgroup of G that contains any proper subgroup of G containing S ; the result follows.

- (4) $\mathbf{Z}(\mathbf{S}) \subseteq \mathbf{Z}(\mathbf{G})$.

By (5.3) one has

$$G = N_G(\hat{J}(S))C_G(Z(S))O_{2'}(G) = MC_G(Z(S)) ;$$

thus $S \subseteq C_G(Z(S)) \not\subseteq M$, whence $C_G(Z(S)) = G$ by (3).

- (5) $\mathbf{G}$ centralizes $\frac{\mathbf{O}_2(\mathbf{G})}{\mathbf{Z}(\mathbf{G})}$.

Let $C = C_G(\frac{O_2(G)}{Z(G)}) \triangleleft G$; then

$$[S, O_2(G)] \subseteq [S, S] \subseteq Z(S) \subseteq Z(G)$$

(by (4) and the hypothesis on S). It follows that $S \subseteq C$, whence

$$G = CN_G(S) ,$$

again by the Frattini argument. If C were different from G , one would have $C \subseteq M$ (because of (3)) and

$$G = CN_G(S) \subseteq MN_G(S) \subseteq M.M = M ,$$

a contradiction. Thus $C = G$.

(6) **The End.**

By (5) one has $[G, O_2(G)] \subseteq Z(G)$, i.e.

$$[G, O_2(G), G] = [O_2(G), G, G] = 1.$$

Philip Hall's Three Subgroups Lemma now yields

$$[G, G, O_2(G)] = 1 ,$$

that is :

$$G' \subseteq C_G(O_2(G)) ,$$

whence $G' \subseteq O_2(G)$ by (2). Therefore $H = \frac{G}{O_2(G)}$ is an abelian group with $O_2(H) = 1$, i.e. an abelian $2'$ -group ; it appears that $S = O_2(G) \triangleleft G$, whence $\hat{J}(S) \triangleleft G$, thus $G = M$ and again a contradiction ensues. This concludes the proof. $\square$

Remark 5.4 It seems difficult to generalize directly Corollary 5.2, and even Corollary 5.3, as the counter-examples to the ZJ -Theorem for $p = 2$ given by Glauberman in the last paragraph of [2] show. Such a counterexample G is solvable, with Sylow 2-subgroup S of nilpotence class 3 (this is not difficult to see), and S possesses a unique abelian subgroup of maximal order A , that is elementary abelian. Therefore $J_e(S)$, $J_R(S)$, $J(S)$ and $ZJ(S)$ all coincide with A , and neither is normal in G .

References

- [1] S.F.Bauman, *Groups with p -length 1* Journal of Algebra **8** (1968), 388–392.
- [2] G.Glauberman, *A characteristic subgroup of a p -stable group* Canad. J. Math. **20** (1968), 1101–1135.
- [3] G.Glauberman, *Factorizations in local subgroups of finite groups* CBMS series, vol.33, Amer. Math. Soc., 1977.
- [4] D. Gorenstein, *Finite simple groups* Plenum Press, New York, 1982.
- [5] M.Hayashi, *2-factorization in finite groups* Pacific J. Math. **84(1)** (1979), 97–142.
- [6] P.Lescot, *Sur la factorisation de Thompson* Revue de Mathématiques Spéciales **99(5)** (1989), 197–198.
- [7] B.Stellmacher, *A characteristic subgroup of Σ_4 -free groups* Israel Journal of Mathematics **94** (1996), 367–379.
- [8] J.G.Thompson, *Normal p -complements for finite groups* J.Algebra **1** (1964), 43–46.
- [9] J.G.Thompson, *Factorizations of p -solvable groups* Pacific J. Math. **16** (1966) 371–372.

INSSET–UNIVERSITÉ DE PICARDIE, 48 Rue Raspail, 02100 Saint–Quentin (FRANCE), paul.lescot@insset.u-picardie.fr, Fax 00 33 (0)3 23 62 89 35
LAMFA, Faculté de Mathématiques et d’Informatique, 33, Rue Saint-Leu, 80039 Amiens Cédex, Phone 03 22 82 79 70, Fax 03 22 82 78 38, paul.lescot@u-picardie.fr