

Exploiting $\mathcal{N} = 2$ in consistent coset reductions of type IIA

Davide Cassani^{a,b} and Amir-Kian Kashani-Poor^c

^a *Laboratoire de Physique Théorique*, École Normale Supérieure,
24 rue Lhomond, 75231 Paris Cedex 05, France*

^b *Dipartimento di Fisica, Università di Roma “Tor Vergata”
Via della Ricerca Scientifica, 00133 Roma, Italy*

^c *Institut des Hautes Études Scientifiques
Le Bois-Marie, 35, route de Chartres
91440 Bures-sur-Yvette, France*

Abstract

We study compactifications of type IIA supergravity on cosets exhibiting $SU(3)$ structure. We establish the consistency of the truncation based on left-invariance, providing a justification for the choice of expansion forms which yields gauged $\mathcal{N} = 2$ supergravity in 4 dimensions. We explore $\mathcal{N} = 1$ solutions of these theories, emphasizing the requirements of flux quantization, as well as their non-supersymmetric companions. In particular, we obtain a no-go result for de Sitter solutions at string tree level, and, exploiting the enhanced leverage of the $\mathcal{N} = 2$ setup, provide a preliminary analysis of the existence of de Sitter vacua at all string loop order.

*Unité mixte du CNRS et de l'École Normale Supérieure associée à l'Université Pierre et Marie Curie Paris 6, UMR 8549.

1 Introduction

In the era of LHC, much effort is being invested in finding phenomenologically viable string vacua. Much of this work takes place by considering compactifications to $\mathcal{N} = 1$ theories in 4d. In this paper, we will focus instead on a framework which yields 4 dimensional theories that have $\mathcal{N} = 2$ symmetry realized off-shell. While the $\mathcal{N} = 1$ setup allows for more flexibility in choosing the various ingredients of the theory, and hence (currently) permits the construction of more realistic vacua, the increased rigidity of the $\mathcal{N} = 2$ setup has the advantage of allowing a more exhaustive treatment of α' , string loop, and foreseeably even brane instanton corrections. An impressive example of the power of the $\mathcal{N} = 2$ framework is the recent proof [1] that $\mathcal{N} = 2$ gauged supergravities without vector multiplets do not permit de Sitter vacua, in spite of the presence of such solutions in the one-brane-instanton approximation [2]. Studying theories in the $\mathcal{N} = 2$ framework hence presents one promising avenue towards assessing the viability of the approximations that are necessary to get off the ground in less supersymmetric frameworks.

The best studied example of $\mathcal{N} = 2$ theories obtained from string theory are type II Calabi-Yau compactifications [3, 4]. The differential operators governing the geometric moduli problem of the internal Calabi-Yau manifolds turn out to coincide with the mass operators of the supergravity theory. Unobstructed deformations hence give rise to massless excitations, resulting in the beautiful identification between the massless scalar fields of these theories, whose VEVs parametrize a family of supergravity solutions, and the geometric moduli of the Calabi-Yau. The masslessness of the scalars is protected by supersymmetry, as $\mathcal{N} = 2$ forbids a potential in the case of uncharged matter. In [5], the study of type II compactifications on $SU(3)$ structure manifolds was initiated (recall that Calabi-Yau manifolds satisfy the stronger condition of $SU(3)$ holonomy). This setup is more akin to the phenomenologically motivated $\mathcal{N} = 1$ analyses: solutions of the supergravity equations of motion on these internal manifolds require the presence of background fluxes [6, 7], and compactification gives rise to 4d $\mathcal{N} = 2$ gauged supergravity theories [8, 9], which, in contrast to the Calabi-Yau case with uncharged matter, exhibit a potential for the scalar fields in the theory. The increased phenomenological viability comes at a price: the very presence of a potential makes it unlikely that the choice of light degrees of freedom of the theory can be associated to a geometric moduli problem. Indeed, a systematic approach to a reduction ansatz for these theories is still lacking. Following our work in [10] and [11], we here pursue an alternative approach towards justifying the reduction ansatz, that of consistent truncation: obtaining a field theory with a finite number of fields upon compactification requires truncating most of the degrees of freedom of the higher dimensional theory; this truncation is called consistent when all solutions to the lower dimensional equations of motion lift to solutions of the higher dimensional theory. Note the contrast to a Kaluza-Klein reduction [12], which is an expansion valid around a single 10 dimensional solution (hence referred to as a base-point dependent reduction in [13]).

Consistently truncated lower dimensional field theories are powerful allies in studying the

vacuum structure of the higher dimensional *string* theory. This is partially a consequence of computational techniques being more refined in lower dimensions. E.g., various leading non-trivial contributions in α' to the 10d type II supergravity action have been determined [14, 15, 16]. One may hope to establish the complete action to this order by 10d supersymmetric completion [17]. However, the 10d supersymmetry equations have simply proved too cumbersome to date. By contrast, the supersymmetric completion of the contribution of these terms to the 4d $\mathcal{N} = 2$ supergravity action is readily available, yielding the full string tree level and one loop corrected action. In fact, in 4d we can, as we will discuss, even draw conclusions regarding the all string loop corrected action. Studying the lower dimensional theory is however not merely a question of computational convenience. An effective higher dimensional description of worldsheet or brane instantons is even conceptually problematic.

In [11], it was shown that expansion forms can be defined on Nearly Kähler manifolds that satisfy the conditions of [13], implying that the reduction of the type IIA action based on these forms yields $\mathcal{N} = 2$ gauged supergravity in 4d. It was further demonstrated that the truncation in this setting is consistent in the supersymmetric sector (i.e. 4d solutions preserving $\mathcal{N} = 1$ supersymmetry lift). In this paper, we shift our focus to certain coset spaces which subsume the currently known set of 6d Nearly Kähler manifolds. We introduce these spaces in section 2. Considering the emphasis on base point independence of the reduction, it was perhaps somewhat disappointing that the theories based on Nearly Kähler reduction yielded a single supersymmetric vacuum for a given choice of fluxes. Cosets by contrast permit multiple $\mathcal{N} = 1$ solutions for a given choice, which are all accessible via the 4d theory. We demonstrate this in section 3. Due to flux quantization, the solutions come in a discrete family. We perform the required K-theory analysis. In section 4, we demonstrate that the left-invariant coset reductions represent a consistent truncation by establishing that the 10d equations of motion reduce to the 4d equations following from the appropriate $\mathcal{N} = 2$ action. This extends the analysis of [10] beyond the RR sector and overcomes the restriction to consistency merely of the supersymmetric sector [11, 18]. Fueled by this result, we turn to the study of non-supersymmetric vacua of the 4d theories in sections 5 and 6. We find several non-supersymmetric Nearly Kähler companions to the solution of section 3 and study their stability, in particular with regard to deformations away from the Nearly Kähler locus. We also consider the question of the existence of de Sitter vacua, which has received some attention recently in the type IIA context [19, 1, 20, 21]. We demonstrate that such vacua are absent at string tree level (we prove this result in greater generality than the coset context: it is valid for any gauged supergravity with merely the universal tree-level hypermultiplet, irrespective of the specifics of the vector multiplet sector). Due to the increased leverage in the $\mathcal{N} = 2$ setup, we are able to push this analysis beyond tree level. We obtain the full string loop corrected potential, which evades the tree-level no-go theorem, and uncover a necessary condition on the contribution of the NS sector to the potential for de Sitter vacua to be possible.

2 Introducing the internal geometries

We consider dimensional reductions of massive type IIA supergravity on left coset spaces $M_6 = G/H$ endowed with a left-invariant $SU(3)$ structure. An exhaustive list of such cosets was provided in ref. [22] (see section 1 and in particular table 1 therein). In the following, we are going to focus on the cosets whose $SU(3)$ structure cannot be further reduced to $SU(2)$, namely

$$\frac{SU(3)}{U(1) \times U(1)} \quad , \quad \frac{Sp(2)}{S(U(2) \times U(1))} \quad , \quad \frac{G_2}{SU(3)} \quad , \quad (2.1)$$

where $S(U(2) \times U(1))$ is non-maximally embedded in $Sp(2)$.

It is easy to see that a reduction performed on these manifolds by expanding the higher dimensional fields in a basis of left-invariant forms satisfies the constraints of [13] and therefore yields a gauged $\mathcal{N} = 2$ supergravity in 4d.

The remaining cosets listed in [22] have vanishing Euler characteristic and admit a left-invariant vector: their $SU(3)$ structure group is therefore further reduced to at least $SU(2)$. For these cosets, the $\mathcal{N} = 2$ reduction ansatz based on the presence of $SU(3)$ structure can be more naturally enlarged to include the whole set of left-invariant forms, possibly yielding a further extended supergravity ($\mathcal{N} \geq 4$) in 4d.

The only non-vanishing torsion classes¹ characterizing the $SU(3)$ structure of the cosets (2.1) are W_1 and W_2 , i.e. the $SU(3)$ invariant 2- and 3-form J and Ω satisfy

$$\begin{aligned} dJ &= \frac{3}{2} \text{Im}(\bar{W}_1 \Omega) \quad , \\ d\Omega &= W_1 J \wedge J + W_2 \wedge J \quad . \end{aligned} \quad (2.2)$$

In fact, $\frac{G_2}{SU(3)}$ allows just $W_1 \neq 0$ and is therefore a Nearly Kähler manifold. The cosets $\frac{SU(3)}{U(1) \times U(1)}$ and $\frac{Sp(2)}{S(U(2) \times U(1))}$ also admit a region in the $SU(3)$ structure parameter space in which they are Nearly Kähler, but in general, their W_2 torsion class does not vanish. Since W_1 and W_2 can be chosen purely imaginary, these cosets fall into the class of ‘half-flat’ manifolds, characterized by $\text{Re } W_1 = \text{Re } W_2 = W_4 = W_5 = 0$ [24].

A description of the coset spaces (2.1) was given e.g. in [25]. In the context of $SU(3)$ structure compactifications of (massive) type IIA supergravity, supersymmetric AdS_4 backgrounds on these manifolds have recently been found in [22, 26, 27, 28] and further discussed in [29], while refs. [30, 20] study the properties of the associated effective 4d $\mathcal{N} = 1$ supergravity in the presence of orientifold projections (see also [27] for a previous work considering the coset $\frac{SU(3)}{U(1) \times U(1)}$). Type IIA reduction on Nearly Kähler manifolds has been worked out in [11]. The cosets (2.1) have also been employed in [31] for heterotic dimensional reductions.

¹For a review of $SU(3)$ structures and their torsion classes, see e.g. subsection 3.2 of ref. [23].

2.1 The expansion forms

In the following we provide the most general left-invariant positive-definite metric for each coset (2.1), as well as a basis for all the left-invariant differential forms, on which we are going to expand the supergravity fields.

We define the 6d coset spaces (2.1) as in ref. [22], and in particular adopt the set of group structure constants listed therein. The same reference also provides a summary of the needed mathematical notions about coset spaces, while a more extended review can be found e.g. in [25].

Using the local coframe² $\{e^{\underline{m}}\}$ inherited from G , a differential form on the coset G/H reads $\omega_k = \frac{1}{k!} \omega_{\underline{m}_1 \dots \underline{m}_k} e^{\underline{m}_1} \wedge \dots \wedge e^{\underline{m}_k}$. This is invariant under the left action of G if its components are constant and satisfy the following relation involving the G structure constants

$$f^{\underline{p}}_{i[\underline{m}_1} \omega_{\underline{m}_2 \dots \underline{m}_k] \underline{p}} = 0 , \quad (2.3)$$

where the index i is associated with the generators of the algebra $\mathfrak{h}$, while the underlined indices label a basis for the complement of $\mathfrak{h}$ in $\mathfrak{g}$. For the coset metric $ds^2 = g_{\underline{mn}} e^{\underline{m}} \otimes e^{\underline{n}}$ the relation is analogous to (2.3), with a symmetrization of indices replacing the antisymmetrization. The action of the exterior derivative preserves left-invariance, and is also determined by the structure constants of G .

None of the cosets we consider admits left-invariant 1- or 5-forms.

We define the ‘standard volume’ of the cosets as

$$I := \int e^{123456} .$$

2.1.1 $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$

Left-invariant metric:

$$g_{\underline{mn}} = \text{diag}(v^1, v^1, v^2, v^2, v^3, v^3) , \quad v^1 > 0, v^2 > 0, v^3 > 0 . \quad (2.4)$$

The left-invariant forms are spanned by

$$\begin{aligned} \omega_0 &= 1 , & \omega_1 &= -e^{12} , & \omega_2 &= e^{34} , & \omega_3 &= -e^{56} , \\ \alpha &= \frac{1}{2\sqrt{I}}(e^{135} + e^{146} - e^{236} + e^{245}) , & \beta &= \frac{1}{2\sqrt{I}}(-e^{136} + e^{145} - e^{235} - e^{246}) , \\ \tilde{\omega}^0 &= \frac{1}{I}e^{123456} , & \tilde{\omega}^1 &= \frac{1}{I}e^{3456} , & \tilde{\omega}^2 &= -\frac{1}{I}e^{1256} , & \tilde{\omega}^3 &= \frac{1}{I}e^{1234} . \end{aligned} \quad (2.5)$$

²Here and in the following (see in particular subsection 4.3), frame indices are underlined.

2.1.2 $\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$

Left-invariant metric:

$$g_{mn} = \text{diag}(v^1, v^1, v^1, v^1, v^2, v^2), \quad v^1 > 0, v^2 > 0. \quad (2.6)$$

Basis of left-invariant forms:

$$\begin{aligned} \omega_0 &= 1, & \omega_1 &= -e^{12} - e^{34}, & \omega_2 &= e^{56}, \\ \alpha &= \frac{1}{2\sqrt{I}}(e^{135} + e^{146} + e^{236} - e^{245}), & \beta &= \frac{1}{2\sqrt{I}}(e^{136} - e^{145} - e^{235} - e^{246}), \\ \tilde{\omega}^0 &= \frac{1}{I}e^{123456}, & \tilde{\omega}^1 &= \frac{1}{2I}(e^{1256} + e^{3456}), & \tilde{\omega}^2 &= -\frac{1}{I}e^{1234}. \end{aligned} \quad (2.7)$$

2.1.3 $\frac{\text{G}_2}{\text{SU}(3)}$

Left-invariant metric:

$$g_{mn} = \text{diag}(v^1, v^1, v^1, v^1, v^1, v^1), \quad v^1 > 0. \quad (2.8)$$

Basis of left-invariant forms:

$$\begin{aligned} \omega_0 &= 1, & \omega_1 &= -e^{12} + e^{34} - e^{56}, \\ \alpha &= \frac{1}{2\sqrt{I}}(e^{135} + e^{146} - e^{236} + e^{245}), & \beta &= \frac{1}{2\sqrt{I}}(-e^{136} + e^{145} - e^{235} - e^{246}), \\ \tilde{\omega}^0 &= \frac{1}{I}e^{123456}, & \tilde{\omega}^1 &= \frac{1}{3I}(e^{3456} - e^{1256} + e^{1234}). \end{aligned} \quad (2.9)$$

2.1.4 Properties

The overall factors in the basis forms (2.5), (2.7), and (2.9) have been chosen in such a way that

$$\int \langle \omega_A, \tilde{\omega}^B \rangle = \delta_A^B, \quad \int \alpha \wedge \beta = 1, \quad (2.10)$$

where $A = (0, a)$, $B = (0, b)$ and a, b label the left-invariant 2- and 4-forms. The antisymmetric pairing $\langle, \rangle$ is defined on even forms ρ, σ as $\langle \rho, \sigma \rangle = [\lambda(\rho) \wedge \sigma]_{\text{top}}$, with $\lambda(\rho_k) = (-)^{\frac{k}{2}} \rho_k$, k being the degree of ρ .

The basis forms define a closed differential system,

$$\begin{aligned} d\omega_a &= q_a \alpha, \\ d\alpha &= 0, & d\beta &= q_a \tilde{\omega}^a, \\ d\tilde{\omega}^A &= 0, \end{aligned} \quad (2.11)$$

which is also closed under the action of the Hodge star operator,

$$*\alpha = \beta, \quad *\tilde{\omega}^0 = \frac{1}{\text{Vol}}, \quad *\tilde{\omega}^a = -\frac{1}{4\text{Vol}} \mathcal{G}^{ab} \omega_b.$$

	$\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$	$\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$	$\frac{\text{G}_2}{\text{SU}(3)}$
range of a :	1, 2, 3	1, 2	1
geometric flux q_a :	$q_1 = q_2 = q_3 = -\sqrt{I}$	$q_1 = 2\sqrt{I}, q_2 = \sqrt{I}$	$q_1 = 2\sqrt{3I}$
$\mathcal{G}^{ab} =$	$\text{diag}\left(4(v^1)^2, 4(v^2)^2, 4(v^3)^2\right)$	$\text{diag}\left(2(v^1)^2, 4(v^2)^2\right)$	$\frac{4}{3}(v^1)^2$
$\text{Vol} =$	$v^1 v^2 v^3 I$	$(v^1)^2 v^2 I$	$(v^1)^3 I$
$I =$	$2^5 \pi^3$	$\frac{2^7 \pi^3}{3}$	$\frac{144 \pi^3}{5}$

Table 1: Values of the different quantities introduced in this subsection.

Here, the q_a encode what are sometimes referred to as geometric fluxes, Vol denotes the volume of the coset, and the matrix $\mathcal{G}^{ab}$ is the inverse of

$$\mathcal{G}_{ab} = \frac{1}{4\text{Vol}} \int \omega_a \wedge * \omega_b, \quad (2.12)$$

corresponding to the special Kähler metric on the space of the internal metric and B-field deformations [13]; see subsection A.1 of the appendix for more details.

In table 1, we give the values of the quantities introduced above for each coset. The standard volume I was computed following ref. [25].³ Its evaluation requires knowledge of the Euler characteristic of our cosets. Since the harmonic forms on a compact coset reside among the left-invariant forms, we can read off the cohomology from the differential relations (2.11). We immediately conclude that all our cosets have trivial odd cohomology. Concerning the even cohomology, for $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$, with

$$\omega'_1 = \omega_1 - \omega_3, \quad \omega'_2 = \omega_2 - \omega_3, \quad (2.13)$$

we have

$$H^2 = \text{Span}([\omega'_1], [\omega'_2]), \quad H^4 = \text{Span}([\tilde{\omega}^1], [\tilde{\omega}^2]),$$

hence the Euler characteristic is $\chi = 6$.

For $\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$, we have $b_2 = 1$ and $\chi = 4$, while for $\frac{\text{G}_2}{\text{SU}(3)}$, $b_2 = 0$ and $\chi = 2$.

2.2 The SU(3) structure

For each coset in (2.1), the pair of left-invariant forms parametrized by v^a ,

$$J = v^a \omega_a, \quad \Omega = 2\sqrt{\text{Vol}}(\alpha + i\beta), \quad (2.14)$$

³We have a 2^6 supplementary factor in I with respect to [25]. This is due to the fact that for the normalization of the group structure constants we follow the choice of [22], and this differs from the one of [25] by a factor 1/2.

satisfies the relations $J \wedge \Omega = 0$ and $\frac{3i}{4}\Omega \wedge \bar{\Omega} = J \wedge J \wedge J$ and hence determines a left-invariant $SU(3)$ structure. The metric specified by J and Ω is precisely the one given in eq. (2.4), (2.6), and (2.8) respectively for the three cosets. Using the properties of the basis forms listed in subsection 2.1.4 above, one can see that the differential relations (2.2) are satisfied, with torsion classes⁴

$$\begin{aligned} W_1 &= -\frac{iv^a q_a}{3\sqrt{Vol}} , \\ W_2 &= -\frac{2i}{3\sqrt{Vol}} q_a (v^a v^b - \frac{3}{4} \mathcal{G}^{ab}) \omega_b . \end{aligned} \tag{2.15}$$

Substituting the quantities given in the table of subsection 2.1.4, we see that the Nearly Kähler condition $W_2 = 0$ is identically satisfied on $\frac{G_2}{SU(3)}$. For $\frac{Sp(2)}{S(U(2) \times U(1))}$ and $\frac{SU(3)}{U(1) \times U(1)}$, this condition is satisfied on a line in the parameter space determined by $v^1 = v^2$ and $v^1 = v^2 = v^3$ respectively. In this Nearly Kähler limit the cosets are Einstein manifolds (the only other loci at which the Einstein condition is satisfied are $2v^1 = v^2$ for $\frac{Sp(2)}{S(U(2) \times U(1))}$ and $2v^1 = 2v^2 = v^3$, or cyclic permutations of this, for $\frac{SU(3)}{U(1) \times U(1)}$ [25]).

The forms (2.14) are the most general left-invariant pair satisfying the $SU(3)$ structure defining relations (the unphysical overall phase freedom in Ω has been chosen in order to make the torsion classes purely imaginary). In particular, since the volume Vol is fixed by the v^a , we see that Ω identifies a rigid $SL(3, \mathbb{C})$ structure, and there are no almost complex structure moduli.

2.3 An alternative basis?

In [13], conditions on the expansion forms were emphasized that arise when these are moduli dependent, as is the case with the basis of harmonic forms on which Calabi-Yau reductions are based (the *-ed conditions in section 2 of [13]). For the set of expansion forms that we have introduced above, these conditions are trivially satisfied, as the forms are moduli independent. In this sense, our expansion ansatz here is technically simpler than in the Calabi-Yau case. However, in a small flux approximation, the laplacian $\Delta = - * d * d - d * d *$ becomes the mass operator for the modes of the 10d supergravity fields, and an expansion in eigenforms of it is physically motivated. Can we replace the forms introduced above by such a basis of eigenforms?

In the Nearly Kähler case the expansion in eigenforms of the laplacian is further motivated by the fact that both J and Ω are themselves eigenforms of Δ [11]. In the more general case $W_2 \neq 0$, this is still true for Ω ,⁵

$$\Delta \Omega = (3|W_1|^2 + \frac{1}{4} W_2 \lrcorner \bar{W}_2) \Omega , \tag{2.16}$$

⁴The evaluation of W_2 is performed rewriting the second line of (2.2) as $W_2 = 2W_1 J - *d\Omega$.

⁵One needs the relation $dW_2 = \frac{i}{4} (W_2 \lrcorner \bar{W}_2) \text{Re } \Omega$, satisfied by the cosets (2.1).

but not for J , which instead satisfies

$$\Delta J = 3|W_1|^2 J - \frac{3}{2} \text{Re}(\bar{W}_1 W_2) .$$

Considering e.g. the coset $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$, a change of basis sending the 2-forms introduced in (2.5) to a set of eigenforms of the laplacian is

$$\omega'_1 = \omega_1 - \omega_3 \quad , \quad \omega'_2 = \omega_2 - \omega_3 \quad , \quad \omega'_3 = \frac{\sum_a (v^a)^2 \omega_a}{\sum_b (v^b)^2} , \quad (2.17)$$

where $\Delta\omega'_1 = \Delta\omega'_2 = 0$, while $\Delta\omega'_3 = \frac{(v^1)^2 + (v^2)^2 + (v^3)^2}{v^1 v^2 v^3} \omega'_3$. The harmonic 4-forms are spanned by

$$*\omega'_1 \propto \frac{v^3}{v^1} \tilde{\omega}^1 - \frac{v^1}{v^3} \tilde{\omega}^3 \quad , \quad *\omega'_2 \propto \frac{v^3}{v^2} \tilde{\omega}^2 - \frac{v^2}{v^3} \tilde{\omega}^3 , \quad (2.18)$$

while $*\omega'_3 \propto -\sqrt{I}(\tilde{\omega}^1 + \tilde{\omega}^2 + \tilde{\omega}^3) = d\beta$ is exact.

The condition $v^a \partial_{v^b} \omega_a$ (*7 of [13]) gives rise to a complicated set of equations for possible v^a dependent normalization factors of the primed basis. However, it is easy to see upon inspection that the moduli independence of the triple intersection product (condition *8 of [13]) cannot be satisfied for any such choice. The question whether the choice of left-invariant expansion forms can be motivated from a Kaluza-Klein reduction point of view hence remains an interesting open question.

3 Supersymmetric 10d solutions parametrized by fluxes

In this section, we will rewrite the family of $\mathcal{N} = 1$ solutions of the 10d supergravity equations found in [22] in a manner which makes the discreteness of this family as a result of flux quantization manifest. By [10] and [11], these solutions can be recovered from the 4d point of view. After proving the full consistency of our reduction in section 4, we will proceed to complement these solutions with their non-supersymmetric relatives in section 6.

3.1 Flux quantization and K -theory

RR-fields are classified topologically by K -theory classes. This has two consequences for the choice of fluxes associated to the RR-field strengths,. Firstly, the naive integer quantization of fluxes must be replaced by quantization in multiples of fractions determined also by the topology of the compactification manifold. Secondly, not every choice of flux number satisfying these quantization conditions will possess a K -theory lift and hence be permissible. We will now study these two points in turn.

In [32], fluxes were conjectured to take values in the image of the map

$$\sqrt{\hat{A}(X)} \text{ch}(\cdot) : K(X) \rightarrow H^{\text{even}}(X, \mathbb{Q}) .$$

$\text{ch}(x)$ is the Chern character as extended to a K -theory element $x = E - F$ via $\text{ch}(x) = \text{ch}(E) - \text{ch}(F)$. Hence,

$$\frac{[F(x)]}{2\pi} = \sqrt{\hat{A}} \text{ch}(x), \quad (3.1)$$

where $F = \sum_{i=0}^5 F_{2i}$ denotes a formal sum of all RR-field strengths, and $[\cdot]$ indicates rational cohomology class (rational rather than integral due to the fractional coefficients of Chern classes that appear in the expansion of the Chern character). When $H \neq 0$, the equations of motion and Bianchi identity of F are modified from the naive Maxwell form, enforcing harmonicity of F , to a version of these equations twisted by H . In particular, F now satisfies $(d - H)F = 0$. When H is exact, as will be the case in our study, H -twisted cohomology maps to ordinary cohomology via $F \rightarrow e^{-B}F$, where $H = dB$. It hence proves convenient to introduce a basis of RR fields given by $G = e^{-B}F$. Equation (3.1) then holds for G rather than F , and the term ‘fluxes’ refers to the cohomology classes $[G]$.

To decide which fluxes we can choose as boundary conditions of our physical system (and then parametrize our solutions by this choice), we need to decide on electric vs. magnetic variables. Ignoring subtleties related to torsion, which does not enter in a supergravity analysis, we can choose the electric basis to lie in $\oplus_{i=1}^3 H^{2i}(X, \mathbb{Q})$.

Let us now consider the question of flux quantization. To this end, we expand the right hand side of (3.1) in terms of Chern classes for x the class of a vector bundle F on X ,

$$\begin{aligned} \text{ch}_0(F) &= \text{rank}(F), & \text{ch}_1(F) &= c_1(F), & \text{ch}_2(F) &= \frac{1}{2}[c_1(F)^2 - 2c_2(F)], \\ \text{ch}_3(F) &= \frac{1}{3!}[c_1(F)^3 - 3c_1(F)c_2(F) + 3c_3(F)], \\ \hat{A} &= 1 - \frac{p_1}{24} + \dots \end{aligned}$$

Hence,

$$\begin{aligned} \frac{[G_0]}{2\pi} &= \text{rank}(F), & \frac{[G_2]}{2\pi} &= c_1(F), & \frac{[G_4]}{2\pi} &= \frac{1}{2}[c_1(F)^2 - 2c_2(F)], \\ \frac{[G_6]}{2\pi} &= \frac{1}{3!}[c_1(F)^3 - 3c_1(F)c_2(F) + 3c_3(F)] - \frac{p_1(X)}{48}c_1(F). \end{aligned}$$

As Chern classes take value in integral cohomology, it follows that in the presence of G_2 flux, $G_4/2\pi$ is generically half-integrally quantized. Neglecting gravitational effects, $G_6/2\pi$ is quantized in multiples of $\frac{1}{6}$, incorporating the $\hat{A}$ -genus generically yields quantization in multiples of $\frac{1}{48}$. In particular, for the cosets we are considering, the Pontrjagin classes are given by

$$p\left(\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}\right) = 1 \quad , \quad p\left(\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}\right) = (1+x^2)^4 \quad , \quad p\left(\frac{\text{G}_2}{\text{SU}(3)}\right) = 1. \quad (3.2)$$

	$\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$	$\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$	$\frac{\text{G}_2}{\text{SU}(3)}$
G_0	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
G_2	$\mathbb{Z}$	$\mathbb{Z}$	—
G_4	$\frac{1}{2}\mathbb{Z}$	$\frac{1}{2}\mathbb{Z}$	—
G_6	$\frac{1}{6}\mathbb{Z}$	$\frac{1}{12}\mathbb{Z}$	$\frac{1}{3}\mathbb{Z}$

Table 2: Quantization condition on fluxes.

The first result follows from a theorem of Borel and Hirzebruch, according to which the Pontrjagin class of a coset G/U , with U a maximal torus of G , is trivial. The latter two follow from the identification of the two cosets topologically with $\mathbb{CP}^3$ and S^6 respectively. The x that occurs is the generator of the integer cohomology of $\mathbb{CP}^3$. It follows that $G_6/2\pi$ is quantized in multiples of $\frac{1}{6}$ for the cosets $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$ and $\frac{\text{G}_2}{\text{SU}(3)}$, and in multiples of $\frac{1}{12}$ for $\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$. For $\frac{\text{G}_2}{\text{SU}(3)}$, we can go further. In [33], the following mod 2 relation among Chern classes is derived

$$c_3(E) = c_1(E)c_2(E) + Sq^2c_2(E) \pmod{2}.$$

Since $\frac{\text{G}_2}{\text{SU}(3)}$ has no 2- and 4-cohomology, it follows that $c_3(E)$ must be even for any vector bundle on this space ([33] provide an index theory argument for this conclusion). We conclude that on this coset, G_6 is quantized in units of $\frac{1}{3}$.

We turn to the second question raised above: given an element of $H^*(X, \mathbb{Q})$ satisfying the integrality conditions just discussed, when does it lie in the image of the map $\sqrt{\hat{A}} \text{ch}(\cdot)$, thus qualifying as a viable choice of flux? We will not provide a general answer, but address the following two subquestions which will be relevant in the next subsection.

Is it possible to have only G_0 and G_6 non-vanishing? It is a theorem (see e.g. Thm. V.3.25 in [34]) that the map (3.1) provides an isomorphism when the domain is extended to rational K -theory, $K(X) \otimes \mathbb{Q}$. It follows that any class in $H^*(X, \mathbb{Q})$ lifts to a fractional K -theory class. Multiiplying our choice of G_0 and G_6 with an appropriate integer hence always provides a viable choice of flux.

Given $G_2 = 0$, which G_4 are permissible? When G_2 vanishes, G_4 is integrally quantized. For the two cosets with non-trivial 2- and 4-cohomology, this is the only restriction on G_4 , i.e. all of $H^4(X, \mathbb{Z})$ is a permissible choice for this flux. As pointed out in [33], this situation arises whenever the cohomology of the manifold is generated in second degree. If we call the generators x_i , line bundles L_i exist with $c_1(L_i) = x_i$. The K -theory classes $x_{ij} = L_i \otimes L_j - L_i \oplus L_j$ can then be used as building blocks for lifting any G_4 -flux, by

$$\text{ch}(x_{ij}) = x_i x_j + \frac{1}{2}(x_i^2 x_j + x_i x_j^2).$$

3.2 The solution

The $\mathcal{N} = 1$ supersymmetry conditions for an AdS_4 vacuum have been determined by [35]. A nontrivial warp factor is not allowed, and the dilaton ϕ has to be constant. Furthermore, in our conventions⁶ the equations governing the H-field and the internal RR field strengths read

$$\begin{aligned} H &= \frac{2m}{5} e^\phi \text{Re } \Omega, \\ F_0 &= m, \quad F_2 = -\frac{f}{9} J + i e^{-\phi} W_2, \quad F_4 = \frac{3m}{10} J \wedge J, \quad F_6 = \frac{f}{6} J \wedge J \wedge J, \end{aligned} \quad (3.3)$$

where the only nonvanishing purely imaginary torsion classes are $W_1 = \frac{4i}{9} e^\phi f$ and W_2 . The only Bianchi identity which is not automatically satisfied is $dF_2 - H F_0 = 0$. This imposes

$$dW_2 = i e^{2\phi} \left(\frac{2}{27} f^2 - \frac{2}{5} m^2 \right) \text{Re } \Omega. \quad (3.4)$$

The AdS curvature is determined by

$$\Lambda = -3e^{2\phi} \left(\frac{m^2}{25} + \frac{f^2}{9} \right). \quad (3.5)$$

Following work of [28], [22] showed that these equations can be solved on the cosets we introduced in the previous section, by expanding all fields in forms invariant under the left group action. We will repeat this analysis, but parametrize the solutions by the fluxes $[G]$, as introduced in the previous subsection, rather than the parameter f and the dilaton. This is the favored approach as it allows us to take flux quantization into account naturally (from a 4d point of view, the distinction between fluxes and parameters such as f and the dilaton is most striking, as the former correspond to charges, the latter to VEVs; in 10d, while fluxes can also be considered as VEVs, they are distinguished by encoding topological information).

We will focus on $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$ for concreteness. This example is the most rich among the three cosets we are considering, as it has the largest set of left-invariant forms, and the largest cohomology.

The ansatz (2.14) already led to the expressions (2.15) for W_1 and W_2 in terms of the metric parameters v^a . It will prove convenient for this section to express the internal component b of the B-field using the closed 2-forms (2.13),

$$b = b^1 \omega'_1 + b^2 \omega'_2 + b^3 \omega_3.$$

⁶Our supergravity field strengths are as in [36]. We derive the susy conditions starting from an ansatz for the two type IIA susy parameters ϵ^1, ϵ^2 which assigns negative chirality to ϵ^1 and positive chirality to ϵ^2 . For the gamma matrices and the $\text{SU}(3)$ structure we adopt the conventions listed in subsection A.2 of [10]. The resulting equations (3.3) and the $\text{SU}(3)$ torsion classes differ from the ones in [22] by just a few minus signs.

Thus, b'^1 and b'^2 capture topological information about the B -field, while b'^3 enters in H . Likewise, our ansatz for G is

$$\begin{aligned} G_0 &= m, \\ G_2 &= m'^1 \omega'_1 + m'^2 \omega'_2, \\ G_4 &= -e_1 \tilde{\omega}^1 - e_2 \tilde{\omega}^2 - \tilde{\xi} d\beta, \\ G_6 &= -e \tilde{\omega}^0. \end{aligned}$$

The equations of motion for G are complicated, and are encoded in the equations (3.3). By contrast, the Bianchi identities are already guaranteed by the ansatz (hence the use of primed forms).

To solve (3.3) in terms of the flux parameters, we begin by solving (3.4) in term of ϕ , invoking the relation between W_1 and f ,

$$e^{2\phi} = \frac{5}{16m^2 v^1 v^2 v^3} [6 \sum_{a < b} v^a v^b - 5 \sum (v^a)^2].$$

For the rest of this section, ϕ will denote this solution.

Utilizing the equation for H , this allows us to solve for b'^3 in terms of the metric parameters,

$$\begin{aligned} b'^3 &= -\frac{4m}{5} \sqrt{v^1 v^2 v^3} e^\phi \\ &= -\frac{m}{|m|} \sqrt{5 \left(6 \sum_{a < b} v^a v^b - 5 \sum (v^a)^2 \right)}. \end{aligned}$$

We next want to solve for f , starting with

$$F_6 = G_6 + B \wedge G_4 + \frac{1}{2} B^2 \wedge G_2 + \frac{1}{3!} B^3 \wedge G_0 = \frac{f}{6} J \wedge J \wedge J. \quad (3.6)$$

We eliminate the B^3 term via

$$\begin{aligned} F_4 &= G_4 + B \wedge G_2 + \frac{1}{2} B \wedge B \wedge G_0 = \frac{3m}{10} J \wedge J \\ \Leftrightarrow m B^3 &= \frac{3m}{5} B \wedge J \wedge J - 2B \wedge G_4 - 2B^2 \wedge G_2. \end{aligned}$$

Hence,

$$\frac{f}{6} J \wedge J \wedge J = G_6 + \frac{2}{3} B \wedge G_4 + \frac{1}{6} B^2 \wedge G_2 + \frac{m}{10} B \wedge J \wedge J,$$

and substituting f into

$$F_2 = G_2 + B \wedge G_0 = -\frac{f}{9} J + i e^{-\phi} W_2^-$$

yields three equations which can be solved for b^1 , b^2 and $\tilde{\xi}$,

$$\begin{aligned} b^1 &= \frac{(5v^1 - 3(v^2 + v^3))\sqrt{v^1 v^2 v^3}}{4v^2 v^3 m} e^{-\phi} - \frac{m'^1}{m}, \\ b^2 &= \frac{(5v^2 - 3(v^1 + v^3))\sqrt{v^1 v^2 v^3}}{4v^1 v^3 m} e^{-\phi} - \frac{m'^2}{m} \end{aligned}$$

We omit the expression for $\tilde{\xi}$, which is lengthy and not illuminating.

At this stage, we have expressed $\tilde{\xi}, b^a, e^\phi$ in terms of v^a . Substituting these into the F_4 equation,

$$G_4 + G_2 \wedge B + \frac{1}{2} B \wedge B \wedge G_0 = \frac{3m}{10} J \wedge J, \quad (3.7)$$

yields three independent equations for v^a , two of which take the simple form

$$\begin{aligned} \frac{(v^1 - v^3)(v^1 v^2 + v^2 v^3 - 3v^1 v^3)}{v^1 v^3} - e^{2\phi} \left(\frac{m e_1}{I} + m'^2 (2m'^1 + m'^2) \right) &= 0, \\ \frac{(v^2 - v^3)(v^1 v^2 + v^1 v^3 - 3v^2 v^3)}{v^2 v^3} - e^{2\phi} \left(\frac{m e_2}{I} + m'^1 (m'^1 + 2m'^2) \right) &= 0. \end{aligned} \quad (3.8)$$

The main new feature we wish to demonstrate, as compared to the Nearly Kähler analysis of [11], is the presence of several supersymmetric vacua of a given theory, i.e. upon a fixed choice of fluxes. This phenomenon already occurs at $e_a = m'^a = 0$, which is a permissible choice of flux by the previous subsection. The third equation following from (3.7) here takes the form

$$15e^\phi \sqrt{v^1 v^2 v^3} e - 8I v^2 v^3 (v^2 v^3 - 3v^1 v^2 - 3v^1 v^3) = 0.$$

It is easy to see that this system of equations has, aside from the Nearly Kähler solution at

$$v^1 = v^2 = v^3 = -\frac{\sqrt{15}}{2} \left(\frac{1}{20I} \frac{e}{|m|} \right)^{\frac{1}{3}},$$

the solution

$$v^1 = v^2 = 2v^3 = -\frac{\sqrt{15}}{4} \left(\frac{1}{2I} \frac{e}{|m|} \right)^{\frac{1}{3}},$$

as well as two others which arise upon cyclic permutation of v^1, v^2, v^3 . Note that physical supersymmetric solutions exist only for $e < 0$.

The symmetry between the three metric parameters v^1, v^2, v^3 can be broken by considering backgrounds with G_4 flux. E.g., maintaining $G_2 = 0$, we obtain from (3.8)

$$\begin{aligned} e_1 \neq 0 &\rightarrow v^1 \neq v^3, \\ e_2 \neq 0 &\rightarrow v^2 \neq v^3, \\ e_1 \neq e_2 &\rightarrow v^1 \neq v^2. \end{aligned}$$

We have checked numerically that e.g. at $e_1 \neq 0, e_2 = 0$, solutions with $v^2 = v^3$ exist.

4 The dimensional reduction

4.1 The truncation scheme

As announced, we will adopt a reduction prescription in which the higher dimensional supergravity fields are expanded on a basis for the left-invariant tensors admitted by the coset. This expansion basis was introduced in subsect. 2.1 for the three cosets (2.1).

We stress again that this G -invariant truncation does not coincide with a massless Kaluza-Klein ansatz. We can illustrate the differences between the two schemes e.g. by considering the gauge vectors of the dimensionally reduced theory arising from the decomposition of the higher dimensional metric. The conventional massless Kaluza-Klein ansatz associates a gauge vector of the truncated theory to each Killing vector on the compact manifold, the gauge symmetry being inherited from the reparameterization invariance of the higher dimensional spacetime.⁷ On the other hand, the G -invariant ansatz preserves just a subgroup of the full isometry group of the internal manifold G/H . The theory of compact left coset spaces endowed with a left-invariant metric (such are the cosets we consider) states that in general the isometry group of G/H is $G \times N(H)/H$, where $N(H)$ is the normalizer of H in G , defined as $N(H) := \{g \in G : gH = Hg\}$. The G factor in $G \times N(H)/H$ is associated with the left action of G on the coset, while the $N(H)/H$ factor derives from the right action of G . The Killing vectors generating the right isometries are left-invariant, while this is not the case for the ones generating the left isometries.⁸ It follows that a left-invariant reduction ansatz keeps only the former, and the gauge group descending from the higher dimensional metric sector is just $N(H)/H$.

For the cosets we consider the G -invariant ansatz is particularly simple, because $N(H)/H$ turns out to be trivial. This can be seen either by observing that $\text{rank } G = \text{rank } H$ [25], or by noticing that our cosets do not admit left-invariant vectors at all. We conclude that no gauge vectors will descend from the dimensional reduction of the type II supergravity NSNS sector, and the whole (abelian) gauge group will be provided by the RR sector. This is analogous to what is realized in Calabi-Yau compactifications.

Though physically well motivated, dimensional reductions based on the full massless KK ansatz have a drawback: they are generically inconsistent [38, 12]. Rare exceptions are known, an example being the S^7 reduction of [39] (see [40] for a discussion of consistent KK sphere reductions). The G -invariant reduction scheme is instead believed to provide consistent truncations, due to the fact that the preserved invariant fields never generate the truncated non-invariant modes. A further argument for consistency is that the substitution of a G -invariant ansatz guarantees the dropping of the dependence on the internal coordinates y from the higher dimensional Lagrangian, see e.g. [41, 12] for more details. The consistency of the G -invariant scheme was explicitly shown in ref. [42] for a reduction

⁷In principle, nonvanishing background values of the non-metric supergravity fields may break the gauge symmetry to a subgroup of the isometry group, however this is guaranteed not to happen as far as these vevs are invariant under the isometries [12, pag. 16].

⁸A detailed discussion of the isometries of G/H can be found in section 2 of ref. [37].

of the pure gravity action. Recent related discussions can be found in [43] (for coset space reductions of Einstein-Yang-Mills theories), in [44, 45] (for Scherk-Schwarz reductions on group manifolds), and in [46, 47] (for consistent reductions on spaces supporting AdS solutions, and their relation with the dual SCFT). However, an explicit check of consistency in the context of $SU(3)$ structure compactifications with fluxes had not been performed to date. In subsection 4.3 we will work out the reduction of the higher dimensional equations of motion in detail, and prove the consistency of the truncation of the full type IIA bosonic sector for the cosets (2.1).

4.2 The 4d action

Following the reduction prescription for type IIA on $SU(3)$ structure manifolds initiated in [5], the complete 4d gauged $\mathcal{N} = 2$ bosonic action has by now been derived [48, 49, 50, 27, 13]. Here, we will use the notation of ref. [10]. Separating the contributions of the NSNS and RR sectors, the action $S^{(4)}$ arising from a reduction on our cosets (2.1) reads $S^{(4)} = S_{\text{NS}}^{(4)} + S_{\text{RR}}^{(4)}$, with

$$S_{\text{NS}}^{(4)} = \int_{M_4} \left(\frac{1}{2} R_4 * 1 - \frac{1}{4} e^{-4\varphi} dB \wedge *dB - d\varphi \wedge *d\varphi - \mathcal{G}_{ab} dt^a \wedge *dt^b - V_{\text{NS}} * 1 \right), \quad (4.1)$$

$$S_{\text{RR}}^{(4)} = \int_{M_4} \left\{ \frac{1}{4} \text{Im} \mathcal{N}_{AB} F^A \wedge *F^B + \frac{1}{4} \text{Re} \mathcal{N}_{AB} F^A \wedge F^B - \frac{e^{2\varphi}}{4} (D\xi \wedge *D\xi + d\tilde{\xi} \wedge *d\tilde{\xi}) \right. \\ \left. + \frac{1}{4} dB \wedge [\xi d\tilde{\xi} - \tilde{\xi} D\xi + 2e_A A^A + \tilde{\xi} q_a A^a] - \frac{1}{4} m^A e_A B \wedge B - V_{\text{RR}} * 1 \right\}. \quad (4.2)$$

The different quantities appearing in this 4d action are introduced in appendix A, where we also give some details about the derivation from the higher dimensional supergravity. The 4d degrees of freedom descending from the NSNS sector are the metric $g_{\mu\nu}$, the 2-form B , the complex scalars $t^a = b^a + iv^a$ and the 4d dilaton φ , defined in (A.2). The RR sector yields the scalars ξ and $\tilde{\xi}$ introduced in the first line of (A.12), as well as the gauge potentials A^A , whose modified field strengths F^A are defined in (A.13) (recall that the index A runs over $(0, a)$).

The $\mathcal{N} = 2$ action $S^{(4)}$ contains the gravitational multiplet $(g_{\mu\nu}, A^0)$, a number of vector multiplets (t^a, A^a) (see table 1 for the coset dependent range of a), and one tensor multiplet $(B, \varphi, \xi, \tilde{\xi})$. When $m^A = 0$ the antisymmetric tensor B becomes massless and can be dualized to a scalar, yielding the universal hypermultiplet. From $D\xi = d\xi - q_a A^a$ it follows that ξ is charged under the A^a , the charges being provided by the geometric fluxes q_a given in table 1. The graviphoton A^0 instead does not participate to this gauging (due to the fact that the compactification manifolds (2.1) don't allow for a flux of the NSNS 3-form [5]).

The special Kähler metric $\mathcal{G}_{ab}$ governing the kinetic terms for the scalars in the vector multiplets is given in table 1, and further discussed in subsection A.1 of the appendix,

together with the period matrix $\mathcal{N}_{AB}$ describing the kinetic and topological terms for the gauge potentials.

The full 4d scalar potential reads $V = V_{\text{NS}} + V_{\text{RR}}$. Reduction of the internal NSNS sector on our coset spaces yields⁹

$$\begin{aligned} V_{\text{NS}} &\equiv -\frac{e^{2\varphi}}{2} \left(R_6 - \frac{1}{2} H \lrcorner H \right) \\ &= \frac{e^{2\varphi}}{4\text{Vol}} q_a q_b \left(\mathcal{G}^{ab} - 3v^a v^b + b^a b^b \right), \end{aligned} \quad (4.3)$$

where the 6d Ricci scalar R_6 has been evaluated in terms of the torsion classes expressed in eq. (2.15) via the formula¹⁰ [51]

$$R_6 = \frac{15}{2} |W_1|^2 - \frac{1}{2} W_2 \lrcorner \overline{W}_2, \quad (4.4)$$

while for the internal NSNS 3-form we have $H = d_6 b = b^a q_a \alpha$.

The RR contribution to the scalar potential, obtained from the general expression given in eq. (A.14) of the appendix, is

$$V_{\text{RR}} = -\frac{e^{4\varphi}}{4} \left[m^A \text{Im} \mathcal{N}_{AB} m^B + (e_A + q_A \tilde{\xi} - m^C \text{Re} \mathcal{N}_{CA}) (\text{Im} \mathcal{N})^{-1 AB} (e_B + q_B \tilde{\xi} - \text{Re} \mathcal{N}_{BD} m^D) \right], \quad (4.5)$$

where $q_A = (0, q_a)$. Notice that while $\tilde{\xi}$ appears in the potential, the other RR scalar ξ is a flat direction. Since the matrix $\text{Im} \mathcal{N}$ is negative, V_{RR} is positive semi-definite.

4.3 Consistency of the truncation

We now prove the consistency of the dimensional reduction leading to the 4d action $S^{(4)}$ introduced in the previous subsection. To this end, we plug the G -invariant reduction ansatz into the bosonic equations of motion (EoM) of type IIA supergravity, and show that these yield the EoM following from the reduced action $S^{(4)}$.

The reduction of the equations for the RR degrees of freedom was already described in the general analysis of [10] and is summarized, for the specific compactification on the coset spaces (2.1), in subsection A.2 of the appendix. In fact, the piece (4.2) of the 4d action has been established requiring its compatibility with the EoM for the 4d fields $A^A, \xi, \tilde{\xi}$ as obtained from the higher dimensional equations (A.9), (A.10). It follows that, as far the RR sector is concerned, the reduction is consistent by construction.

⁹For any pair of forms P, Q of degree k we define the contraction $P \lrcorner Q := \frac{1}{k!} P_{m_1 \dots m_k} Q^{m_1 \dots m_k}$, so that $(P \lrcorner Q) * 1 = P \wedge * Q$. This also holds for the 10d spacetime equations of the forthcoming subsection.

¹⁰An equivalent expression for R_6 was given in [25] using a general formula relating the Riemann tensor of G/H to the structure constants of G . The 4 factor mismatch we have w.r.t. that expression is due to the different normalization of the $\text{SU}(3)$ structure constants already mentioned in footnote 3.

Hence, we just have to analyse the equations of motion for the NSNS degrees of freedom, namely the B -field, the Einstein and the dilaton equations. For the democratic formulation of type IIA supergravity [36] in string frame, these read

$$d(e^{-2\phi} * \hat{H}) - \frac{1}{2}[\hat{\mathbf{F}} \wedge * \hat{\mathbf{F}}]_8 = 0, \quad (4.6)$$

$$\hat{R}_{MN} + 2\hat{\nabla}_M \partial_N \phi - \frac{1}{2}\iota_M \hat{H} \lrcorner \iota_N \hat{H} - \frac{e^{2\phi}}{4} \sum_{k=0}^{10} \iota_M \hat{F}_{(k)} \lrcorner \iota_N \hat{F}_{(k)} = 0, \quad (4.7)$$

$$\hat{R} - \frac{1}{2}\hat{H} \lrcorner \hat{H} + 4(\hat{\nabla}^2 \phi - \partial_M \phi \hat{\partial}^M \phi) = 0, \quad (4.8)$$

where the hat denotes 10d quantities, $\hat{\mathbf{F}} \equiv \sum_{k=0}^{10} \hat{F}_{(k)}$ is the sum of the RR field-strengths, and M, N are 10d spacetime indices.

$\hat{B}$ -field EoM

The $\hat{B}$ -field EoM (4.6) is an 8-form equation. Its expansion in the left-invariant forms on M_6 yields two independent equations: the first exhibiting two indices along 4d spacetime M_4 and 6 indices along M_6 , and the second with 4 indices along M_4 and 4 indices along M_6 . We get no equation with 5 indices along M_6 due to the absence of invariant 5-forms on the cosets (2.1). Concretely, recalling (A.10) we rewrite the RR piece of (4.6) as

$$[\hat{\mathbf{F}} \wedge * \hat{\mathbf{F}}]_8 = [\hat{\mathbf{F}} \wedge \lambda(\hat{\mathbf{F}})]_8 = [\hat{\mathbf{G}} \wedge \lambda(\hat{\mathbf{G}})]_8.$$

Expanding $\hat{B}$ as in (A.4) and $\hat{\mathbf{G}}$ as in (A.11), we see that eq. (4.6) reduces to

$$\left[d(e^{-4\varphi} * dB) + G_{(0)}^A \tilde{G}_{(2)A} - \tilde{G}_{(0)A} G_{(2)}^A + \tilde{G}_{(1)} \wedge G_{(1)} \right] \tilde{\omega}^0 = 0 \quad (4.9)$$

and

$$\begin{aligned} & -4d_4(\mathcal{G}_{ab} *_4 d_4 b^b) \tilde{\omega}^a + e^{-2\phi+4\varphi} \text{vol}_4 \wedge d_6(*_6 d_6 b) + \\ & + \left[G_{(0)}^0 \tilde{G}_{(4)a} + G_{(4)}^0 \tilde{G}_{(0)a} - \mathcal{K}_{abc} G_{(0)}^b G_{(4)}^c - G_{(2)}^0 \wedge \tilde{G}_{(2)a} + \frac{1}{2} \mathcal{K}_{abc} G_{(2)}^b \wedge G_{(2)}^c \right] \tilde{\omega}^a = 0, \end{aligned} \quad (4.10)$$

where the 4d forms $G_{(p)}, \tilde{G}_{(p)}$ are expressed in (A.12), and we used $\omega_a \wedge \omega_b = -\mathcal{K}_{abc} \tilde{\omega}^c$, with the $\mathcal{K}_{abc}$ given in (A.6).

Eq. (4.9) provides the EoM for the 2-form B in 4d. It already appeared in section 5 of ref. [10], where it was employed in order to deduce the 4d action $S^{(4)}$ written in subsection 4.2 above. It follows that, on the same footing as the RR equations, consistency of this equation with the action $S^{(4)}$ is guaranteed by construction.

Eq. (4.10) (which was not analysed in [10]) corresponds to the EoM for the 4d scalars b^a defined by the expansion of the internal B-field b on the basis 2-forms. Using $d_6 *_6 d_6 b =$

$q_b b^b q_a \tilde{\omega}^a$, substituting the expressions (A.12) for $G_{(2)}, \tilde{G}_{(2)}, G_{(4)}, \tilde{G}_{(4)}$ and the definition (A.13) of F^A , eq. (4.10) reads

$$4\nabla_\mu (\mathcal{G}_{ab} \partial^\mu b^b) - e^{2\varphi} \frac{q_b b^b q_a}{Vol} - \text{Im} \mathcal{N}_{aB} * (F^0 \wedge * F^B) - \text{Re} \mathcal{N}_{aB} * (F^0 \wedge F^B) \\ + \frac{1}{2} \mathcal{K}_{abc} * (F^b \wedge F^c) + e^{4\varphi} [G_{(0)}^0 (\text{Im} \mathcal{N} G_{(0)} - \text{Re} \mathcal{N} L)_a - \tilde{G}_{(0)a} L^0 + \mathcal{K}_{abc} G_{(0)}^b L^c] = 0,$$

where we denote $L \equiv (\text{Im} \mathcal{N})^{-1} (\tilde{G}_{(0)} - \text{Re} \mathcal{N} G_{(0)})$. Recalling the form of $\text{Im} \mathcal{N}$ and $\text{Re} \mathcal{N}$ in (A.7) and (A.8), as well as V_{NS} in (4.3) and V_{RR} in (A.14), one checks that this equation can be reformulated as

$$2\nabla_\mu (\mathcal{G}_{ab} \partial^\mu b^b) - \frac{1}{4} \partial_{b^a} \text{Im} \mathcal{N}_{BC} * (F^B \wedge * F^C) - \frac{1}{4} \partial_{b^a} \text{Re} \mathcal{N}_{BC} * (F^B \wedge F^C) - \partial_{b^a} (V_{\text{NS}} + V_{\text{RR}}) = 0$$

which is precisely the EoM obtained varying $S^{(4)}$ in (4.1), (4.2) with respect to b^a .

10d Einstein equation

We first deal with the term $\hat{R}_{MN} + 2\hat{\nabla}_M \partial_N \phi$ in eq. (4.7). Starting from the G -invariant metric ansatz (A.1) and recalling that the 4d dilaton $\varphi(x)$ satisfies (A.3), we derive the following decomposition¹¹

$$\begin{aligned} \hat{R}_{\mu\nu} + 2\hat{\nabla}_\mu \partial_\nu \phi &= R_{\mu\nu} - \frac{1}{4} g^{mp} g^{nq} \partial_\mu g_{mn} \partial_\nu g_{pq} - 2\partial_\mu \varphi \partial_\nu \varphi - g_{\mu\nu} \nabla_4^2 \varphi, \\ \hat{R}_{\mu n} &= 0 = \hat{\nabla}_\mu \partial_n \phi, \\ \hat{R}_{mn} + 2\hat{\nabla}_m \partial_n \phi &= R_{mn} + \frac{1}{2} e^{-2\varphi} (g^{pq} \partial_\mu g_{mp} \partial^\mu g_{nq} - \nabla_4^2 g_{mn}). \end{aligned} \quad (4.11)$$

Taking the trace, we get

$$\hat{R} + 4\hat{\nabla}^2 \phi - 4\partial_M \phi \hat{\partial}^M \phi = e^{-2\varphi} (R_4 + e^{2\varphi} R_6 - \frac{1}{4} g^{mp} g^{nq} \partial_\mu g_{mn} \partial^\mu g_{pq} - 2\nabla_4^2 \varphi - 2\partial_\mu \varphi \partial^\mu \varphi). \quad (4.12)$$

In the previous expressions, quantities labeled with 4 or 6 are associated to $(M_4, g_{\mu\nu})$ or (M_6, g_{mn}) respectively. The 4d indices on the r.h.s. are raised using the rescaled metric $g^{\mu\nu}$ of eq. (A.1). Notice that all the terms depend just on x^μ : indeed, thanks to G -invariance, the whole dependence on the internal coordinates drops out.

Let's now consider the $\mu\nu$ components of the 10d Einstein equation (4.7). Using (4.11), (4.12) we find (we reinstate in the Einstein equation the term proportional to $\hat{g}_{\mu\nu}$, which

¹¹The nonvanishing higher dimensional Christoffel symbols are:

$$\hat{\Gamma}_{\mu\nu}^\rho = \Gamma_{\mu\nu}^\rho + \partial_\mu \varphi \delta_\nu^\rho + \partial_\nu \varphi \delta_\mu^\rho - g_{\mu\nu} \partial^\rho \varphi, \quad \hat{\Gamma}_{mn}^\rho = -\frac{1}{2} e^{-2\varphi} \partial^\rho g_{mn}, \quad \hat{\Gamma}_{\mu n}^p = \frac{1}{2} g^{pq} \partial_\mu g_{nq}, \quad \hat{\Gamma}_{mn}^p = \Gamma_{mn}^p.$$

In the derivation of $\hat{R}_{\mu n} = 0$ we assume $\nabla_m e_n^p = 0$.

actually vanishes thanks to the dilaton EoM (4.8)),

$$\begin{aligned}
\hat{R}_{\mu\nu} + 2\hat{\nabla}_\mu\partial_\nu\phi - \frac{1}{2}\iota_\mu\hat{H}\lrcorner\iota_\nu\hat{H} - \frac{1}{2}\hat{g}_{\mu\nu}\left(\hat{R} + 4\hat{\nabla}^2\phi - 4\partial_\rho\phi\hat{\partial}^\rho\phi - \frac{1}{2}\hat{H}^2\right) = \\
= R_{\mu\nu} - \frac{1}{4}e^{-4\varphi}H_{\mu\nu\rho}H_\nu{}^{\rho\sigma} - 2\partial_\mu\varphi\partial_\nu\varphi - 2\mathcal{G}_{ab}\partial_{(\mu}t^a\partial_{\nu)}\bar{t}^b \\
- g_{\mu\nu}\left(\frac{1}{2}R_4 - \frac{1}{24}e^{-4\varphi}H_{\mu\nu\rho}H^{\mu\nu\rho} - \partial_\mu\varphi\partial^\mu\varphi - \mathcal{G}_{ab}\partial_\mu t^a\partial^\mu\bar{t}^b - V_{\text{NS}}\right).
\end{aligned} \tag{4.13}$$

For the RR piece, taking into account all the terms of the expansion described in subsection A.2 of the appendix, we arrive at

$$\begin{aligned}
-\frac{e^{2\phi}}{4}\sum_{k=0}^{10}\iota_\mu\hat{F}_{(k)}\lrcorner\iota_\nu\hat{F}_{(k)} &= \frac{1}{2}\text{Im}\mathcal{N}_{AB}\iota_\mu F^A\lrcorner\iota_\nu F^B - \frac{1}{2}e^{2\varphi}(D_\mu\xi D_\nu\xi + \partial_\mu\tilde{\xi}\partial_\nu\tilde{\xi}) \\
&- g_{\mu\nu}\left\{\frac{1}{4}\text{Im}\mathcal{N}_{AB}F^A\lrcorner F^B - \frac{e^{2\varphi}}{4}[(D_\mu\xi)^2 + (\partial_\mu\tilde{\xi})^2] - V_{\text{RR}}\right\}.
\end{aligned} \tag{4.14}$$

From (4.13), (4.14) we see that the equation arising from the $\mu\nu$ components of (4.7) precisely reproduces the 4d Einstein equation following from $S^{(4)}$.

Since there are no left-invariant 1-forms on the cosets (2.1), the 10d Einstein equation with μn indices is trivialized by our left-invariant truncation prescription, and does not yield any constraint at the 4d level. Indeed, one can check that all the μn terms in (4.7) vanish once the truncation ansatz is plugged in.

Finally, we study the purely internal components of (4.7) in flat $\underline{mn}$ indices. Depending on which of the cosets (2.1) we consider, these yield just one, two or three 4d scalar equations, labeled by the index a . On our cosets, any left-invariant symmetric rank-2 tensor has the same diagonal structure as the invariant metric $g_{\underline{mn}}$ given in subsection 2.1. Furthermore, the left-invariant Ricci tensor on coset spaces satisfies $R_{\underline{mn}} = \frac{\partial}{\partial g^{\underline{mn}}}R_6$. Focusing for definiteness on $\frac{\text{SU}(3)}{\text{U}(1)\times\text{U}(1)}$, we have (recall $\mathcal{G}^{ab}$ in table 1)

$$R_{\underline{2a-1}\underline{2a-1}} \equiv R_{\underline{2a}\underline{2a}} = -\frac{1}{8}\mathcal{G}^{ab}\partial_{v^b}R_6, \quad a = 1, 2, 3.$$

Then, using the last line of (4.11), we get

$$\hat{R}_{\underline{2a}\underline{2a}} + 2\hat{\nabla}_{\underline{2a}}\partial_{\underline{2a}}\phi - \frac{1}{2}\iota_{\underline{2a}}\hat{H}\lrcorner\iota_{\underline{2a}}\hat{H} = \frac{e^{-2\varphi}\mathcal{G}^{ab}}{4}\left[-2\nabla_\mu(\mathcal{G}_{bc}\partial^\mu v^c) + \partial_{v^b}\mathcal{G}_{cd}\partial_\mu t^c\partial^\mu\bar{t}^d + \partial_{v^b}V_{\text{NS}}\right]. \tag{4.15}$$

Concerning the RR term, a tedious computation gives

$$-\frac{e^{2\phi}}{4}\sum_{k=0}^{10}\iota_{\underline{2a}}\hat{F}_{(k)}\lrcorner\iota_{\underline{2a}}\hat{F}_{(k)} = \frac{e^{-2\varphi}\mathcal{G}^{ab}}{4}\left[\partial_{v^b}V_{\text{RR}} - \frac{1}{4}\partial_{v^b}(\text{Im}\mathcal{N}_{CD})F^C\lrcorner F^D\right]. \tag{4.16}$$

Analogous steps can be repeated for the cosets $\frac{\text{Sp}(2)}{\text{S}(\text{U}(2)\times\text{U}(1))}$ and $\frac{\text{G}_2}{\text{SU}(3)}$, leading to the same r.h.s. of the above equations.

From (4.15), (4.16) we conclude that the components of the 10d Einstein equation (4.7) with two internal indices precisely match the EoM for the scalars v^a following from $S^{(4)}$:

$$- 2\nabla_\mu(\mathcal{G}_{ab}\partial^\mu v^b) + \partial_{v^a}\mathcal{G}_{bc}\partial_\mu t^b\partial^\mu \bar{t}^c + \partial_{v^a}(V_{\text{NS}} + V_{\text{RR}}) - \frac{1}{4}\partial_{v^a}(\text{Im}\mathcal{N}_{BC})F^B \lrcorner F^C = 0 .$$

Dilaton equation

Subtracting the trace over the $\mu\nu$ components of (4.7) from the 10d dilaton equation (4.8), we eventually obtain

$$2\nabla_4^2\varphi + \frac{1}{6}e^{-4\varphi}H_{\mu\nu\rho}H^{\mu\nu\rho} - \frac{e^{2\varphi}}{2}[(D_\mu\xi)^2 + (\partial_\mu\tilde{\xi})^2] - 2V_{\text{NS}} - 4V_{\text{RR}} = 0 , \quad (4.17)$$

which is the EoM for the 4d dilaton φ following from $S^{(4)}$.

This concludes the consistency proof of the dimensional reduction.

5 The 4d potential via $\mathcal{N} = 2$

In this section, we recast the scalar potential obtained in (4.3) and (4.5) in 4d $\mathcal{N} = 2$ language. In this framework, given the prepotential $\mathcal{F}$ governing the special geometry data of the vector multiplet sector and the quaternionic metric h_{uv} of the hypermultiplet sector, the potential is uniquely determined by the gauged isometries of h_{uv} . This structure allows us to incorporate string loops into our considerations, which correct the hypermultiplet metric. As the 4 dimensional quaternionic metrics with the isometry structure imposed by our compactifications are highly constrained, we use the results of [57, 59] to write down the general form of the all-loop string corrected potential in subsection 5.2. We analyse this potential further in subsection 6.3.

The general form of the potential in 4d $\mathcal{N} = 2$ gauged supergravity is [9, 52, 53, 54]

$$V = 4e^K h_{uv}(X^A k_A^u - \tilde{k}^{uA}\mathcal{F}_A)(\bar{X}^B k_B^u - \tilde{k}^{uB}\bar{\mathcal{F}}_B) - \left[\frac{1}{2}(\text{Im}\mathcal{N})^{-1AB} + 4e^K X^A \bar{X}^B \right] (\mathcal{P}_A^x - \tilde{\mathcal{P}}^{Cx}\mathcal{N}_{CA})(\mathcal{P}_B^x - \tilde{\mathcal{P}}^{Dx}\mathcal{N}_{DB}) . \quad (5.1)$$

The coordinates X , the prepotential $\mathcal{F}$, and the gauge coupling matrix $\mathcal{N}$ encode special geometry data and are discussed further in appendix A. h_{uv} refers to the universal hypermultiplet metric, which is expressed in terms of the quaternionic vielbein components as

$$h = u \otimes \bar{u} + v \otimes \bar{v} .$$

We will denote the quaternionic coordinates collectively by q^u . k_A^u and $\tilde{k}^{uA}$ are the components of the Killing vectors describing the isometries of the hypermultiplet metric being gauged by the A^{th} gauge vector. The $\text{Sp}(1)$ factor ω of the spin connection of the hypermultiplet metric enters in the potential via its relation to the Killing prepotentials. For

the case that the 3 components of the curvature of ω each are invariant under an isometry $k^u \partial_{q^u}$ of the metric, the corresponding Killing prepotential is given by [52, 55]

$$\mathcal{P}^x = \omega_u^x k^u.$$

In this case, one can rewrite the potential in a more convenient form. Introducing

$$Q_A^u = k_A^u - \tilde{k}^{uB} \mathcal{N}_{BA},$$

we obtain

$$V = Q_A^u \bar{Q}_B^v \left[4e^K X^A \bar{X}^B (u \otimes \bar{u} + v \otimes \bar{v})_{uv} - \left(4e^K X^A \bar{X}^B + \frac{1}{2} (\text{Im } \mathcal{N})^{-1 AB} \right) \sum_x (\omega^x \otimes \omega^x)_{uv} \right]. \quad (5.2)$$

5.1 Tree level

At tree level, the quaternionic vielbein is given by [56]¹²

$$\begin{aligned} u &= \frac{1}{2} e^\varphi (d\tilde{\xi} - id\xi), \\ v &= d\varphi - i \frac{e^{2\varphi}}{2} (da + \tilde{\xi} d\xi). \end{aligned}$$

The $\text{Sp}(1)$ connection has the following form in terms of these quaternionic vielbein components¹³

$$\omega^1 = i(\bar{u} - u) \quad , \quad \omega^2 = -(u + \bar{u}) \quad , \quad \omega^3 = \frac{i}{2}(v - \bar{v}).$$

In the class of theories we are considering, the isometries being gauged are described by the following Killing vectors

$$\begin{aligned} k_A &= \sqrt{2} \left(e_A \frac{\partial}{\partial a} + q_A \frac{\partial}{\partial \xi} \right), \\ \tilde{k}^A &= \sqrt{2} m^A \frac{\partial}{\partial a}. \end{aligned}$$

Since Q^u does not contain a non-vanishing entry for $u = \varphi$, the real part of v does not enter upon contraction with Q^u , hence we can substitute

$$\sum_x (\omega^x \otimes \omega^x) \sim 4u \otimes \bar{u} + v \otimes \bar{v}$$

¹² $\varphi, \xi, \tilde{\xi}$ were introduced above. The coordinate a is related to the dual a_B of the spacetime component of the B-field via $a_B = a + \frac{\xi \tilde{\xi}}{2}$.

¹³The components ω^x of the $\text{Sp}(1)$ curvature ω should not be confused with the expansion forms ω_a .

in the potential, obtaining

$$V = Q_A^u \bar{Q}_B^v \left[-e^{2\varphi} \left(\frac{1}{2} (\text{Im } \mathcal{N})^{-1AB} + 3e^K X^A \bar{X}^B \right) (d\xi^2 + d\tilde{\xi}^2)_{uv} - \frac{1}{8} e^{4\varphi} (\text{Im } \mathcal{N})^{-1AB} (da + \tilde{\xi} d\xi)_{uv}^2 \right].$$

This coincides with (4.3) and (4.5) obtained above via reduction from 10 dimensions.

5.2 All string loop

For the case of the universal hypermultiplet with 3 isometries, the quaternionic metric is of the Calderbank-Pedersen form [57], determined by a single function $\sqrt{\rho} F(\rho, \eta) = \rho^2 - c$ [58, 59], such that

$$\begin{aligned} u &= \frac{\sqrt{\rho^2 + c}}{2(\rho^2 - c)} (d\tilde{\xi} - i d\xi), \\ v &= \frac{\rho}{2(\rho^2 - c) \sqrt{\rho^2 + c}} \left[2 \frac{\rho^2 + c}{\rho} d\rho + i(da + \tilde{\xi} d\xi) \right]. \end{aligned} \quad (5.3)$$

This class of metrics hence comes in a 1-parameter family. The metric at string tree level lies at $c = 0$, and the variable identification

$$\rho = e^{-\varphi}$$

takes up back to the expression for the metric introduced above.¹⁴

In terms of the quaternionic vielbein components (5.3), the $\text{Sp}(1)$ connection of the Calderbank-Pedersen metric is [57]

$$\begin{aligned} \omega^1 &= \frac{\rho}{\sqrt{\rho^2 + c}} i(\bar{u} - u) = -\frac{\rho}{\rho^2 - c} d\xi, \\ \omega^2 &= -\frac{\rho}{\sqrt{\rho^2 + c}} (u + \bar{u}) = -\frac{\rho}{\rho^2 - c} d\tilde{\xi}, \\ \omega^3 &= \frac{\sqrt{\rho^2 + c}}{\rho} \frac{i}{2} (v - \bar{v}) = -\frac{1}{2(\rho^2 - c)} (da + \tilde{\xi} d\xi). \end{aligned}$$

The $\mathcal{N} = 2$ potential (5.2) for this choice of metric becomes

$$\begin{aligned} V &= \frac{Q_A^u \bar{Q}_B^v}{(\rho^2 - c)^2} \left[\left(-\frac{1}{2} (\text{Im } \mathcal{N})^{-1AB} - 3e^K X^A \bar{X}^B \right) \rho^2 (d\xi^2 + d\tilde{\xi}^2)_{uv} \right. \\ &\quad - \frac{1}{8} (\text{Im } \mathcal{N})^{-1AB} (da + \tilde{\xi} d\xi)_{uv}^2 + c e^K X^A \bar{X}^B (d\xi^2 + d\tilde{\xi}^2)_{uv} \\ &\quad \left. - \frac{c}{\rho^2 + c} e^K X^A \bar{X}^B (da + \tilde{\xi} d\xi)_{uv}^2 \right]. \end{aligned} \quad (5.4)$$

¹⁴The coordinates used in [59] are related to our choice via $\psi = \frac{a+\xi\tilde{\xi}}{2}$, $\eta = -\frac{\xi}{2}$, $\phi = \tilde{\xi}$.

In the case of Calabi-Yau compactifications, the metric is corrected away from $c = 0$ in passing from tree level to one-loop [59]. Beyond 1-loop, all corrections can be captured by field redefinitions. This means that the quaternionic metric (i.e. the value of c) remains unchanged, the identification $\rho = e^{-\varphi}$ however is modified (note that the isometry structure of the metric determines the identification of the other 3 Calderbank-Pedersen coordinates with the 10d variables as indicated in footnote 14; this is why we have not introduced separate notation for them).

To study perturbative string corrections in the case of interest, let us review the argument of [59]. The 1-loop correction to the four dimensional Einstein-Hilbert term can be determined by reduction of the 1-loop R^4 correction in 10d.¹⁵ In the normalization of [59], this yields

$$S_{\text{Einstein-Hilbert}} = \int d^4x \sqrt{g} \left(e^{-2\phi} - \frac{4\zeta(2)\chi}{(2\pi)^3} \right) R.$$

Unfortunately, the full 1-loop corrected 10d action is not available as a means towards obtaining the 1-loop completion of the 4d action. Nonetheless, after parametrizing the ignorance regarding this action and comparing to the 4d effective action obtained by choosing the Calderbank-Pedersen metric on the universal hypermultiplet scalar manifold, [59] finds that only two possible values for c are possible,

$$c = 0 \quad \text{or} \quad c = -\frac{4\zeta(2)\chi}{(2\pi)^3},$$

with χ the Euler characteristic of the Calabi-Yau. A perturbative string calculation then establishes that it is the latter value that is correct beyond tree level. Such a calculation in the case of the coset backgrounds with RR-flux that we are interested in is very challenging, and beyond the scope of this work. However, the first part of the analysis of [59] goes through also for these more general backgrounds. In particular, the 10d R^4 term is proportional to [59]

$$t_8 t_8 R^4 + \frac{1}{4} E_8.$$

The first term is shorthand for $t_8 t_8 R^4 = t^{M_1 \dots M_8} t^{N_1 \dots N_8} R_{M_1 M_2 N_1 N_2} \dots R_{M_7 M_8 N_7 N_8}$, which is expanded in terms of scalars built out of contractions of four Riemann tensors in eq. (A.12) of [59]. The second term can be written compactly in form notation as

$$E_8 \sim \Omega_{AB} \wedge \Omega_{CD} \wedge \Omega_{EF} \wedge \Omega_{GH} \wedge *(e^A \wedge \dots \wedge e^H),$$

with $\Omega^A{}_B = \frac{1}{2} R^A{}_{BCD} e^C e^D$ the curvature 2-form and e^A , $A = 1, \dots, 10$ a local coframe basis. From the expansion of the t_8 term in [59], we see that in each scalar invariant,

¹⁵As with all such arguments, we are relying on the off-shell continuation of an on-shell string computation. It would be desirable to back this line of reasoning up with an explicit string computation on the background in question. We thank Pierre Vanhove for discussions on this point.

contractions pair at least two Riemann tensors. Hence, this term does not contribute to the 4d Einstein-Hilbert term upon reduction. The contribution from E_8 to the Einstein-Hilbert term stems, exactly as in the Ricci flat case, from

$$\Omega_{ab} \wedge *_4(e^a \wedge e^b) \wedge \Omega_{mn} \wedge \Omega_{pq} \wedge \Omega_{rs} \wedge *_6(e^m \wedge \cdots \wedge e^s),$$

with a, b flat spacetime and $m, n, \dots$ flat internal indices. We recognize the internal contribution as proportional to the 6 dimensional Euler density. The conclusion of our analysis is hence that in generalizing beyond Calabi-Yau manifolds, the same two possibilities for the Calderbank-Pedersen parameter c exist as in the Calabi-Yau case (and await a perturbative string calculation as arbiter).

6 Non-supersymmetric vacua

As an application of our consistent truncation result, we will search for non-supersymmetric vacua of the 4d effective action. By the analysis of section 4, these are guaranteed to lift to 10d solutions.

6.1 Tree level

The potential we obtained at tree level above has the form

$$V = A_1 e^{2\varphi} + A_2 e^{4\varphi}, \quad (6.1)$$

with

$$\begin{aligned} A_1 &= -Q_A^u \bar{Q}_B^v \left(\frac{1}{2} (\text{Im } \mathcal{N})^{-1AB} + 3e^K X^A \bar{X}^B \right) (d\xi^2 + d\tilde{\xi}^2)_{uv}, \\ A_2 &= -Q_A^u \bar{Q}_B^v \frac{1}{8} (\text{Im } \mathcal{N})^{-1AB} (da + \tilde{\xi} d\xi)_{uv}^2. \end{aligned} \quad (6.2)$$

Minimizing the potential with regard to the 4d dilaton yields [60]

$$V_\varphi = -\frac{A_1^2}{4A_2}.$$

As A_2 is positive definite, the potential at tree level is negative semi-definite on-shell. In fact, this result generalizes immediately to any hypermultiplet metric of the general form [56] that arises upon Calabi-Yau and $\text{SU}(3)$ structure compactifications, and the respective gaugings. The corresponding potential is obtained by appropriately modifying u and v in (6.2). A_2 hence remains positive also in this more general case.

We have thus proved that $\mathcal{N} = 2$ gauged supergravity as it arises in Calabi-Yau like compactifications at string tree level (i.e. with hypermultiplet metric as given in [56], and gaugings of axionic isometries) does not permit de Sitter solutions. Due to the consistency

of the truncation, this 4d result also follows from the 10d no-go theorem of Maldacena-Nuñez [61]. Note however that our 4d reasoning continues to hold for an *arbitrary* vector multiplet sector, i.e. including all possible worldsheet instanton corrections.

The two contributions to (6.1) arise upon compactification from the NSNS and the RR sector respectively, see (4.3) and (4.5). The positivity of A_2 is also manifest here.

6.2 Non-supersymmetric Nearly Kähler companions

The 10d analysis of subsection 3.2 reveals that, given a choice of the RR fluxes G_0 and G_6 , with all the other fluxes vanishing, there exists a single Nearly Kähler supersymmetric vacuum on the cosets (2.1). This solution is also recovered adopting the 4d approach, as discussed in [27, 11].

It is possible to show that, under the same conditions, the 4d tree level scalar potential V also admits non-supersymmetric Nearly Kähler extrema. In the following formulae, we introduce the sum of the geometric fluxes $q \equiv \sum_a q_a$, we rename the RR fluxes as $e_0 \rightarrow e$, $m^0 \rightarrow m$, and we call the equal v^a and the equal b^a respectively v and b .

We obtain three Nearly Kähler extrema, lying at

$$v = \frac{\sqrt{15}}{2} \left(\frac{1}{20I} \left| \frac{e}{m} \right| \right)^{1/3}, \quad b = \frac{1}{2} \left(\frac{1}{20I} \frac{e}{m} \right)^{1/3}, \quad \tilde{\xi} = \frac{24Imb^2}{q}, \quad e^{2\varphi} = \frac{5q^2}{48I^2m^2v^4}, \quad (6.3)$$

$$v = \sqrt{3} \left(\frac{1}{20I} \left| \frac{e}{m} \right| \right)^{1/3}, \quad b = - \left(\frac{1}{20I} \frac{e}{m} \right)^{1/3}, \quad \tilde{\xi} = -\frac{12Imb^2}{q}, \quad e^{2\varphi} = \frac{q^2}{12I^2m^2v^4}, \quad (6.4)$$

and

$$v = \left(\frac{1}{\sqrt{5}I} \left| \frac{e}{m} \right| \right)^{1/3}, \quad b = 0 = \tilde{\xi}, \quad e^{2\varphi} = \frac{5q^2}{36I^2m^2v^4}. \quad (6.5)$$

By comparing to section 3.2, we learn that the only extremum preserving supersymmetry is (6.3), provided $e < 0$.

Thanks to the consistency of the reduction, the non-supersymmetric extrema of V found here also solve the 10d equations of motion. Unlike the situation for the supersymmetric solution, stability is of course no longer guaranteed. As in any truncation scheme, a full stability analysis can only take place in the higher dimensional theory. What we can offer in our 4 dimensional theory is a stability analysis with regard to the modes we retain. To this end, we rescale the scalar fields¹⁶ ($v^a, b^a, \varphi, \tilde{\xi}$) to obtain canonically normalized kinetic terms, and then diagonalize the mass matrix at the respective solutions.

The case $\frac{G_2}{\text{SU}(3)}$ is depicted in figure 1: the first two extrema (6.3) and (6.4) are minima, while the remaining extremum is a saddle point. For $\frac{\text{SU}(3)}{\text{U}(1) \times \text{U}(1)}$ and $\frac{\text{Sp}(2)}{\text{S}(\text{U}(2) \times \text{U}(1))}$, (6.4) is a minimum, whereas due to modes leading away from the Nearly Kähler locus $v^a = v$

¹⁶Note that the shift symmetry of a and ξ is gauged, the background value of these fields is hence a gauge choice.




Figure 1: The potential for $\frac{G_2}{SU(3)}$: we plot the rescaled potential $e^{\frac{5}{3}} m^{\frac{1}{3}} I^{\frac{4}{3}} V$ as a function of $(Im/e)^{\frac{1}{3}} b$ and $|Im/e|^{\frac{1}{3}} v$, at the extremum of φ and $\tilde{\xi}$. The deepest minimum corresponds to solution (6.4). The cut of the plot at $V = 0$ is due to the constraint $e^{\varphi(b,v)} > 0$.

for all a , (6.3) is merely a saddle point, as is (6.5). To analyse stability, we compare the magnitude of the negative masses at the saddle points with the Breitenlohner-Freedman bound

$$m_{\text{tachyonic}}^2 \geq -\frac{3}{4}|V|.$$

All extrema (including the saddle point depicted in figure 1) prove stable.

Finally, we remark that α' and string loop corrections can be safely neglected for the solutions above by tuning the RR fluxes e and m in such a way that the internal volume $Vol \equiv v^3 I \sim e/m$ becomes sufficiently large and the string coupling constant $e^\phi \equiv e^\varphi \sqrt{Vol} \sim e^{-\frac{1}{6}} m^{-\frac{5}{6}}$ becomes small (recall the definition (A.2) of the 4d dilaton). We can study moderately large string coupling by invoking the corrected potential (5.4). A preliminary numerical analysis for the coset $\frac{G_2}{SU(3)}$ indicates that all three AdS extrema survive string loop corrections.

6.3 de Sitter vacua at all string loop order?

In face of the no-go result for de Sitter vacua obtained in subsection 6.1, we would like to analyse how loop corrections modify the outcome of this study. Of course, to guarantee the consistency of the truncation, the analysis in section 4 must be extended beyond the two derivative case. However, the arguments put forth in subsection 4.1 in favor of consistency apply to the additional terms as well. We will also assume in this section that $c \neq 0$, as in the Calabi-Yau case. Note that by the results above, we can perform an (almost) complete analysis of the full loop corrected potential. The identification of the physical coordinate φ and the Calderbank-Pedersen coordinate ρ , which is modified order by order in the string coupling and is not available, merely enters in identifying the range of the CP coordinate, see below. Away from very strong coupling (in which brane instanton corrections would have to be considered regardless), this does not affect the

search for de Sitter minima.

Focusing on the ρ dependence of the potential (5.4) and taking the obvious positivity constraints on the coefficients into account does not rule out de Sitter vacua. One can then proceed to derive various constraints on these coefficients. E.g., by noting that the potential (5.4) has the form

$$V(\rho) = P(\rho)Q(\rho),$$

with $P(\rho) = \frac{1}{(\rho^2 - c)^2}$, we obtain

$$\begin{aligned} V(\rho_0) &= -\frac{P^2}{P'}Q'|_{\rho_0} \\ &= \frac{Q_A^u \bar{Q}_B^v}{2(\rho_0^2 - c)} \left[\left(-\frac{1}{2}(\text{Im } \mathcal{N})^{-1AB} - 3e^K X^A \bar{X}^B \right) (4d\xi^2)_{uv} \right. \\ &\quad \left. + \frac{c}{(\rho_0^2 + c)^2} e^K X^A \bar{X}^B (da + \tilde{\xi} d\xi)_{uv}^2 \right], \end{aligned}$$

where ρ_0 signifies the value of ρ at a minimum of the potential. Since c is negative for the cosets we are considering, a de Sitter vacuum requires the first term in the square bracket to be positive at the minimum of the potential. This term is proportional the tree level NSNS contribution to V , given in eq. (4.3). Hence, our necessary condition translates into the following inequality involving the internal NSNS 3-form and Ricci scalar

$$H \lrcorner H - 2R_6 > 0.$$

Recalling eq. (4.4), this is obviously true whenever the non-vanishing $\text{SU}(3)$ torsion classes satisfy $15|W_1|^2 < W_2 \lrcorner \bar{W}_2$. For the simple case of Nearly Kähler manifolds (i.e. when $W_2 = 0$) the inequality is however non-trivial, and reads $3b^2 - 5v^2 > 0$.

We hope to return to a more complete analysis of the all loop corrected potential in the near future.

Acknowledgements

We would like to thank Alessandro Tomasiello for collaboration in the initial stages of this project. We also acknowledge useful discussions with Adel Bilal, Paul Koerber, Luca Martucci, Ruben Minasian, and Pierre Vanhove. DC thanks the Service de Physique Théorique et Mathématique de l'Université Libre de Bruxelles, where part of this work was done, for hospitality and financial support. DC and AK are supported in part by the EU grant MRTN-CT-2004-005104. In addition, DC is partially supported by the EU grant MRTN-CT-2004-512194, by the French grant ANR(CNRS-USAR) no.05-BLAN-0079-01 and by the ‘‘Programme Vinci 2006 de l'Université Franco-Italienne’’. AK is supported in part by *l'Agence Nationale de la Recherche* under the grants ANR-06-BLAN-3_137168 and ANR-05-BLAN-0029-01.

A Details of the dimensional reduction

The G -invariant reduction ansatz strongly constrains the dependence of all the higher dimensional fields on the G/H coordinates, relegating it into the coframe $e^{\underline{m}}$ introduced in subsection 2.1. In particular, the most general G -invariant 10d metric is (here and in the following, the hat denotes 10d fields):

$$d\hat{s}^2 = e^{2\varphi(x)} g_{\mu\nu}(x) dx^\mu \otimes dx^\nu + g_{\underline{mn}}(x) e^{\underline{m}}(y) \otimes e^{\underline{n}}(y) , \quad (\text{A.1})$$

where x^μ and $y^{\underline{m}}$ are respectively coordinates on the 4d spacetime and the internal manifold M_6 , and $g_{\underline{mn}}$ satisfies the G -invariance condition discussed in subsection 2.1. Components of the 10d metric with mixed 4d-6d indices are not allowed since there are no left-invariant 1-forms on our coset manifolds (2.1). Since the invariant scalars on the coset are necessarily constant, a nontrivial warp factor is also not permitted (see [62, 63] for recent discussions of a non-trivial warp factor in the $\mathcal{N} = 1$ context). The Weyl factor $e^{2\varphi(x)}$ in front of the 4d metric is needed in order to obtain a canonical lower dimensional Einstein-Hilbert term $\int_{M_4} \text{vol}_4 R_4$ from the string frame higher dimensional action $\int_{M_{10}} \text{vol}_{10} e^{-2\phi} \hat{R}$, with

$$\varphi(x) = \phi(x) - \frac{1}{2} \log \int_{M_6} d^6 y \sqrt{g_6} , \quad (\text{A.2})$$

where $\phi(x)$ is the 10d dilaton and $\sqrt{g_6} \equiv \sqrt{\det g_{mn}(x, y)} = \sqrt{\det g_{mn}(x)} |\det e^{\underline{p}}_{\underline{q}}(y)|$. Notice that, thanks to this factorization of the x and y dependence, $\partial_\mu \log \sqrt{g_6}$ does not depend on the internal coordinates, and

$$\partial_\mu \varphi = \partial_\mu \phi - \frac{1}{2} \partial_\mu \log \sqrt{g_6} . \quad (\text{A.3})$$

The ansatz for the 10d supergravity field strengths must be chosen consistently with their Bianchi identities. For instance, from the Bianchi identity $d\hat{F}_2 = \hat{H}\hat{F}_0$, one sees that if $\hat{F}_0 \neq 0$, then the NSNS 3-form $\hat{H}$ has to be exact: $\hat{H} = d\hat{B}$, with a globally defined 2-form potential $\hat{B}$. The most general $\hat{B}$ respecting left-invariance on M_6 is

$$\hat{B} = B + b , \quad (\text{A.4})$$

where $B(x)$ is along 4d spacetime, while $b(x, y) = b^a(x) \omega_a(y)$ lives on M_6 (the left-invariant 2-forms ω_a are given in subsection 2.1).

We deal with the expansion of the RR fields in subsection A.2.

A.1 Special Kähler geometry from the NSNS sector

Combining the 2-form J of subsection 2.2 and the internal NS field b we introduce $t = b + iJ$, whose expansion $t = t^a \omega_a$ on the basis 2-forms defines the complex 4d scalars $t^a = b^a + i v^a$. The associated kinetic term is determined by

$$\frac{1}{8} g^{mp} g^{nq} (\partial_\mu g_{mn} \partial^\mu g_{pq} + \partial_\mu b_{mn} \partial^\mu b_{pq}) = \frac{1}{4 \text{Vol}} \int_{M_6} \partial_\mu t \wedge * \partial^\mu \bar{t} = \mathcal{G}_{ab} \partial_\mu t^a \partial^\mu \bar{t}^b , \quad (\text{A.5})$$

where the l.h.s. originates from the reduction of the 10d Ricci scalar and $\hat{H}^2$ terms, while the σ -model metric $\mathcal{G}_{ab}$ was introduced in eq. (2.12). The first equality in (A.5) is derived recalling that the internal metric is fixed by the forms J and Ω defining the $SU(3)$ structure: indeed, calling $\mathcal{I}$ the almost complex structure induced by Ω , we have $g_{mn} = J_{mp}\mathcal{I}^p_n$. Notice that we get no contribution from the variation of $\mathcal{I}$ since the associated Ω , given in eq. (2.14), is rigid.

The metric $\mathcal{G}_{ab}$ is special Kähler: indeed, it can be obtained via $\mathcal{G}_{ab} = \frac{\partial^2 K}{\partial t^a \partial t^b}$ from the Kähler potential

$$K = -\log \frac{4}{3} \int J \wedge J \wedge J = -\log 8Vol .$$

It in turn is determined by a prepotential $\mathcal{F}$ via the special Kähler geometry formula $K = -\log i(\bar{X}^A \mathcal{F}_A - X^A \bar{\mathcal{F}}_A)$, where $X^A \equiv (X^0, X^a) = (1, -t^a)$ and $\mathcal{F}_A = \frac{\partial \mathcal{F}(X)}{\partial X^A}$.

For each of the cosets we consider, the explicit expressions of $\mathcal{G}_{ab}$ and Vol are given in table 1. The (cubic) prepotential reads

$$\mathcal{F}(X) = \frac{1}{6} \mathcal{K}_{abc} \frac{X^a X^b X^c}{X^0} ,$$

where the non-vanishing triple intersection numbers $\mathcal{K}_{abc} := \int \omega_a \wedge \omega_b \wedge \omega_c$ (recall the 2-forms ω_a in subsection 2.1) are

$$\begin{aligned} \mathcal{K}_{123} &= I & \text{for } \frac{SU(3)}{U(1) \times U(1)} \\ \mathcal{K}_{112} &= 2I & \text{for } \frac{Sp(2)}{S(U(2) \times U(1))} \\ \mathcal{K}_{111} &= 6I & \text{for } \frac{G_2}{SU(3)} . \end{aligned} \tag{A.6}$$

The period matrix $\mathcal{N}_{AB}$ of special Kähler geometry is given by the formula (see e.g. [64])

$$\mathcal{N}_{AB} = \bar{\mathcal{F}}_{AB} + 2i \frac{\text{Im}(\mathcal{F}_{AC}) X^C \text{Im}(\mathcal{F}_{BD}) X^D}{X^E \text{Im}(\mathcal{F}_{EF}) X^F} , \quad \text{where } \mathcal{F}_{AB} \equiv \frac{\partial^2 \mathcal{F}}{\partial X^A \partial X^B} .$$

Equivalently, we can directly obtain it from the coset geometry via [18]:

$$\begin{aligned} (\text{Im} \mathcal{N})^{-1 AB} &= - \int \langle \tilde{\omega}^A, *_b \tilde{\omega}^B \rangle , & [\text{Re} \mathcal{N} (\text{Im} \mathcal{N})^{-1}]_A^B &= - \int \langle \omega_A, *_b \tilde{\omega}^B \rangle , \\ [\text{Im} \mathcal{N} + \text{Re} \mathcal{N} (\text{Im} \mathcal{N})^{-1} \text{Re} \mathcal{N}]_{AB} &= - \int \langle \omega_A, *_b \omega_B \rangle , \end{aligned}$$

with $*_b(\cdot) \equiv e^{-b} * \lambda(e^b \cdot)$. The operator λ and the pairing $\langle \cdot, \cdot \rangle$ were defined below (2.10).

We obtain the matrices

$$\text{Im} \mathcal{N} = -Vol \begin{pmatrix} 1 + 4\mathcal{G}_{ab} b^a b^b & 4\mathcal{G}_{ab} b^b \\ 4\mathcal{G}_{ab} b^b & 4\mathcal{G}_{ab} \end{pmatrix} , \tag{A.7}$$

$$\text{Re} \mathcal{N} = - \begin{pmatrix} \frac{1}{3} \mathcal{K}_{abc} b^a b^b b^c & \frac{1}{2} \mathcal{K}_{abc} b^b b^c \\ \frac{1}{2} \mathcal{K}_{abc} b^b b^c & \mathcal{K}_{abc} b^c \end{pmatrix} . \tag{A.8}$$

A.2 The RR sector

In order to reduce the RR sector we specialize the general procedure described in section 5 of ref. [10] for M_6 corresponding to our coset spaces. Adopting the democratic formulation of type IIA supergravity [36], the RR degrees of freedom can be encoded in a field strength $\hat{\mathbf{G}}$ consisting of a formal sum of forms of all possible even degrees, satisfying

$$\text{Bianchi identity : } d\hat{\mathbf{G}} = 0 \quad (\text{A.9})$$

$$\text{self-duality constraint : } \hat{\mathbf{F}} = \lambda(*\hat{\mathbf{F}}), \quad \text{where } \hat{\mathbf{F}} \equiv e^{\hat{B}}\hat{\mathbf{G}} \text{ and } \lambda(\hat{F}_{(k)}) = (-)^{\frac{k}{2}}\hat{F}_{(k)}. \quad (\text{A.10})$$

Due to the self-duality constraint, the equations of motion for the RR degrees of freedom are equivalent to the Bianchi identities.

We implement the reduction ansatz by expanding $\hat{\mathbf{G}}$ on the basis of left-invariant internal forms introduced in subsection 2.1,

$$\hat{\mathbf{G}} = (G_{(0)}^A + G_{(2)}^A + G_{(4)}^A)\omega_A - (\tilde{G}_{(0)A} + \tilde{G}_{(2)A} + \tilde{G}_{(4)A})\tilde{\omega}^A + (G_{(1)} + G_{(3)})\alpha - (\tilde{G}_{(1)} + \tilde{G}_{(3)})\beta. \quad (\text{A.11})$$

$G_{(p)}(x)$ and $\tilde{G}_{(p)}(x)$ are p -forms in 4d spacetime. Plugging this expansion into eqs. (A.9), (A.10), and going through the derivation of [10], one identifies the 4d variables

$$\begin{aligned} G_{(0)}^A &= m^A, & \tilde{G}_{(0)A} &= e_A + q_A \tilde{\xi} \\ G_{(1)} &= D\xi \equiv d\xi - q_a A^a, & \tilde{G}_{(1)} &= d\tilde{\xi} \\ G_{(2)}^A &= dA^A, & \tilde{G}_{(2)A} + B\tilde{G}_{(0)A} &= \text{Im}\mathcal{N}_{AB} * F^B + \text{Re}\mathcal{N}_{AB} F^B \\ G_{(3)} &= -B \wedge D\xi + e^{2\varphi} * d\tilde{\xi}, & \tilde{G}_{(3)} &= -B \wedge d\tilde{\xi} - e^{2\varphi} * D\xi \end{aligned} \quad (\text{A.12})$$

$$\begin{aligned} G_{(4)}^A + B \wedge G_{(2)}^A + \frac{1}{2}B^2 G_{(0)}^A &= e^{4\varphi} [(\text{Im}\mathcal{N})^{-1}(\tilde{G}_{(0)} - \text{Re}\mathcal{N}G_{(0)})]^A * 1 \\ \tilde{G}_{(4)A} + B \wedge \tilde{G}_{(2)A} + \frac{1}{2}B^2 \tilde{G}_{(0)A} &= e^{4\varphi} [-\text{Im}\mathcal{N}G_{(0)} + \text{Re}\mathcal{N}(\text{Im}\mathcal{N})^{-1}(\tilde{G}_{(0)} - \text{Re}\mathcal{N}G_{(0)})]_A * 1 \end{aligned}$$

where the propagating fields are the two real scalars $\xi, \tilde{\xi}$ and the 1-forms A^A . We also introduced the modified field strengths

$$F^A \equiv dA^A + m^A B. \quad (\text{A.13})$$

Furthermore we introduce $q_A = (0, q_a)$, the q_a being the geometric fluxes defined in subsection 2.1.4, while m^A, e_A are constant flux parameters satisfying $q_a m^a = 0$. Notice that one of the e_a is redundant, since it can be eliminated via a constant shift of $\tilde{\xi}$. This reflects the fact that on our cosets the linear combination $q_a \tilde{\omega}^a$ is exact (see eq. (2.11)), and therefore doesn't support any flux.

The residual content of (A.9)–(A.11) not included in eqs. (A.12) consists of a set of equations to be read as the EoM for $\xi, \tilde{\xi}$ and A^A . We use these equations to reconstruct the

4d action $S_{\text{RR}}^{(4)}$ of subsection 4.2. In particular, we infer the RR contribution to the 4d scalar potential,

$$V_{\text{RR}} = -\frac{e^{4\varphi}}{4} [G_{(0)} \text{Im} \mathcal{N} G_{(0)} + (\tilde{G}_{(0)} - G_{(0)} \text{Re} \mathcal{N})(\text{Im} \mathcal{N})^{-1}(\tilde{G}_{(0)} - \text{Re} \mathcal{N} G_{(0)})] . \quad (\text{A.14})$$

Substitution of the explicit expressions for $G_{(0)}$ and $\tilde{G}_{(0)}$ given in (A.12) yields eq. (4.5).

As a last remark, we stress that the whole procedure of section 5 of [10] applies here with no need to take any integral over M_6 . In other words, once the left-invariant truncation ansatz has been plugged in, the dependence of eqs.(A.9), (A.10) on the internal coordinates automatically factorizes out.

References

- [1] M. Gomez-Reino, J. Louis and C. A. Scrucca, *No metastable de Sitter vacua in $N=2$ supergravity with only hypermultiplets*, arXiv:0812.0884 [hep-th].
- [2] F. Saueressig, U. Theis and S. Vandoren, *On de Sitter vacua in type IIA orientifold compactifications*, Phys. Lett. B **633**, 125 (2006) [arXiv:hep-th/0506181].
- [3] M. Bodner, A. C. Cadavid and S. Ferrara, *(2,2) vacuum configurations for type IIA superstrings: $N=2$ supergravity Lagrangians and algebraic geometry*, Class. Quant. Grav. **8** (1991) 789.
- [4] R. Bohm, H. Gunther, C. Herrmann and J. Louis, *Compactification of type IIB string theory on Calabi-Yau threefolds*, Nucl. Phys. B **569** (2000) 229 [arXiv:hep-th/9908007].
- [5] S. Gurrieri, J. Louis, A. Micu and D. Waldram, *Mirror symmetry in generalized Calabi-Yau compactifications*, Nucl. Phys. B **654**, 61 (2003) [arXiv:hep-th/0211102].
- [6] S. B. Giddings, S. Kachru and J. Polchinski, *Hierarchies from fluxes in string compactifications*, Phys. Rev. D **66** (2002) 106006 [arXiv:hep-th/0105097].
- [7] M. Grana, R. Minasian, M. Petrini and A. Tomasiello, *Supersymmetric backgrounds from generalized Calabi-Yau manifolds*, JHEP **0408** (2004) 046 [arXiv:hep-th/0406137].
- [8] J. Polchinski and A. Strominger, *New Vacua for Type II String Theory*, Phys. Lett. B **388** (1996) 736 [arXiv:hep-th/9510227].
- [9] L. Andrianopoli, M. Bertolini, A. Ceresole, R. D’Auria, S. Ferrara, P. Fre and T. Margri, *$N = 2$ supergravity and $N = 2$ super Yang-Mills theory on general scalar manifolds: Symplectic covariance, gaugings and the momentum map*, J. Geom. Phys. **23**, 111 (1997) [arXiv:hep-th/9605032].
- [10] D. Cassani, *Reducing democratic type II supergravity on $SU(3) \times SU(3)$ structures*, JHEP **0806** (2008) 027 [arXiv:0804.0595 [hep-th]].
- [11] A. K. Kashani-Poor, *Nearly Kaehler Reduction*, JHEP **0711** (2007) 026 [arXiv:0709.4482 [hep-th]].

- [12] M. J. Duff, B. E. W. Nilsson and C. N. Pope, *Kaluza-Klein Supergravity*, Phys. Rept. **130** (1986) 1.
- [13] A. K. Kashani-Poor and R. Minasian, *Towards reduction of type II theories on $SU(3)$ structure manifolds*, JHEP **0703**, 109 (2007) [arXiv:hep-th/0611106].
- [14] D. J. Gross and E. Witten, *Superstring Modifications Of Einstein's Equations*, Nucl. Phys. B **277** (1986) 1.
- [15] M. B. Green and J. H. Schwarz, *Supersymmetrical Dual String Theory. 3. Loops And Renormalization*, Nucl. Phys. B **198** (1982) 441.
- [16] N. Sakai and Y. Tanii, *One Loop Amplitudes And Effective Action In Superstring Theories*, Nucl. Phys. B **287** (1987) 457.
- [17] K. Peeters, P. Vanhove and A. Westerberg, *Supersymmetric higher-derivative actions in ten and eleven dimensions, the associated superalgebras and their formulation in superspace*, Class. Quant. Grav. **18** (2001) 843 [arXiv:hep-th/0010167].
- [18] D. Cassani and A. Bilal, *Effective actions and $N=1$ vacuum conditions from $SU(3)\times SU(3)$ compactifications*, JHEP **0709** (2007) 076 [arXiv:0707.3125 [hep-th]].
- [19] S. S. Haque, G. Shiu, B. Underwood and T. Van Riet, *Minimal simple de Sitter solutions*, arXiv:0810.5328 [hep-th].
- [20] C. Caviezel, P. Koerber, S. Kors, D. Lust, T. Wrase and M. Zagermann, *On the Cosmology of Type IIA Compactifications on $SU(3)$ -structure Manifolds*, arXiv:0812.3551 [hep-th].
- [21] R. Flauger, S. Paban, D. Robbins and T. Wrase, *On Slow-roll Moduli Inflation in Massive IIA Supergravity with Metric Fluxes*, arXiv:0812.3886 [hep-th].
- [22] P. Koerber, D. Lust and D. Tsimpis, *Type IIA AdS_4 compactifications on cosets, interpolations and domain walls*, JHEP **0807**, 017 (2008) [arXiv:0804.0614 [hep-th]].
- [23] M. Grana, *Flux compactifications in string theory: A comprehensive review*, Phys. Rept. **423** (2006) 91 [arXiv:hep-th/0509003].
- [24] S. Chiossi and S. Salamon, *The intrinsic torsion of $SU(3)$ and G_2 structures*, arXiv:math/0202282.
- [25] F. Mueller-Hoissen and R. Stuckl, *Coset spaces and ten-dimensional unified theories*, Class. Quant. Grav. **5** (1988) 27.
- [26] K. Behrndt and M. Cvetič, *General $N = 1$ Supersymmetric Flux Vacua of (Massive) Type IIA String Theory*, Phys. Rev. Lett. **95** (2005) 021601 [arXiv:hep-th/0403049].
- [27] T. House and E. Palti, *Effective action of (massive) IIA on manifolds with $SU(3)$ structure*, Phys. Rev. D **72**, 026004 (2005) [arXiv:hep-th/0505177].
- [28] A. Tomasiello, *New string vacua from twistor spaces*, Phys. Rev. D **78** (2008) 046007 [arXiv:0712.1396 [hep-th]].
- [29] G. Aldazabal and A. Font, *A second look at $\mathcal{N} = 1$ supersymmetric AdS_4 vacua of type IIA supergravity*, JHEP **0802** (2008) 086 [arXiv:0712.1021 [hep-th]].
- [30] C. Caviezel, P. Koerber, S. Kors, D. Lust, D. Tsimpis and M. Zagermann, *The*

- effective theory of type IIA AdS₄ compactifications on nilmanifolds and cosets*, Class. Quant. Grav. **26**, 025014 (2009) [arXiv:0806.3458 [hep-th]].
- [31] A. Chatzistavrakidis, P. Manousselis and G. Zoupanos, *Reducing the Heterotic Supergravity on nearly-Kahler coset spaces* arXiv:0811.2182 [hep-th].
 - [32] G. W. Moore and E. Witten, *Self-duality, Ramond-Ramond fields, and K-theory*, JHEP **0005** (2000) 032 [arXiv:hep-th/9912279].
 - [33] D. E. Diaconescu, G. W. Moore and E. Witten, *E(8) gauge theory, and a derivation of K-theory from M-theory*, Adv. Theor. Math. Phys. **6** (2003) 1031 [arXiv:hep-th/0005090].
 - [34] M. Karoubi, *K-theory*, Springer Verlag, 2008.
 - [35] D. Lust and D. Tsimpis, *Supersymmetric AdS(4) compactifications of IIA supergravity*, JHEP **0502** (2005) 027 [arXiv:hep-th/0412250].
 - [36] E. Bergshoeff, R. Kallosh, T. Ortin, D. Roest and A. Van Proeyen, *New formulations of D = 10 supersymmetry and D8 – O8 domain walls*, Class. Quant. Grav. **18** (2001) 3359 [arXiv:hep-th/0103233].
 - [37] L. Castellani, L. J. Romans and N. P. Warner, *Symmetries Of Coset Spaces And Kaluza-Klein Supergravity*, Annals Phys. **157** (1984) 394.
 - [38] M. J. Duff, B. E. W. Nilsson, C. N. Pope and N. P. Warner, *On The Consistency Of The Kaluza-Klein Ansatz*, Phys. Lett. B **149** (1984) 90.
 - [39] B. de Wit and H. Nicolai, *The Consistency of the S⁷ Truncation in D=11 Supergravity*, Nucl. Phys. B **281** (1987) 211.
 - [40] M. Cvetič, H. Lu and C. N. Pope, *Consistent Kaluza-Klein sphere reductions*, Phys. Rev. D **62** (2000) 064028 [arXiv:hep-th/0003286].
 - [41] M. J. Duff and C. N. Pope, *Consistent Truncations In Kaluza-Klein Theories*, Nucl. Phys. B **255** (1985) 355.
 - [42] R. Coquereaux and A. Jadczyk, *Consistency of the G invariant Kaluza-Klein scheme*, Nucl. Phys. B **276** (1986) 617.
 - [43] A. Chatzistavrakidis, P. Manousselis, N. Prezas and G. Zoupanos, *On the consistency of coset space dimensional reduction*, Phys. Lett. B **656** (2007) 152 [arXiv:0708.3222 [hep-th]]; *Coset Space Dimensional Reduction of Einstein–Yang–Mills theory*, Fortsch. Phys. **56** (2008) 389 [arXiv:0712.2717 [hep-th]].
 - [44] C. M. Hull and R. A. Reid-Edwards, *Flux compactifications of string theory on twisted tori*, arXiv:hep-th/0503114.
 - [45] G. Dall’Agata and N. Prezas, *Scherk-Schwarz reduction of M-theory on G₂-manifolds with fluxes*, JHEP **0510** (2005) 103 [arXiv:hep-th/0509052].
 - [46] J. P. Gauntlett and O. Varela, *Consistent Kaluza-Klein Reductions for General Supersymmetric AdS Solutions*, Phys. Rev. D **76**, 126007 (2007) [arXiv:0707.2315 [hep-th]].
 - [47] J. P. Gauntlett, S. Kim, O. Varela and D. Waldram, *Consistent supersymmetric Kaluza–Klein truncations with massive modes*, arXiv:0901.0676 [hep-th].

- [48] R. D'Auria, S. Ferrara, M. Trigiante and S. Vaula, *Gauging the Heisenberg algebra of special quaternionic manifolds*, Phys. Lett. B **610** (2005) 147 [arXiv:hep-th/0410290].
- [49] A. Tomasiello, *Topological mirror symmetry with fluxes*, JHEP **0506** (2005) 067 [arXiv:hep-th/0502148].
- [50] M. Grana, J. Louis and D. Waldram, *Hitchin functionals in $N = 2$ supergravity*, JHEP **0601**, 008 (2006) [arXiv:hep-th/0505264].
- [51] L. Bedulli and L. Vezzoni *The Ricci tensor of $SU(3)$ -manifolds*, J. Geom. Phys. 57 (2007), n. 4, 1125 [arXiv:math/0606786].
- [52] J. Michelson, *Compactifications of type IIB strings to four dimensions with non-trivial classical potential*, Nucl. Phys. B **495** (1997) 127 [arXiv:hep-th/9610151].
- [53] G. Dall'Agata, R. D'Auria, L. Sommovigo and S. Vaula, *$D = 4$, $N = 2$ gauged supergravity in the presence of tensor multiplets*, Nucl. Phys. B **682**, 243 (2004) [arXiv:hep-th/0312210].
- [54] R. D'Auria, L. Sommovigo and S. Vaula, *$N = 2$ supergravity Lagrangian coupled to tensor multiplets with electric and magnetic fluxes*, JHEP **0411**, 028 (2004) [arXiv:hep-th/0409097].
- [55] R. D'Auria, S. Ferrara and P. Fre, *Special and quaternionic isometries: general couplings in $N=2$ supergravity and the scalar potential*, Nucl. Phys. B **359** (1991) 705.
- [56] S. Ferrara and S. Sabharwal, *Quaternionic Manifolds for Type II Superstring Vacua of Calabi-Yau Spaces*, Nucl. Phys. B **332**, 317 (1990).
- [57] D. M. J. Calderbank and H. Pedersen, *Selfdual Einstein metrics with torus symmetry*, arXiv:math/0105263.
- [58] A. Strominger, *Loop corrections to the universal hypermultiplet*, Phys. Lett. B **421**, 139 (1998) [arXiv:hep-th/9706195].
- [59] I. Antoniadis, R. Minasian, S. Theisen and P. Vanhove, *String loop corrections to the universal hypermultiplet*, Class. Quant. Grav. **20**, 5079 (2003) [arXiv:hep-th/0307268].
- [60] S. Kachru and A. K. Kashani-Poor, *Moduli potentials in type IIA compactifications with RR and NS flux*, JHEP **0503** (2005) 066 [arXiv:hep-th/0411279].
- [61] J. M. Maldacena and C. Nunez, *Supergravity description of field theories on curved manifolds and a no go theorem*, Int. J. Mod. Phys. A **16** (2001) 822 [arXiv:hep-th/0007018].
- [62] P. Koerber and L. Martucci, *From ten to four and back again: how to generalize the geometry*, JHEP **0708**, 059 (2007) [arXiv:0707.1038 [hep-th]].
- [63] G. Shiu, G. Torroba, B. Underwood and M. R. Douglas, *Dynamics of Warped Flux Compactifications*, JHEP **0806** (2008) 024 [arXiv:0803.3068 [hep-th]].
- [64] B. Craps, F. Roose, W. Troost and A. Van Proeyen, *What is special Kähler geometry?*, Nucl. Phys. B **503** (1997) 565 [arXiv:hep-th/9703082].