

NOT SO NON-RENORMALIZABLE GRAVITY*

DIRK KREIMER

ABSTRACT. We review recent ideas [1] how gravity might turn out to be a renormalizable theory after all.

1. INTRODUCTION

Renormalizable perturbative quantum field theories are embarrassingly successful in describing observed physics. Whilst their mathematical structure is still a challenge albeit an entertaining one, they are testimony to some of the finest achievements in our understanding of nature. The physical law as far as is insensitive to the surrounding geometry seems completely described by such theories. Alas, if we incorporate gravity, and want to quantize it, we seem at a loss.

In this talk, we report on some recent work [1] which might give hope. Our main purpose is to review the basic idea and to put it into context.

As in [1], we will proceed by a comparison of the structure of a renormalizable theory, quantum electrodynamics in four dimensions, and gravity.

It is the role of the Hochschild cohomology [2] in those two different situations which leads to surprising new insights. We will discuss them at an elementary level for the situation of pure gravity. We also allow, in the spirit of the workshop, for the freedom to muse about conceptual consequences at the end.

Acknowledgments. It is a pleasure to thank Bertfried Fauser and Eberhard Zeldner for inviting me to this workshop.

2. THE STRUCTURE OF DYSON-SCHWINGER EQUATIONS IN QED₄

2.1. The Green functions. Quantum electrodynamics in four dimensions of space-time (QED₄) is described in its short-distance behaviour by four Green functions

$$(1) \quad G^{\bar{\psi}\gamma\cdot\partial\psi}, G^{m\bar{\psi}\psi}, G^{\bar{\psi}\gamma\cdot A\psi}, G^{\frac{1}{4}F^2},$$

corresponding to the four monomials in its Lagrangian

$$(2) \quad L = \bar{\psi}\gamma\cdot\partial\psi - \bar{\psi}m\psi - \bar{\psi}\gamma\cdot A\psi - \frac{1}{4}F^2.$$

Here, $G^i = G^i(\alpha, L)$, with α the fine structure constant and $L = \ln q^2/\mu^2$, so that we work in a MOM scheme, subtract at $q^2 = \mu^2$, project the vertex function to its scalar formfactor $G^{\bar{\psi}\gamma\cdot\partial\psi}$ with UV divergences evaluated at zero photon momentum. Similarly the other Green functions are normalized as to be the multiplicative quantum corrections to the tree level monomials above, in momentum space.

*Talk given at "Max Planck Institute for Mathematics in the Natural Sciences, Leipzig". Work supported in parts by grant NSF-DMS/0603781. Author supported by CNRS.

In perturbation theory, the degree of divergence of a graph Γ with f external fermion lines and m external photon lines in D dimensions is

$$(3) \quad \omega_D(\Gamma) = \frac{3}{2}f + m - D - (D-4)(|\Gamma| - 1) \Rightarrow \omega_4(\Gamma) = \frac{3}{2}f + m - 4.$$

This is independent of the loop number for QED₄, $D = 4$, and is a sole function of the number and type of external legs. $\omega_D(\Gamma)$ determines the number of derivatives with respect to masses or external momenta needed to render a graph logarithmically divergent, and hence identifies the top-level residues which drive the iteration of Feynman integrals according to the quantum equations of motion [3].

We define these four Green functions as an evaluation by renormalized Feynman rules of a series of one-particle irreducible (1PI) Feynman graphs $\Gamma \in \mathcal{FG}_i$. These series are determined as a fixpoint of the following system in Hochschild cohomology.

$$(4) \quad X^{\bar{\psi}\gamma \cdot \partial\psi} = 1 - \sum_{k=1}^{\infty} \alpha^k B_+^{\bar{\psi}\gamma \cdot \partial\psi, k} (X^{\bar{\psi}\gamma \cdot \partial\psi} Q^{2k}(\alpha)),$$

$$(5) \quad X^{\bar{\psi}\gamma \cdot A\psi} = 1 + \sum_{k=1}^{\infty} \alpha^k B_+^{\bar{\psi}\gamma \cdot A\psi, k} (X^{\bar{\psi}\gamma \cdot A\psi} Q^{2k}(\alpha)),$$

$$(6) \quad X^{\bar{\psi}m\psi} = 1 - \sum_{k=1}^{\infty} \alpha^k B_+^{\bar{\psi}m\psi, k} (X^{\bar{\psi}m\psi} Q^{2k}(\alpha)),$$

$$(7) \quad X^{\frac{1}{4}F^2} = 1 - \sum_{k=1}^{\infty} \alpha^k B_+^{\frac{1}{4}F^2, k} (X^{\frac{1}{4}F^2} Q^{2k}(\alpha)).$$

Here,

$$(8) \quad B_+^{i, k} = \sum_{|\gamma|=k, \Delta'(\gamma)=0, \gamma \in \mathcal{FG}_i} B_+^{\gamma}, \quad \forall i \in \mathcal{R}_{\text{QED}},$$

a sum over all Hopf algebra primitive graphs with given loop number k and contributing to superficially divergent amplitude i , and

$$(9) \quad B_+^{\gamma}(X) = \sum_{\Gamma \in \langle \Gamma \rangle} \frac{\text{bij}(\gamma, X, \Gamma)}{|X|_{\vee}} \frac{1}{\text{maxf}(\Gamma)} \frac{1}{(\gamma|X)} \Gamma,$$

where $\text{maxf}(\Gamma)$ is the number of maximal forests of Γ , $|X|_{\gamma}$ is the number of distinct graphs obtainable by permuting edges of X , $\text{bij}(\gamma, X, \Gamma)$ is the number of bijections of external edges of X with an insertion place in γ such that the result is Γ , and finally $(\gamma|X)$ is the number of insertion places for X in γ [4], and

$$(10) \quad \mathcal{R}_{\text{QED}} = \{\bar{\psi}\gamma \cdot \partial\psi, \bar{\psi}\gamma \cdot A\psi, m\bar{\psi}\psi, \frac{1}{4}F^2\}.$$

Also, we let

$$(11) \quad Q = \frac{X^{\bar{\psi}\gamma \cdot A\psi}}{X^{\bar{\psi}\gamma \cdot \partial\psi} \sqrt{X^{\frac{1}{4}F^2}}}.$$

The resulting maps $B_+^{i, K}$ are Hochschild closed

$$(12) \quad bB_+^{i, K} = 0,$$

in the sense of [5]. We have in fact

$$(13) \quad \Delta(B_+^\gamma(X)) = \sum_{\Gamma} n_{\Gamma, X, \gamma} \Gamma$$

where $n_{\Gamma, X, \gamma}$ can be determined from (9,12).

Furthermore, one can choose a basis of primitives γ [3] such that their Mellin transforms $M_\gamma(\rho)$ have the form

$$(14) \quad M_\gamma(\rho) = \int \iota_\gamma(k_i; q) |_{q^2=\mu^2} \prod_{s=1}^{|\gamma|} \frac{[k_s^2/\mu^2]^{-\rho/|\gamma|} d^4 k_i}{(2\pi)^4} \text{ for } 1 > \Re(\rho) > 0,$$

where the integrand ι_γ is a function of internal momenta k_i and an external momentum q , subtracted at $q^2 = \mu^2$.

The Dyson Schwinger equations then take the form

$$(15) \quad G_R^i(\alpha, L) = 1 \pm \lim_{\rho \rightarrow 0} \left[\sum_k \alpha^k \sum_{|\gamma|=k} G_r^i(\alpha, \partial_\rho) \mathcal{Q}(\alpha, \partial_\rho) M_\gamma(\rho) \left[\left(\frac{q^2}{\mu^2} \right)^{-\rho} - 1 \right] \right],$$

where

$$(16) \quad \Phi_R(X^i) = G_R^i(\alpha, L),$$

and

$$(17) \quad \Phi_R(Q) = \mathcal{Q}(\alpha, L),$$

is the invariant charge, all calculated with renormalized Feynman rules in the MOM scheme.

2.2. Gauge theoretic aspects. Using Ward identities, we can reduce the set $\mathcal{R}_{\text{QED}} = \{\bar{\psi}\gamma \cdot \partial \psi, m\bar{\psi}\psi, \bar{\psi}\gamma \cdot A\psi, \frac{1}{4}F^2\}$ to three elements upon identifying $G^{\bar{\psi}\gamma \cdot \partial \psi} = G^{\bar{\psi}\gamma \cdot A\psi}$. Using the Baker–Johnson–Willey gauge [6] we can furthermore trivialize

$$(18) \quad G^{\bar{\psi}\gamma \cdot \partial \psi} = G^{\bar{\psi}\gamma \cdot A\psi} = 1.$$

Using their work again [7], we have that $m\bar{\psi}\psi$ can be ignored in $\mathcal{R}_{\text{QED}}$.

We are hence left with the determination of a single gauge-independent Green function $G^{\frac{1}{4}F^2}$ which in the MOM scheme takes the form

$$(19) \quad G^{\frac{1}{4}F^2}(\alpha, L) = 1 - \sum_{k=1}^{\infty} \gamma_k(\alpha) L^k,$$

and the renormalization group determines [8]

$$(20) \quad \gamma_k(\alpha) = \frac{1}{k} \gamma_1(\alpha) (1 - \alpha \partial_\alpha) \gamma_{k-1}(\alpha).$$

Here, $\gamma_1(\alpha) = 2\psi(\alpha)/\alpha$, where $\psi(\alpha)$ is the MOM scheme β -function of QED, which is indeed half of the anomalous dimension γ_1 of the photon field in that scheme.

One can show that $\gamma_1(\alpha)$ as a perturbative series $(\gamma_1(\alpha) = \sum_{j=1}^{\infty} \gamma_{1,j} \alpha^j)$ is Gevrey-1 and that the series $\sum_{j=1}^{\infty} \gamma_{1,j} \alpha^j / j!$ has a finite radius of convergence, with a bound involving the lowest order contribution of the β -function and the one-instanton action [3].

Furthermore, $\gamma_1(\alpha)$ fulfills [3]

$$(21) \quad \gamma_1(\alpha) = P(\alpha) - \gamma_1(\alpha)(1 - \alpha \partial_\alpha) \gamma_1(\alpha),$$

an equation which has been studied in numerical detail recently [9], with more of its analytic structure to be exhibited there. In this equation, $P(\alpha)$ is obtained from the primitives of the Hopf algebra

$$(22) \quad P(\alpha) = \sum_{\gamma} \alpha^{|\gamma|} \lim_{\rho \rightarrow 0} \rho M_{\gamma}(\rho),$$

and $P(\alpha)$ is known perturbatively as a fifth order polynomial [10] and its asymptotics have been conjectured long ago [11].

This finishes our summary of QED₄ as a typical renormalizable theory.

2.3. Non-abelian gauge theory. The above approach to Green functions remains valid for a non-abelian gauge theory with the definition of a single invariant charge $\mathcal{Q}(\alpha, L)$ being the crucial requirement. This can be consistently done, [4], upon recognizing that the celebrated Slavnov–Taylor identities for the couplings fulfill

$$(23) \quad \frac{S_R^{\phi}(X^{\bar{\psi}\gamma \cdot A\psi})}{S_R^{\phi}(X^{\bar{\psi}\gamma \cdot \partial\psi})} = \frac{S_R^{\phi}(X^{AA\partial A})}{S_R^{\phi}(X^{\partial A\partial A})} = \frac{S_R^{\phi}(X^{AAAA})}{S_R^{\phi}(X^{AA\partial A})} = \frac{S_R^{\phi}(X^{\bar{\phi}A \cdot \partial\phi})}{S_R^{\phi}(X^{\bar{\phi}\square\phi})},$$

for the set of amplitudes

$$(24) \quad \mathcal{R}_{\text{QCD}} = \{DADA, \bar{\psi}\gamma \cdot \partial\psi, \bar{\phi}\square\phi, AADA, AAAA, \bar{\phi}A \cdot \partial\phi, \bar{\psi}\gamma \cdot A\psi\},$$

needing renormalization in QCD.

This allows to define a Hochschild cohomology on the sum of graphs at a given loop order, and hence to obtain multiplicative renormalization in this language from the resulting coideals in the Hopf algebra [4, 12].

Note that the structure of the sub-Hopf algebras underlying this approach [5, 8] implies that the elements $X^i(\alpha)$ close under the coproduct. A general classification of related sub-Hopf algebras has been recently obtained by Loic Foissy [13]. He considers only the case that the lowest order Hochschild cocycle is present in the combinatorial Dyson–Schwinger equations, but his study is rather complete when augmented by the results of [2].

3. GRAVITY

We consider pure gravity understood as a theory based on a graviton propagator and n -graviton couplings as vertices. A fuller discussion incorporating ghosts and matter fields is referred to future work.

3.1. Summary of results of [1].

Corollary 1. *Let $|\Gamma| = k$. Then $\omega(\Gamma) = -2(|\Gamma| + 1)$.*

This is a significant change from the behavior of a renormalizable theory: in the renormalizable case, each graph contributing to the same amplitude i has the same powercounting degree regardless of the loop number. Here, we have the dual situation: the loop number determines the powercounting degree, regardless of the amplitude.

Theorem 2. *The set $d_{\omega(\Gamma)}$ contains no primitive element beyond one loop.*

The set $d_{\omega(\Gamma)}$ is determined as a set of dotted graphs, with dots representing $\omega(\Gamma)$ derivatives with respect to masses or external momenta such that the corresponding integrand ι_{Γ} is overall log-divergent. Whilst in a renormalizable theory, we find for

each amplitude in the finite set $\mathcal{R}$ primitives at each loop order in $d_{\omega(\Gamma)}$, here we have an infinite set $\mathcal{R}$, but only one primitive in it.

Proposition 3. *The relations*

$$(25) \quad \frac{X^{n+1}}{X^n} = \frac{X^n}{X^{n-1}}, \quad n \geq 3,$$

define a sub-Hopf algebra with Hochschild closed one-cocycles $B_+^{1,n}$.

Here, X^n is the sum of all graphs with n external graviton lines. One indeed finds that the combinatorial Dyson–Schwinger equations for gravity provide a sub-Hopf algebra upon requiring these relations, in straightforward generalization of the situation in a non-abelian gauge theory.

3.2. Comments.

3.2.1. *Gauss-Bonnet.* The Gauss–Bonnet theorem ensures here, in the form

$$(26) \quad 0 = \int_{\mathbb{M}} \sqrt{g} (R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - 4R_{\alpha\beta} R^{\alpha\beta} + R^2)$$

the vanishing of the one-loop renormalization constants. This does not imply the vanishing of the two-loop renormalization constants as their one-loop subdivergences are off-shell. But it implies that the two-loop counter term has only a first order pole by the scattering type formula, in agreement with the vanishing of $\phi_{\text{off-shell}}(\gamma)\phi_{\text{on-shell}}(\Gamma/\gamma)$. Here, γ , Γ/γ is the decomposition of Γ into one-loop graphs and $\phi_{\text{on/off-shell}}$ denotes suitable Feynman rules.

3.2.2. *Two-loop counterterm.* Also, the universality of the two-loop counterterm suggests that indeed

$$(27) \quad \frac{Z^{\text{gr}_{n+1}}}{Z^{\text{gr}_n}} = \frac{Z^{\text{gr}_n}}{Z^{\text{gr}_{n-1}}}, \quad \text{with } Z^{\text{gr}_n} = S_R^\phi(X^n),$$

holds for off-shell counterterms. In particular, if we compute in a space of constant curvature and conformally reduced gravity which maintains many striking features of asymptotic safe gravity [14, 15], the above identities should hold for suitably defined characters: indeed, in such circumstances we can renormalize using a graviton propagator which is effectively massive with the mass $\sqrt{R/6}$ provided by the constant curvature R , and hence can renormalize at zero external momentum. Using the KLT relations [16], this reduces the above identities to a (cumbersome) combinatorial exercise on one-loop graphs to be worked out in the future.

Continuing this line of thought one expects that a single quantity, the β function of gravity, exhibits short-distance singularities. If this expectation bears out, it certainly is in nice conceptual agreement with the expectation that in theories where gravity has a vanishing β function, gravity is indeed a finite theory [17].

3.2.3. *Other instances of gravity powercounting.* The appearance of Feynman rules such that the powercounting of vertex amplitudes in $\mathcal{R}_V$ cancels the powercounting of propagator amplitudes in $\mathcal{R}_E$, $\mathcal{R} = \mathcal{R}_V \cup \mathcal{R}_E$, is not restricted to gravity. It indeed appears for example also in the field theoretic description of Brownian fluids and glass possibly, which were recently described at tree-level as a field theory [18], with dynamics beyond tree level which will involve renormalization with powercounting properties similar to the present discussion.

REFERENCES

- [1] D. Kreimer, *A remark on quantum gravity*, Annals Phys. **323** (2008) 49 [arXiv:0705.3897 [hep-th]].
- [2] C. Bergbauer and D. Kreimer, *Hopf algebras in renormalization theory: Locality and Dyson-Schwinger equations from Hochschild cohomology*, IRMA Lect. Math. Theor. Phys. **10** (2006) 133 [arXiv:hep-th/0506190].
- [3] D. Kreimer and K. Yeats, *An etude in non-linear Dyson-Schwinger equations*, Nucl. Phys. Proc. Suppl. **160** (2006) 116 [arXiv:hep-th/0605096].
- [4] D. Kreimer, *Anatomy of a gauge theory*, Annals Phys. **321** (2006) 2757 [arXiv:hep-th/0509135].
- [5] D. Kreimer, *Dyson-Schwinger Equations: From Hopf algebras to Number Theory*, in *Universality and Renormalization*, I. Binder, D. Kreimer, eds., Fields Inst Comm. **50** (2007) 225, AMS.
- [6] K. Johnson, M. Baker and R. Willey, *Selfenergy Of The Electron*, Phys. Rev. **136** (1964) B1111.
- [7] M. Baker and K. Johnson, *Asymptotic form of the electron propagator and the selfmass of the electron*, Phys. Rev. D **3** (1971) 2516.
- [8] D. Kreimer and K. Yeats, *Recursion and growth estimates in renormalizable quantum field theory*, Commun. Math. Phys. (2008) DOI:10.1007/s00220-008-0431-7; arXiv:hep-th/0612179.
- [9] D. Kreimer, K. Yeats, G. Van Baalen, D. Uminsky, in preparation, see also Karen Yeats homepage at <http://math.bu.edu/people/kayeats/papers/picturetalkshort4up.pdf>
- [10] S. G. Gorishnii, A. L. Kataev, S. A. Larin and L. R. Surguladze, *The Analytical four loop corrections to the QED Beta function in the MS scheme and to the QED psi function: Total reevaluation*, Phys. Lett. B **256** (1991) 81.
- [11] C. Itzykson, G. Parisi and J. B. Zuber, *Asymptotic Estimates In Quantum Electrodynamics*, Phys. Rev. D **16** (1977) 996;
R. Balian, C. Itzykson, G. Parisi and J. B. Zuber, *Asymptotic Estimates In Quantum Electrodynamics. 2*, Phys. Rev. D **17** (1978) 1041.
- [12] W. D. van Suijlekom, *Renormalization of gauge fields: A Hopf algebra approach*, Commun. Math. Phys. **276** (2007) 773 [arXiv:hep-th/0610137].
- [13] L. Foissy, *Faa di Bruno subalgebras of the Hopf algebra of planar trees from combinatorial Dyson-Schwinger equations*, arXiv:0707.1204 [math.RA].
- [14] M. Reuter, *Nonperturbative Evolution Equation for Quantum Gravity*, Phys. Rev. D **57** (1998) 971 [arXiv:hep-th/9605030];
O. Lauscher and M. Reuter, *Ultraviolet fixed point and generalized flow equation of quantum gravity*, Phys. Rev. D **65** (2002) 025013 [arXiv:hep-th/0108040];
M. Niedermaier, *The asymptotic safety scenario in quantum gravity: An introduction*, arXiv:gr-qc/0610018.
- [15] M. Reuter, H. Weyer, *Background Independence and Asymptotic Safety in Conformally Reduced Gravity*, arXiv:0801.3287 [hep-th].
- [16] Z. Bern, D. C. Dunbar and T. Shimada, *String based methods in perturbative gravity*, Phys. Lett. B **312** (1993) 277 [arXiv:hep-th/9307001];
Z. Bern, L. J. Dixon, D. C. Dunbar, M. Perelstein and J. S. Rozowsky, *On the relationship between Yang-Mills theory and gravity and its implication for ultraviolet divergences*, Nucl. Phys. B **530** (1998) 401 [arXiv:hep-th/9802162];
Z. Bern and A. K. Grant, *Perturbative gravity from QCD amplitudes*, Phys. Lett. B **457** (1999) 23 [arXiv:hep-th/9904026].
- [17] L. Dixon, *Is $N=8$ supergravity finite?*, available via <http://www.slac.stanford.edu/lance/Neq8.ppt>.
- [18] A. Velenich, C. Chamon, L. Cugliandolo, D. Kreimer, *On the Brownian gas: a field theory with a Poissonian ground state*, preprint.

KREIMER@IHES.FR, IHES, 35 RTE. DE CHARTRES, 91440 BURES-SUR-YVETTE, FRANCE ([HTTP://WWW.IHES.FR](http://WWW.IHES.FR)) AND BOSTON U. ([HTTP://MATH.BU.EDU](http://MATH.BU.EDU))